


The impact of forest fragmentation on effective population sizes, mating systems and genetic diversity of forest trees in Guanacaste province, Costa Rica
The role of remote sensing in monitoring tropical forest fragments in Costa Rica

**FINAL REPORT TO THE CENTRE FOR
INTERNATIONAL FORESTRY RESEARCH**

Jonathan Cornelius
Centre for Research and Higher Education in Tropical Agriculture
Turrialba, Costa Rica

2002

SUMMARY



Recent estimates suggest that as much as half of the world's forest area has been cleared or degraded since the beginning of the Holocene, recent losses being particularly serious in the tropics. Complete forest removal leads to loss of those unique genetic characteristics were previously present. However, deforestation, rather than being always complete, often leads to the formation of forest fragments set in an unforested or partially forested matrix. The ultimate impact of deforestation depends on the permanence of such fragments, which in turn depends on the viability of their plant and animal populations. Such viability is not assured, because small populations are subject to a number of pressures which are absent or of less importance in large populations. With population extinction, resident genetic diversity is lost; as inbreeding depression due to depleted genetic variation is one consequence of small population size, reduced genetic diversity is at once both causal agent in, and effect of, population extinction.

Clearly, the impact of forest fragmentation on genetic diversity will partly determine the contribution that fragments can make to management and conservation of biodiversity, *i.e.* whether they can mitigate the many negative effects of deforestation, or whether they simply constitute a transient phase in forest destruction. Neither outcome is likely to apply to all species in all circumstances. Rather, given the diversity of plant and animal life, diversity of response would seem to be the most rational expectation. However, this also implies that certain characteristics and fragmentation scenarios are likely to be associated with the degree of resilience to fragmentation effects. The identification of such characteristics would seem to be the most promising means of arriving at meaningful generalizations on the effect of fragmentation

The project 'The impact of forest fragmentation on effective population sizes, mating systems and genetic diversity of forest trees in Guanacaste province, Costa Rica', addressed this question. Its general objective was to contribute to the effective in situ management and conservation of forest genetic resources. The specific objectives, as detailed in the original project document were (1) estimation of genetic diversity, outcrossing rates, and effective population sizes in a range of fragments of differing characteristics; (2) quantification of the influence of physical characteristics of fragments on genetic processes, including gene flow, drift and mating; (3) construction of a model that would permit the identification of key factors influencing population genetic processes in forest fragments. The complementary project 'The role of remote sensing in monitoring tropical forest fragments in Costa Rica' was approved in 1997 in order to provide funding for aerial photography of the study sites, in order to 'support interpretation of data collected by ground-based methods, specifically, to assess the value of airborne sensing in interpreting the impacts of habitat fragmentation on components of biodiversity, including genetic diversity and reproductive dynamics'.

The research was carried out in a 350km² area of seasonally-dry, lowland Guanacaste province. Chapter One explains the justification of the research, in terms of the importance of genetic diversity and the threat posed by fragmentation. The latter is explored more fully

in Chapter Two, which reviews expected and observed genetic consequences of fragmentation. Reductions in variation, often attributed to founder effects, have frequently been observed. Impacts on pollinators, fertility and mating have potential to exert further downward pressure on effective population sizes, whilst the actual mitigatory efficacy of migration is not clear. The genetic impact of fragmentation depends partly on fragment historical-spatial characteristics, as explored in Chapter Three, which traces the deforestation history of the zone, culminating in an analysis of forest change since the 1940s and a discussion of genetic implications of key findings, particularly evidence of increased forest linearity. Chapters Five and Seven report studies of genetic impacts on two native species, based partly on isozyme analysis (Chapters Four and Six describe inheritance, neutrality and linkage of the markers). Fragment populations of *Anacardium excelsum* (Bertero & Balbis) Skeels (Anacardiaceae) exhibit tendencies which suggest relatively high susceptibility to fragmentation and associated disturbance, particularly density-correlated outcrossing, high within-fragment variation in fertility, and reduced seedling growth rates, whilst several factors indicate gene flow may be distance-limited. *Plumeria rubra* L. (Apocynaceae) appears generally less susceptible. Self-incompatibility precludes outcrossing effects, whilst low population differentiation suggests that migration between populations, presumably facilitated by highly mobile hawkmoth pollinators, has been sufficient to avoid loss of variation. Nevertheless, low variation in one isolated population indicates that this resilience is not unlimited. In Chapter Eight, general implications are discussed, with reference both to characteristics of species and forests of the zone, and to genetic and reproductive processes discussed previously. Species groups at particular genetic risk are identified. The mitigation of fragmentation impacts by improved husbandry of existing pastureland and riparian trees is considered.

TABLE OF CONTENTS

SUMMARY.....	2
TABLE OF CONTENTS	4
LIST OF TABLES	8
LIST OF FIGURES.....	12
LIST OF ABBREVIATIONS.....	16
1. INTRODUCTION.....	17
2. THE GENETIC EFFECTS OF FOREST FRAGMENTATION: AN OVERVIEW OF EXPECTATIONS AND FINDINGS	20
Introduction.....	20
Reduction in population size.....	
Expected genetic consequences of reductions in population size.....	21
Fragmentation and pollination.....	24
Fragmentation and gene flow.....	27
Genetic variation in fragmented plant populations	30
Conclusions.....	34
3. HISTORICAL AND SPATIAL PATTERNS OF DEFORESTATION AND FRAGMENTATION IN GUANACASTE PROVINCE, COSTA RICA: CHARACTERIZATION AND GENETIC IMPLICATIONS.....	35
Introduction.....	35
The study zone	36
General patterns of deforestation in lowland Guanacaste	39
Phase One: feral cattle ranching (1560-to late 19 th century).....	39
Phase Two: Timber exploitation, introduction of exotic pasture and cattle (1880- 1930s)	40
Phase Three: Deforestation in the 1950s and after.....	40
Deforestation in the study zone	41
Phases One and Two.....	41
Quantitative analysis.....	43
Methods	43
Results	44
Discussion.....	45
General discussion	47
4. INHERITANCE, LINKAGE AND NEUTRALITY OF ALLOZYMES OF THE NEOTROPICAL TREE <i>ANACARDIUM EXCELSUM</i> (BERTERO & BALBIS EX KUNTH) SKEELS (ANACARDIACEAE)	53
Introduction.....	53
Materials and Methods	54
Linkage disequilibrium.....	56
Neutrality	57
Results	57
Inheritance.....	57
Adenylate kinase	57
Leucine aminopeptidase	58

Malate dehydrogenase.....	59
Phosphogluconate dehydrogenase.....	59
Phosphoglucomutase.....	60
UTP-glucose-1-phosphate uridylyltransferase	60
Linkage.....	61
Neutrality.....	61
Discussion	62
Inheritance.....	62
Linkage disequilibrium.....	66
Neutrality.....	67
Applications.....	67
5. THE EFFECTS OF FOREST FRAGMENTATION ON GENETICS AND REPRODUCTION OF THE TREE <i>ANACARDIUM EXCELSUM</i> (BERTERO & BALBIS) SKEELS (ANACARDIACEAE) IN NORTHWESTERN COSTA RICA 68	
Introduction.....	68
Methods.....	70
Study zone and fragments.....	70
Genetic variation, gene flow and mating systems	72
<i>Field and laboratory procedures</i>	72
<i>Population genetic analysis</i>	72
Current gene flow.....	73
Estimation of mating system parameters	75
Flowering.....	76
Seed weight and growth rate	78
Results	81
Genetic variation and structure.....	81
Maternal generation.....	81
Progeny generation.....	84
Gene flow.....	87
Mating system parameters.....	87
Flowering.....	88
Common garden experiments.....	90
1997 experiment	90
1999 experiment	91
Discussion	94
Founder effects.....	95
Flowering and seed weight.....	95
Inbreeding	96
Seedling growth rate.....	98
Gene flow	98
Selection.....	100
6. INHERITANCE, LINKAGE AND NEUTRALITY OF ALLOZYMES OF THE NEOTROPICAL TREE <i>PLUMERIA RUBRA</i> L. (APOCYNACEAE).....	102
Introduction.....	102
Materials and Methods	103
Field and laboratory procedures	103
Inheritance.....	104
Linkage disequilibrium.....	105

Neutrality	106
Results	106
Inheritance.....	106
Aspartate aminotransferase.....	106
Alcohol dehydrogenase	107
Glucose-phosphate isomerase.....	108
Phosphoglucomutase (PGM; EC 5.4.2.2).....	109
Phosphogluconate dehydrogenase (PGD, EC 1.1.1.43)	110
Linkage disequilibrium.....	111
Selective neutrality.....	111
Discussion	112
Inheritance.....	112
Linkage disequilibrium.....	112
Neutrality	114
7. THE EFFECTS OF FOREST FRAGMENTATION ON GENETICS AND REPRODUCTION OF THE TREE <i>PLUMERIA RUBRA</i> L. IN NORTHWESTERN COSTA RICA	115
Introduction.....	115
Methods.....	117
Study zone and fragments.....	117
Field and laboratory procedures	119
Specific research questions	120
Parental generation.....	121
Current gene flow.....	123
Mating system	123
Fruit production	124
Results.....	124
Genetic variation	124
Maternal generation.....	124
Progeny: 1997.....	125
Progeny: 1998.....	126
Gene flow	128
Mating system parameters.....	128
Fruit production	129
Discussion	132
Population differentiation.....	132
Within-population genetic variation.....	132
Gene flow.....	135
Mating system parameters.....	136
Fruit production	137
Implications.....	138
8. GENERAL CONCLUSIONS: GENETIC IMPLICATIONS OF FOREST FRAGMENTATION FOR SPECIES OF LOWLAND GUANACASTE PROVINCE	140
Introduction.....	140
Species of the study zone.....	140
Effects on outcrossing	142
High variance in reproductive output.....	143

Founder effects.....	143
Gene flow and random genetic drift.....	144
BIBLIOGRAPHY	148
APPENDICES	170
Appendix I: Tables.....	171
Appendix II: Details of laboratory procedures	237
Anacardium excelsum.....	237
Enzyme extraction	237
Gel preparation.....	237
Gel loading and running.....	238
Staining.....	238
Plumeria rubra.....	240
Enzyme extraction	240
Gel preparation.....	241
Gel loading and running.....	241
Staining.....	241
Appendix III: Modified maternal genotypes of <i>Plumeria rubra</i>	243
Appendix IV: Classified species lists for <i>hacienda La Pacifica</i>	245
Introduction	245
Species by breeding system and self-compatibility	245
Species by pollen and seed dispersal vectors	245
Species density	246
Appendix V: financial reports	262
Appendix VI: Project personnel.....	271

LIST OF TABLES

Table 3-1. Two-way contingency tables of location with respect to nearest watercourse ($\leq 100\text{m}$, $>100\text{m}$) and presence of forest in 1945 (southern sector) and 1945-61 (northern sector)	171
Table 3-2. Two-way contingency tables of location with respect to nearest watercourse ($\leq 100\text{m}$, $>100\text{m}$) and presence of forest, 1998.....	172
Table 3-3. Two-way contingency tables of presence of trees within radius of 27.5m in agricultural and pastureland, 1998	172
Table 4-1. Segregation ratios in pooled progeny arrays of five enzyme systems of <i>Anacardium excelsum</i>	173
Table 4-2. Segregation ratios and associated probabilities in progeny of mother trees of <i>Anacardium excelsum</i> putatively heterozygous for the AK2 locus.....	174
Table 4-3. Segregation ratios and associated probabilities in progeny of mother trees of <i>Anacardium excelsum</i> putatively heterozygous for the LAP locus.....	175
Table 4-4. Segregation ratios and associated probabilities in progeny of mother trees of <i>Anacardium excelsum</i> putatively heterozygous for the PGD locus.	177
Table 4-5. Segregation ratios and associated probabilities in progeny of mother trees of <i>Anacardium excelsum</i> putatively heterozygous for the PGM1 locus.....	178
Table 4-6. Segregation ratios and associated probabilities of putatively heterozygous mother trees of <i>Anacardium excelsum</i> , UTP-glucose-1-phosphate uridylyltransferase ...	179
Table 4-7. Significant estimates of Burrows's composite linkage disequilibrium coefficient, correlation coefficient and significance of associated chi-squared test in populations of <i>Anacardium excelsum</i> in Guanacaste province, Costa Rica.....	180
Table 4-8. Two by three classification of <i>Anacardium excelsum</i> progeny by PGD putative genotype and presence of putative AK1-A allele, with result of chi-square test of association.	180
Table 4-9. Estimates of Ohta's multiple population linkage disequilibrium coefficients for 12 populations of <i>Anacardium excelsum</i> in northwestern Costa Rica.	181
Table 4-10. Results of Ewens-Watterson test of selective neutrality for 12 populations of <i>Anacardium excelsum</i> in northwestern Costa Rica.	182
Table 4-11. Estimates of proportions of extraneous seed in collections of putatively heterozygous mother trees of <i>Anacardium excelsum</i>	183
Table 5-1. Description of populations sampled in a study of genetic effects of forest fragmentation on <i>Anacardium excelsum</i> in northwestern Costa Rica.	185
Table 5-2. Sampling of populations of in a study of effects of forest fragmentation on genetic and reproduction of <i>Anacardium excelsum</i> in northwestern Costa Rica.	187
Table 5-3. Representation by block of seed sources (fragments) and open-pollinated families within fragments in a common garden experiment of <i>Anacardium excelsum</i>	188
Table 5-4. Estimates of allele frequencies, allelic richness, expected heterozygosity, fixation indices and G-test for Hardy-Weinberg disequilibria in populations of <i>Anacardium excelsum</i> from 10 forest fragments in northwestern Costa Rica, based on inferred maternal genotypes.....	190
Table 5-5. Homogeneity statistics for five loci of inferred maternal genotypes in populations of <i>Anacardium excelsum</i> from 10 forest fragments in northwestern Costa Rica	192

Table 5-6. Estimates of Wright's statistics and mN for five loci of inferred maternal genotypes in populations of <i>Anacardium excelsum</i> from 10 forest fragments in northwestern Costa Rica.....	193
Table 5-7. Pairwise estimates of Nei's unbiased genetic distance (below diagonal) and geographic distances (km, above diagonal) between ten populations of <i>Anacardium excelsum</i> located in northwestern Costa Rica, based on inferred maternal genotypes...	194
Table 5-8. Estimates of allele frequencies, allelic richness, expected heterozygosity, fixation indices and G-test for Hardy-Weinberg disequilibria in populations of <i>Anacardium excelsum</i> from 12 forest fragments in northwestern Costa Rica (progeny data).....	195
Table 5-9. Homogeneity statistics for five loci of progeny genotypes in populations of <i>Anacardium excelsum</i> from 12 forest fragments in northwestern Costa Rica.....	198
Table 5-10. Estimates of Wright's statistics and mN for five loci of progeny genotypes in populations of <i>Anacardium excelsum</i> from 10 forest fragments in northwestern Costa Rica.	199
Table 5-11. Pairwise estimates of Nei's unbiased genetic distance (below diagonal) and geographic distances (km, above diagonal) between ten populations of <i>Anacardium excelsum</i> located in northwestern Costa Rica	200
Table 5-12. Estimates of current (1999) gene flow into the Paso Hondo population based on two alleles absent from the Paso Hondo population.....	200
Table 5-13. Estimates of mating system parameters of <i>A. excelsum</i> in populations located in 10 forest fragments in northwestern Costa Rica.....	201
Table 5-14. Mean population flowering indices and associated data in 24 forest fragments of <i>Anacardium excelsum</i> in northwestern Costa Rica.	202
Table 5-15. Linear regression of fragment mean tree flowering index on fragment type and mean dbh of sampled trees in 24 fragment populations of <i>A. excelsum</i> in northwestern Costa Rica	202
Table 5-16. Linear regressions of flowering equitability on fragment type and mean dbh and equitability on type and dbh standard deviation of sampled trees in 24 fragment populations of <i>A. excelsum</i> in northwestern Costa Rica.	203
Table 5-17. Analysis of variance of effect of origin (fragment) on weight of seeds collected from seven populations of <i>Anacardium excelsum</i> located in northwestern Costa Rica.	204
Table 5-19. Analysis of variance of effect of origin (fragment) on arcsin germination percentage, four-month height and number of days to germination of seedlings from seven populations of <i>Anacardium excelsum</i> located in northwestern Costa Rica.	204
Table 5-20. Germination percentage, least square means for height at four months (cm) and number of days to germination for <i>Anacardium excelsum</i> seedlings from seven forest fragments in northwestern Costa Rica	205
Table 5-21. Analysis of covariance of effect of origin (fragment), seed weight and germination day on four-month of seedlings from seven populations of <i>Anacardium excelsum</i> located in northwestern Costa Rica.	206
Table 5-22. Analysis of variance of a greenhouse experiment comparing growth of <i>Anacardium excelsum</i> seedlings from 9 forest fragments in northwestern Costa Rica	207
Table 5-23. Least square means of total height at 81 and 167 days and diameter 2cm above root collar at 167 days of <i>Anacardium excelsum</i> seedlings from 9 forest fragments in northwestern Costa Rica.....	208
Table 5-24. Analysis of covariance of a greenhouse experiment comparing growth of <i>Anacardium excelsum</i> seedlings from 9 forest fragments in northwestern Costa Rica	208

Table 5-25. Least square means of total height at 81 and 167 days and diameter 2cm above root collar at 167 days of <i>Anacardium excelsum</i> seedlings from 9 forest fragments in northwestern Costa Rica (based on analysis of covariance).....	209
Table 5-26. Numbers of albino and normal seedlings in progeny of three trees of <i>Anacardium excelsum</i> from populations located in forest fragments in northwestern Costa Rica	209
Table 6-1. Segregation ratios in pooled progeny arrays of five loci of <i>Plumeria rubra</i> .	210
Table 6-2. Segregation ratios and associated probabilities in progeny of mother trees of <i>Plumeria rubra</i> putatively heterozygous for the AAT1 locus.....	211
Table 6-3. Segregation ratios and associated probabilities in progeny of mother trees of <i>Plumeria rubra</i> putatively heterozygous for ADH3 alleles A and B.	213
Table 6-4. Segregation ratios and associated probabilities in progeny of mother trees of <i>Plumeria rubra</i> putatively heterozygous for locus PGI3.....	213
Table 6-5. Segregation ratios and associated probabilities in progeny of mother trees of <i>Plumeria rubra</i> putatively heterozygous for the PGM1 locus.....	214
Table 6-6. Segregation ratios and associated probabilities in progeny of mother trees of <i>Plumeria rubra</i> putatively heterozygous for locus PGD1.....	215
Table 6-7. Significant estimates of Burrows's composite linkage disequilibrium coefficient, correlation coefficient and significance of associated chi-squared test in 7 populations of <i>Plumeria rubra</i> in Guanacaste province, Costa Rica.	217
Table 6-8. Estimates of Ohta's multiple population linkage disequilibrium coefficients for 7 populations of <i>Plumeria rubra</i> in northwestern Costa Rica.....	218
Table 7-1. Description of populations sampled in a study of the genetic effects of forest fragmentation on <i>Plumeria rubra</i> in northwestern Costa Rica.....	219
Table 7-2. Sampling schedule in a study of genetic effects of forest fragmentation in <i>Plumeria rubra</i> in northwestern Costa Rica.....	221
Table 7-3. Estimates of allele frequencies, number of alleles, expected heterozygosity, fixation indices and G-test for Hardy-Weinberg disequilibria in populations of <i>P. rubra</i> from 7 forest fragments in northwestern Costa Rica, based on inferred maternal genotypes.	223
Table 7-4. Homogeneity statistics for five loci of inferred maternal genotypes seven populations in a study of effects of forest fragmentation on genetics of <i>Plumeria rubra</i> in northwestern Costa Rica.....	224
Table 7-5. Estimates of Wright's statistics and <i>mN</i> between seven populations (inferred maternal genotypes) (estimates in parentheses based on data set without Pachanga and Sandillal populations).....	225
Table 7-6. Pairwise estimates of Nei's unbiased genetic distance (below diagonal) and geographic distances (km, above diagonal) between seven populations of <i>Plumeria rubra</i> located in Guanacaste province, northwestern Costa Rica.....	226
Table 7-7. Estimates of allele frequencies, number of alleles, expected heterozygosity, fixation indices and G-test for Hardy-Weinberg disequilibria in 1997 progeny of four populations of <i>P. rubra</i> in northwestern Costa Rica.....	226
Table 7-8. Homogeneity statistics for six loci of four <i>Plumeria rubra</i> populations in a study of effects of forest fragmentation in northwestern Costa Rica (1997 progeny) ..	227
Table 7-9. Estimates of Wright's statistics and <i>mN</i> between 1997 progeny of four populations of <i>P. rubra</i> located in Guanacaste province, northwestern Costa Rica.	227
Table 7-10. Pairwise estimates of Nei's unbiased genetic distance (below diagonal) and geographic distances (km, above diagonal) between 1997 progeny of four populations of <i>Plumeria rubra</i> located in Guanacaste province, northwestern Costa Rica.....	227

Table 7-11. Estimates of allele frequencies, number of alleles, expected heterozygosity, fixation indices and G-test for Hardy-Weinberg disequilibria in 1998 progeny of six populations of <i>P. rubra</i> in northwestern Costa Rica.....	228
Table 7-12. Estimates of Wright's statistics and mN between 1998 progeny of six populations of <i>P. rubra</i> located in Guanacaste province, northwestern Costa Rica.	229
Table 7-13. Pairwise estimates of Nei's unbiased genetic distance (below diagonal) and geographic distances (km) between six populations of <i>P. rubra</i> located in Guanacaste province, Costa Rica.....	230
Table 7-14. Estimates of allele frequencies, number of alleles, expected heterozygosity, fixation indices and G-test for Hardy-Weinberg disequilibria in 1999 progeny of five populations of <i>P. rubra</i> in northwestern Costa Rica.....	231
Table 7-15. Estimates of Wright's statistics and mN between 1999 progeny of five populations.....	232
Table 7-18. Estimates of mating system parameters in seven population of <i>Plumeria rubra</i> located in forest fragments in northwestern Costa Rica.....	233
Table 7-19. Linear regression of log of number of capsules tree ⁻¹ on dbh and neighbourhood density index of trees of <i>Plumeria rubra</i> located in six forest fragments in northwestern Costa Rica.....	234
Table 7-20. Mean numbers of capsules tree ⁻¹ in trees of <i>Plumeria rubra</i> located in seven forest fragments in northwestern Costa Rica	234
Table 7-21. Analysis of covariance of effects of fragment and dbh on mean of log numbers of capsules tree ⁻¹ in six forest fragments of <i>Plumeria rubra</i> located in northwestern Costa Rica.....	234
Table 8-1. Summary of breeding systems and self-compatibility of 114 tree species of Hacienda La Pacífica, Cañas, Guanacaste province, Costa Rica.....	235
Table 8-2. Summary pollen and diaspore vectors of 114 tree species of Hacienda La Pacífica, Cañas, Guanacaste province, Costa Rica.....	235
Table 8-3. Two by two classification of pollen and diaspore dispersal vectors of x forest tree species in Cañas, Guanacaste province, Costa Rica.....	236
Table A3-1. Reassigned maternal inferred genotypes of <i>Plumeria rubra</i>	243
Table A4-1. Tree species of Hacienda La Pacífica, Guanacaste province, Costa Rica, classified according to breeding system and self-compatibility.....	247
Table A4-2. Tree species of Hacienda La Pacífica, Cañas, Guanacaste, Costa Rica, classified by pollinator type and diaspore dispersal vector.....	252
Table A4-3. Population densities of tree species of Hacienda La Pacífica, Cañas, Guanacaste, Costa Rica (data based on Glander and Nisbett, 1996; Jiménez et al., 1987)..	258

LIST OF FIGURES

- Figure 3-1.** Costa Rica: major cities, relief and location of the study zone..... 37
- Figure 3-2.** The study zone and surrounding localities (scale: 1:100000) (source: IGN 1988a, 1988b, reproduced by permission)..... 38
- Figure 3-3.** Land use in the study zone, 1945-1998: estimated percentage cover (standard error) by use category..... 45
- Figure 3-4.** Las Lomas and Llanos de San Pedro area, Hacienda Taboga, in 1956 (left) and 1986 (right)..... 46
- Figure 4-1.** Adenylate kinase zymogram of *Anacardium excelsum* (3 families), showing putative loci (marked '1' and '2'). 'C' indicates *Pinus resinosa* control..... 58
- Figure 4-2.** Adenylate kinase zymogram of *Anacardium excelsum* (2 families). 'X' indicates lane with most common AK2 allele absent and intense staining cathodal of most common AK1 band (see text). 'C' indicates *Pinus resinosa* control..... 58
- Figure 4-3.** Leucine aminopeptidase zymogram of *Anacardium excelsum*, showing putative genotypes AA (marked '1'), BB (2), AB (3), CC (4), AC (5) in four arrays, and *Pinus resinosa* controls (C)..... 58
- Figure 4-4.** Malate dehydrogenase zymogram of *Anacardium excelsum*, showing variant phenotypes (marked V) in two progeny arrays and *Pinus resinosa* controls (C). 59
- Figure 4-5.** Phosphogluconate dehydrogenase zymogram of *Anacardium excelsum*, showing putative genotypes AA (marked '1'), BB (2), AB (3) in two progeny arrays, and *Pinus resinosa* controls (C)..... 59
- Figure 4-6.** Phosphoglucomutase zymogram of *A. excelsum*, showing putative genotypes AA (marked '1'), BB (2), AB (3) in three arrays, faint artefactual banding in PGM-B position (BB genotypes marked '2' and those adjacent (right)) and *Pinus resinosa* controls (C) 60
- Figure 4-7.** UTP-glucose-1-phosphate uridylyltransferase zymogram of *Anacardium excelsum*, showing putative genotypes AA (marked '1'), BB (2), AB (3) in two progeny arrays, and *Pinus resinosa* controls (C). 61
- Figure 4-8.** Adenylate kinase (AK) (above) and phosphogluconate dehydrogenase (PGD) (below) zymograms for *A. excelsum* (family Marcela 418), showing apparent linkage (putative PGD genotypes: 1=AA, 2=AB, 3=BB. C indicates *Pinus resinosa* control..... 62
- Figure 5-1.** The study zone and populations in a study of genetic effects of forest fragmentation on *Anacardium excelsum* populations located near Cañas, Guanacaste province, Costa Rica. Top: a. northern sector; bottom: b. southern sector 71
- Figure 5-2.** Location of Bosque Duquesa tree 706 in relation to other sampled trees (scale approximately 1:7400) (photo: Aerofotos de Costa Rica S.A.)..... 72
- Figure 5-3.** Scatter plot of log population size (log N) and gene diversity of adult trees in ten populations of *Anacardium excelsum* located in forest fragments in northwestern Costa Rica..... 82
- Figure 5-4.** Scatter plot of pairwise relationship between geographic and Nei's unbiased genetic distances in 10 populations of *Anacardium excelsum* located in northwestern Costa Rica (based on inferred maternal genotypes) 83
- Figure 5-5.** UPGMA dendrogram based on Nei's unbiased genetic distance between maternal trees in ten populations of *A. excelsum* located in northwestern Costa Rica. 83

Figure 5-6. UPGMA dendrogram based on Nei's unbiased genetic distance between maternal trees in four populations of <i>A. excelsum</i> (Corobici group) located in northwestern Costa Rica.....	84
Figure 5-7. UPGMA dendrogram based on Nei's unbiased genetic distance between maternal trees in three populations of <i>A. excelsum</i> (Taboga group) located in northwestern Costa Rica.....	84
Figure 5-8. Scatter plot of log population size (log N) and gene diversity of progeny in twelve populations of <i>Anacardium excelsum</i> located in forest fragments in northwestern Costa Rica.....	85
Figure 5-9. Scatter plot of pairwise relationship between geographic and Nei's unbiased genetic distances in 10 populations of <i>Anacardium excelsum</i> located in northwestern Costa Rica (progeny genotypes).....	86
Figure 5-10. UPGMA dendrogram based on Nei's unbiased genetic distance between progeny in 12 populations of <i>Anacardium excelsum</i> located in northwestern Costa Rica.	86
Figure 5-11. UPGMA dendrogram based on Nei's unbiased genetic distance between progeny in four populations of <i>A. excelsum</i> (Corobici group) located in northwestern Costa Rica.....	87
Figure 5-12. UPGMA dendrogram based on Nei's unbiased genetic distance between progeny in three populations of <i>Anacardium excelsum</i> (Taboga group) located in northwestern Costa Rica.....	87
Figure 5-13. Scatter plot of observed inbreeding coefficients (F_{is}) and equilibrium inbreeding coefficients (F_e) based on estimated outcrossing rates ($\hat{F}_e = (1-t_m)/(1+t_m)$) (one-tailed) in populations of <i>Anacardium excelsum</i> located in 10 forest fragments in northwestern Costa Rica (p-value is one-tailed).....	88
Figure 5-14. Linear regression of multilocus population outcrossing rates on population mean neighborhood density index (see text) in ten populations of <i>Anacardium excelsum</i> located in northwestern Costa Rica.....	89
Figure 5-15. Scatter plot of flowering equitability and mean flowering index in 24 populations of <i>Anacardium excelsum</i> located in forest fragments in northwestern Costa Rica.	90
Figure 5-16 Scatter plot of least squares mean height at 81 days (based on ANOVA) and estimated outcrossing rate in populations of <i>Anacardium excelsum</i> from nine forest fragments in northwestern Costa Rica (p-value is one-tailed).....	91
Figure 5-17. Scatter plot of least square mean height at 167 days (based on ANOVA)and estimated outcrossing rate in populations of <i>Anacardium excelsum</i> from nine forest fragments in northwestern Costa Rica (p-value is one-tailed).....	91
Figure 5-18. Scatter plot of least square mean diameter at 167 days (based on ANOVA) and estimated outcrossing rate in populations of <i>Anacardium excelsum</i> from nine forest fragments in northwestern Costa Rica (p-value is one-tailed).....	92
Figure 5-19. Scatter plot of least square mean height at 81 days (based on ANCOVA) and estimated outcrossing rate in populations of <i>Anacardium excelsum</i> from nine forest fragments in northwestern Costa Rica (p-value is one-tailed).....	92
Figure 5-20. Scatter plot of least square mean height at 167 days (based on ANCOVA) and estimated outcrossing rate in populations of <i>Anacardium excelsum</i> from nine forest fragments in northwestern Costa Rica (p-value is one-tailed).....	93

- Figure 5-21.** Scatter plot of least square mean diameter at 167 days (based on ANCOVA) and estimated outcrossing rate in populations of *Anacardium excelsum* from nine forest fragments in northwestern Costa Rica (p-value is one-tailed) 93
- Figure 5-22.** Albino progeny of *Anacardium excelsum* tree 21, El Ojoche population, northwestern Costa Rica..... 94
- Figure 6-1.** Aspartate aminotransferase zymogram of *Plumeria rubra*, showing (1) faint banding in A-locus (most cathodal) position in BC and BB genotypes (marked respectively 1 and 2); C indicates the control (*Pinus resinosa*). 106
- Figure 6-2.** Aspartate aminotransferase zymogram of *Plumeria rubra*, showing asymmetry of AC heterodimers in two families. 'C' indicates the control (*Pinus resinosa*)..... 107
- Figure 6-3.** Alcohol dehydrogenase zymogram of *Plumeria rubra*. ADH2 (intermediate mobility) shows an apparently monomeric banding pattern and apparent close linkage with the dimeric, most anodal locus. 'C' indicates the control (*Pinus resinosa*). 107
- Figure 6-4.** Example of less complex glucose-phosphate isomerase zymogram of *Plumeria rubra*, showing four zones of activity (putative loci, labelled 1 to 4). 'C' indicates the control (*Pinus resinosa*). 108
- Figure 6-5.** Glucose-phosphate isomerase zymogram of *Plumeria rubra*, showing putative genotypes of locus 3: 1=AA, 2=AB, 3=BB; C indicates control (*Pinus resinosa*).109
- Figure 6-6.** Glucose-phosphate isomerase zymogram of *Plumeria rubra*, showing putative genotypes of locus 3: 2=AC, 1=AA; C indicates control (*Pinus resinosa*); families 2026 (first lane), 2034 (final two lanes), 2027 (other lanes). 109
- Figure 6-7.** Glucose-phosphate isomerase zymogram of *Plumeria rubra*, showing putative genotypes of locus 3: 1=AA; 2=AC; 3=CC. C indicates control (*Pinus resinosa*). Family QP12. 109
- Figure 6-8.** Phosphoglucosmutase zymogram of *Plumeria rubra*, showing putative genotypes at locus PGM1: AA (marked 1), AB (marked 4). 110
- Figure 6-9.** Phosphoglucosmutase zymogram of *Plumeria rubra*, showing putative genotypes at locus PGM1: AA (marked 1), CC (marked 2, AC (marked 3). 110
- Figure 6-10.** Phosphogluconate dehydrogenase zymogram of *Plumeria rubra*, showing putative genotypes: 1=AA, 2=BB, 3=AB, 4=double-banded phenotypes scored as AB (see text); 'C' indicates control (*Pinus resinosa*). 111
- Figure 6-11.** Phosphogluconate dehydrogenase zymogram of *Plumeria rubra*, showing putative genotypes: 1=AA, 2=BB, 3=AB, 4=double-banded phenotypes scored as AB (see text) (family 1004); 'C' indicates control (*Pinus resinosa*). 111
- Figure 7-1.** Study zone and location of populations in a study of the effects of forest fragmentation on *Plumeria rubra* in northwestern Costa Rica..... 118
- Figure 7-2.** UPGMA dendrogram based on Nei's unbiased genetic distance between maternal trees in seven population of *P. rubra* located in northwestern Costa Rica. 125
- Figure 7-3.** UPGMA dendrogram based on Nei's unbiased genetic distance between four population of *P. rubra* located in northwestern Costa Rica (1997 progeny) 126
- Figure 7-4.** UPGMA dendrogram based on Nei's unbiased genetic distance between four population of *P. rubra* located in northwestern Costa Rica (1998 progeny) 127
- Figure 7-5.** UPGMA dendrogram based on Nei's unbiased genetic distance between four population of *P. rubra* located in northwestern Costa Rica (1999 progeny) 128
- Figure 7-6.** Frequency distributions of estimates of individual-tree outcrossing rates in seven populations of *Plumeria rubra* located in northwestern Costa Rica..... 129
- Figure 7-6.** Frequency distributions of estimates of individual-tree outcrossing rates in seven populations of *Plumeria rubra* located in northwestern Costa Rica (continued) 129

- Figure 7-7.** Scatter plot of relationship between neighbourhood density index and individual-tree outcrossing rate in trees of *Plumeria rubra* from 6 forest fragments in northwestern Costa Rica 131
- Figure 7-8.** Scatter plots of log number of capsules tree-1 against dbh and neighbourhood density index rate in trees of *Plumeria rubra* from 6 forest fragments in northwestern Costa Rica 131

LIST OF ABBREVIATIONS

A	allelic richness
AAT	aspartate aminotransferase
ADH	alcohol dehydrogenase
AP	allelic richness of polymorphic loci
a.s.l.	above sea level
A_e	effective allelic richness
AK	adenylate kinase
d.f.	degrees of freedom
E.C.	enzyme commission
F_e	estimated equilibrium value of inbreeding coefficient for a given outcrossing rates
F_{is}	correlation between uniting gametes relative to the subpopulations
F_{it}	correlation between uniting gametes relative to the population as a whole
F_{st}	correlation between two gametes drawn randomly within subpopulations relative to that of two gametes drawn randomly from the population as a whole
H_e	expected heterozygosity, gene diversity
hr	hour
IBD	isolation by distance
J	Shannon's equitability index
LAP	leucine aminopeptidase
m	migration rate
MDH	malate dehydrogenase
N	(1)census population size; (2)north
NDI	neighbourhood density index
N_e	effective population size
OMPG	one migrant per generation
p	(1)allele frequency; (2)probability
P	proportion of loci that are polymorphic
PGD	phosphogluconate dehydrogenase
PGI	glucose-phosphate isomerase
PGM	phosphoglucomutase
pop.	population
q	allele frequency;
r	(1)allele frequency; (2)Pearson's correlation coefficient
R^2	coefficient of determination
s	(1)standard deviation; (2)allele frequency;
S.E.	standard error
UGPU	UTP-glucose-1-phosphate uridylyltransferase
W	west

1. INTRODUCTION

Selection between genetic variants is the logical and practical base of the domestication and genetic improvement of alimentary animals and plants. Consequently, genetic diversity, without which effective selection is impossible, has long been, and remains, basic to human welfare. However, genetic diversity also has a wider importance: the lack of breeding potential and the susceptibility to pests and diseases common in many genetically depauperate domesticates (Smith *et al.*, 1992) are paralleled in depauperate wild species by reduced evolutionary potential and fitness, implying that genetically impoverished populations and species are likely to be more susceptible to extinction (Brook *et al.*, 2002; Frankham, 1998). This connection of genetic diversity to species and, ultimately, community and ecosystem persistence, implies that the value of genetic diversity also must include the wide range of values associated with biodiversity in general, i.e. those encapsulated in the Convention on Biological Diversity (CBD, 1992) as 'intrinsic... ecological, genetic, social, economic, scientific, educational, cultural, recreational and aesthetic values'.

Much of the world's biodiversity is lodged in forests, particularly tropical forests, which are estimated to contain up to 70 per cent of the world's species (Groom, 1994). However, over the last two thousand years or so, the exponential growth of human populations, coupled with the growth of cities, of industrialization, and with the requirements of domesticated animals and plants, has led to widespread destruction and degradation of forested and other natural ecosystems. Approximately half of the world's forest area has been cleared or degraded since the beginning of the Holocene (Groombridge and Jenkins, 2002), with about 30 per cent of the world's land area currently under forest (FAO, 2001). In recent decades, loss has been particularly severe in the tropics, where, in the ten years to 2000 alone, the area of natural forest declined from 1,945 million ha to 1,803 million ha (FAO, 2001).

Complete forest removal leads to loss of whatever unique genetic characteristics were previously present and, concomitantly, contributes to the loss of the set of biodiversity values mentioned above. However, the threat to genetic diversity posed by deforestation may be both more insidious and more pervasive than suggested by the above estimates.

Recent studies suggest that only about one-third of remaining world forest is unfragmented (Riitters *et al.* 2000). The ultimate impact of deforestation depends on the permanence of such fragments, which in turn depends on the viability of their plant and animal populations. Such viability is not assured, because small populations are subject to a number of pressures which are absent or of less importance in large populations. When small population effects such as genetic and demographic stochasticity, Allee effects and inherent susceptibility to catastrophe combine (Lacy, 1993), an 'extinction vortex' (Gilpin and Soulé, 1986) may arise, whereby population numbers decline from generation to generation, partly because in any given generation population sizes depend partly on those in that immediately preceding it. With population extinction, resident genetic diversity is lost; as inbreeding depression due to depleted genetic variation is one consequence of small population size, reduced genetic diversity is at once both causal agent in, and effect of, population extinction.

Clearly, the impact of forest fragmentation on genetic diversity will partly determine the contribution that fragments can make to management and conservation of biodiversity, *i.e.* whether they can mitigate the many negative effects of deforestation, or whether they simply constitute a transient phase in forest destruction. Neither outcome is likely to apply to all species in all circumstances. Rather, given the diversity of plant and animal life, diversity of response would seem to be the most rational expectation. However, this also implies that certain characteristics and fragmentation scenarios are likely to be associated with the degree of resilience to fragmentation effects. The identification of such characteristics would seem to be the most promising means of arriving at meaningful generalizations on the effect of fragmentation

The project 'The impact of forest fragmentation on effective population sizes, mating systems and genetic diversity of forest trees in Guanacaste province, Costa Rica', addressed this question. Its general objective was to contribute to the effective in situ management and conservation of forest genetic resources. The specific objectives, as detailed in the original project document were:

- estimation of genetic diversity, outcrossing rates, and effective population sizes in a range of fragments of differing characteristics;

- quantification of the influence of physical characteristics of fragments on genetic processes, including gene flow, drift and mating;
- construction of a model that would permit the identification of key factors influencing population genetic processes in forest fragments.

The complementary project 'The role of remote sensing in monitoring tropical forest fragments in Costa Rica' was approved in 1997 in order to provide funding for aerial photography of the study sites, in order to 'support interpretation of data collected by ground-based methods, specifically, to assess the value of airborne sensing in interpreting the impacts of habitat fragmentation on components of biodiversity, including genetic diversity and reproductive dynamics'.

The present document reports the findings of these Projects in eight chapters. In Chapter Two, the context of the research is clarified by reviewing both the expected genetic consequences of forest fragmentation and results of studies to date. In Chapter Three, temporal and spatial patterns of forest removal in the study zone, together with their expected genetic consequences, are described. Subsequently, in Chapters Five and Seven, research on the effect of forest fragmentation on genetic and reproduction of two tree species of the zone, *Anacardium excelsum* (Anacardiaceae) and *Plumeria rubra* (Apocynaceae) is described; the preceding Chapters Four and Six describe the inheritance, linkage relationships and selective neutrality of the allozyme markers used in the studies. In the final chapter, the wider genetic implications of forest fragmentation in lowland Guanacaste are considered, taking into account both the results detailed in Chapters Three, Five and Seven, and the characteristics of tree species of the study zone.

Previous, interim reports were filed on 15th February 2000, 30th March 1998 and 30th June 1997. The final financial reports are included as Appendix V. Project personnel are listed in Appendix VI.

2. THE GENETIC EFFECTS OF FOREST FRAGMENTATION: AN OVERVIEW OF EXPECTATIONS AND FINDINGS

INTRODUCTION

High rates of deforestation prevail in many tropical countries (FAO, 2001). However, in many cases, one or more tracts within formerly continuous tree cover remain forested (Riitters *et al.*, 2000; Schelhas, 1996). Deforestation converts such tracts to fragments set in an unforested matrix. If viable, communities and populations in such fragments have the potential to substantially mitigate the impact of deforestation on biodiversity.

The viability of any population is partly dependent on the maintenance of genetic variation, as the magnitude of the latter may be correlated with both individual and population fitness (Crmokrak and Roff, 1999; Frankham and Ralls, 1998). In fragmented populations, genetic variation and, therefore, viability is threatened due to pressures on key parameters and processes such as population sizes, pollinator behaviour, mating systems and gene flow. In the first part of the present review, these pressures and the mechanisms by which they are expected to affect genetic diversity are described. Subsequently, the results of field studies of effects of fragmentation on genetic diversity are reviewed in the context of these expectations.

REDUCTION IN POPULATION SIZE

For many species, forest fragmentation is likely to lead to reduction in population sizes. There are at least three ways in which this can occur. Firstly, population members may be eliminated by the forest removal that gives rise to fragmentation. When migration, *i.e.* pollen and seed movement, is maintained between fragments, such population reduction is defined simply by the proportion of the population formerly located in the now deforested matrix. By contrast, total interruption of gene flow leads to a second, more severe, type of population reduction, in which a large, continuous population is reduced to a set of small populations, these corresponding to the fragments. Thirdly, in both cases, initial population reduction may quickly be reinforced by pressures resulting from the profound

environmental changes often wrought by fragmentation. Essentially, these occur because, when a tract in continuous forest becomes a fragment, there is necessarily a change in the adjoining habitat(s): it is logical to expect that the newly conjoined habitats will cause changes in each other, particularly if they differ markedly. There is growing evidence on the importance of the effects of unforested matrices on forested fragments, to such a degree that Gascon and Lovejoy (1998) remark that 'recent knowledge about...how the tropical rainforest ecosystem is affected by landscape fragmentation suggests that much of the ecological degradation can be accounted for by the influence of edge effects and the surrounding matrix'. Laurance *et al.* (1997) describe how such effects are strong enough to lead to biomass reductions of up to 36 per cent within 100m of fragment edges, in spite of increased biomass of lianas. Although such effects may increase population size of 'gap species' (e.g. Sizer and Tanner, 1999; Cunningham, 2000), they imply degradation of habitat of later successional species. For example, Jules (1998) found 'almost no recruitment' to *Trillium* populations within 65m of fragment edges in the Siskeyou Mountains, Oregon (interior recruitment was higher). Similarly, Mesquita *et al.* (1999) found a highly significant relationship between tree mortality and distance to edge in Amazonian forest fragments, whilst Gigord *et al.* (1999) found reduced numbers of juveniles in small populations of the Madagascan endemic *Dombeya acutangula* ssp. *acutangula*, due to increased competition from invasives.

Expected genetic consequences of reductions in population size

In finite populations, formation of successive generations tends necessarily to involve an element of sampling, because the number of gametes and/or zygotes is generally much greater than the adult population to which these give rise. Although viability selection may influence the genetic composition of the survivors, in general random factors will determine survival before natural selection is able to act. As with any sample, there is no guarantee that it will accurately reflect the genetic constitution of the population (in statistical and biological sense) from which it has been drawn. Consequently, unless gametes are genetically invariable, there will be random fluctuations in allele and genotype frequencies from generation to generation, *i.e.* random genetic drift. Eventually, random genetic drift leads to fixation and loss of alleles; fixation probability of each allele is equal to its initial frequency (Hartl and Clark, 1989), whilst expected time to loss depends both on this parameter and

the effective population size (N_e), *i.e.* the number of individuals in a hypothetical ‘ideal’ population with the same magnitude of random drift as the actual population (Hartl and Clark, 1989, Savolainen and Kuittinen, 2000). For example, for $p=0.5$, expected time to fixation is $2.8N_e$ generations, whilst for $p=0.1$ it is about $1.3N_e$ generations (Yeh, 2000). In the interim, allele frequencies tend to become more extreme. As a consequence, heterozygosity is expected to decrease in each generation by a proportion of $1/2N_e$, such that in generation t , expected value of gene diversity is:

$$H_{a(t)} = \left(1 - \frac{1}{2N_e}\right)^t H_{a(0)},$$

where $H_{a(0)}$ is initial gene diversity (Hartl and Clark, 1989). Additive genetic variance is expected to decline at the same rate (Lande and Barrowclough, 1987, Savolainen and Kuittinen, 2000). Otherwise, however, the course of drift in an individual population is impossible to predict. For example, if the least common allele increases in frequency, expected heterozygosity will show a transient increase, although eventually a chance sequence of events is expected to lead to fixation and therefore loss of all heterozygosity.

In practice, and for two reasons, complete fixation is unlikely to occur. Firstly, mutation may reintroduce lost alleles, although, given the low frequency of mutation events, for small population sizes equilibrium values of heterozygosity in closed finite populations will still be close to zero. Secondly, analogously, and more potently, migration can also introduce alleles. This second factor is of greater importance: both generally, given that migration rates are generally several orders of magnitude higher than mutation rates (Hart and Clark, 1989), and in the present context, because the genetic effect of forest fragmentation has been seen as depending on the interaction of genetic drift (reducing variation) and migration (introducing variation). This interaction is further discussed below (‘Fragmentation and gene flow’).

Post-fragmentation decline in population numbers further increases susceptibility to drift. However, such effects do not derive only from direct reductions of population size, because the magnitude of genetic drift can be predicted as a simple function of population size only when the population characteristics meet the assumptions of the Fisher-Wright drift model

(Caballero, 1994). Usually, this is not the case (Hartl and Clark, 1989), and consequently N_e tends strongly to be smaller than N (Frankham, 1995). Several factors are responsible for this tendency, including differing numbers of females and males, non-Poisson distribution of fecundity and population size fluctuations (Falconer, 1989; Futuyma, 1986; Hartl and Clark, 1989; Lande and Barrowclough, 1987; Nunney, 1993; Wright, 1938; Yeh, 2000). Of these, the most important is generally considered to be variance in fecundity (Nunney, 1993; Falconer, 1989). There are good reasons to expect fragmentation to affect this ratio, principally by introducing more highly variable environmental conditions, typified by the edge-interior distinction: just as fragmentation may increase or decrease fertility, so it can be expected to increase variation in fertility, because the environmental changes which lead to the increases or decreases may not be uniform within fragments. Aldrich and Hamrick's (1998) work illustrate this effect. They found that reproduction of *Symphonia globulifera* (Clusiaceae) in a 38.5ha circular plot was dominated by a numerically small group of remnant pastureland trees, which experienced a post-fragmentation increase in fecundity, leading to a 'secondary constriction' of the fragmentation bottleneck. Similarly, in the New Zealand mistletoe *Peraxilla tetrapetala* (Loranthaceae) pollination and seed set were more than fourfold higher in isolated than continuous-forest individuals (Kelly *et al.*, 2000).

When fragmentation leads to immediate reductions in population size, 'one-time drift' or 'founder effects' (Yeh, 2000) may occur, because remaining trees constitute a sample of the original population. The expected average number of alleles remaining after a founder event is:

$$E = m - \sum_j (1 - p_j)^{2N} ,$$

where m = the number of alleles before population reduction, p_j = frequency of the j^{th} allele, N = population size after reduction (Meffe and Carroll, 1994). That is, it depends on the initial number and frequency of alleles. Under the infinite alleles model of mutation, for likely values of θ ($\theta = 4N_e\mu$, where μ = mutation rate), populations tend to have no more than 2-4 alleles at frequencies > 0.05 (Marshall and Brown, 1975). Frankel and Soulé (1981) provide examples of the effects of one-time drift on initial allele frequencies consistent with these theoretical expectations. For example, for $p=0.94$ and $q=r=s=0.02$, two of four alleles

are lost on reduction of population size to 10 individuals, whilst reduction to 50 individuals leads to 10% loss of allelic richness. By contrast, gene diversity is expected to be relatively little affected by bottlenecks, because rare alleles contribute little to total heterozygosity; as the expected reduction is $\frac{1}{2N}$, 95 percent of gene diversity is conserved in a bottleneck of ten individuals (Frankel and Soulé, 1981). The latter authors comment 'unless the number of founders is of the order of two pairs or fewer, the bottleneck, *per se*, is not the villain...most of the loss that ensues is attributable to events following the bottleneck'.

The above formulation implies that the fragmentation-mediated 'sampling' process is analogous to simple, unbiased random sampling. In practice, this is unlikely to be the case. Firstly, fragmentation is patently not analogous to simple random sampling, because it 'samples' spatially clustered groups of trees, not individuals distributed randomly in space. As limited seed dispersal in forest trees frequently leads to small-scale spatial genetic structure (Hamrick *et al.*, 1993; Loiselle *et al.*, 1995; Hamilton, 1999), fragmentation will lead to loss of more genetic variation than if a similar number of trees was randomly removed from the population. Secondly, if, as is frequently the case, fragmentation is non-random with respect to environmental conditions (e.g. soil type) (Laurance, 1999; Chapter Three), local adaptation to specific sites (Gilbert *et al.*, 1996; Linhart and Grant, 1996) could generate correlations between spatial genetic structures and deforestation criteria (*i.e.* those factors, such as soil conditions, which determine whether a given forest tract is removed), leading to loss of particular non-neutral alleles or allelic complexes (Nason *et al.*, 1997). S-alleles, *i.e.* those controlling self-incompatibility mechanisms, may be particularly susceptible to loss through founder effects as, due to frequency-dependent selection, allelic richness tends to be high, *i.e.* allele frequencies tend to be low. A decrease in allelic richness at incompatibility loci will tend to reduce fertility, as well as exerting selective pressure in favour of occasional self-compatible individuals; similar effects could occur in low density populations of dioecious and distylous species (Murcia, 1996).

FRAGMENTATION AND POLLINATION

Fragmentation leads to changes in abundance of many insects, including pollinators (Didham *et al.*, 1996). Particularly in tropical forests, where more than 90 per cent of trees

are thought to be animal-pollinated (Bawa *et al.*, 1985), effects of fragmentation on pollination are therefore to be expected. At least five specific mechanisms may be identified. Firstly, small fragments may be too small to support resident pollinator populations: possible causal factors include low numbers of particular plant species (specialist pollinators) or absence of high density resources (many generalists), lack of suitable nesting habitat and larval host plants, pesticide contamination from agricultural matrices, or invasion of competitors or predators (Murcia, 1996; Nason and Hamrick, 1997; Rathke and Jules, 1993; Roubik, 1989). Secondly, non-resident pollinators are less likely to visit small, isolated populations, as they offer lower resource levels, require more energy to reach, and may not even be detected. For example, Mustajärvi *et al.* (2001) found that bumble-bees preferred larger artificially established fragments of the outbreeding, self-compatible *Lychnis viscaria*. Groom (1998) reported that small patches of *Clarkia concinna* received fewer pollinator visits when certain isolation thresholds were reached, leading to reproductive failure due to lack of pollination. Jennersten (1998) found that *Dianthus deltoides* flowers in a fragmented area received fewer pollinator visits and set less seed than in a mainland area; hand pollination increased seed set, an indication of pollinator limitation (Vaughton and Ramsey, 1995). Similarly, Ågren (1996) found positive relationships between population size and seed production per flower and plant in the herb *Lythrum salicaria*, caused by pollinator limitation. Thirdly, forest destruction may lead to increased mean pollinator movements, as pollinators are forced to forage further afield. Fourthly, fragmentation-induced changes in plant density (which may result from reductions of population size within a finite area) may cause changes in pollinator behaviour. For example, in both the wetland perennial *Mimulus ringens* (Karron *et al.*, 1995) and the tropical tree *Shorea siamensis* (Ghazoul *et al.*, 1998), the proportion of pollinator flights between (as opposed to within) plants was negatively correlated with plant density. Finally, similar effects could also be caused through changes in pollinator species. For example, both Aizen and Feinsinger (1994a) and Dick (2001) reported fragmentation-caused increases in importance of feral honey-bees. Such changes could be beneficial in some cases. Dick's report, for example, describes the positive role of *Apis* as long-distance pollinators able to effect pollination in small and isolated fragments. Aizen and Feinsinger (1994a), by contrast, found that *Apis* bees tended to move between trees more rarely than native bees, whilst Gross and Mackay (1998) found that honey-bees

reduced fitness in the Australian pioneer shrub *Melastoma affine* by disturbing more effective native pollinators.

It should not be assumed that fragmentation will always be associated with changes in pollinator abundance and behaviour. Rather, the incidence of such effects is likely to depend on particular circumstances, e.g. fragment size, isolation and pollinator mobility. For example, for 14 species present in Andean cloud forest remnants, Murcia (1996) reported no differences in pollination between small and medium-sized fragments. She suggested that it is likely that pollination will be affected only when fragments reach extremely small size. Similarly, Aizen and Feinsinger's (1994b) study of effects of fragmentation on pollination of 10 species of Argentinian dry forest revealed that pollination decline was greatest in those species which receive few visits even in continuous forest.

Patterns of pollen transfer between trees affect both gene flow and mating systems. In the case of the latter, in self-compatible species, increases in within-plant movements would be expected to reduce outcrossing rates, due to increased geitonogamous selfing. Empirical results are consistent with this expectation. For example, Murawski and Hamrick (1992) and Murawski *et al.* (1990) established that outcrossing rate in *Cavanillesia platanifolia* is density-dependent: isolated trees had high selfing, whereas trees in groups were predominantly outcrossed. Similarly, Prober and Brown (1994) found negative relationships between tree density and values of Wright's fixation index (F_{is}) in fragments of *Eucalyptus albens*, although Wahlund effects provide an alternative explanation in this case. Aldrich and Hamrick (1998) found that pasture trees of *Symphonia glabra* showed higher selfing rates than neighbouring conspecifics in continuous forests, and attributed this to changes in pollinator foraging patterns. Population density effects have also been implicated in interpopulation variation in outcrossing rates in *Pterocarpus macrocarpa* in Thailand (Liengsiri *et al.*, 1998). Increased selfing leads to decreases in effective population size (Yeh, 2000) and increased homozygosity, with the attendant risk of inbreeding depression.

The possible impact on gene flow of changes in pollinator abundance and behaviour is discussed below, following a brief outline of migration theory.

FRAGMENTATION AND GENE FLOW

The magnitude and patterns of gene flow are central to the genetic effects of forest fragmentation, because gene flow may act to counter random genetic drift. This is not simply because immigration can reintroduce alleles lost by drift. As outlined below, if migration is sufficiently prevalent, in effect 'separate' subpopulations may behave as one larger population, thus reducing the importance of genetic stochasticity.

Theoretical formulation and investigation of the interaction between migration and drift has relied on three principal models, *i.e.* Wright's Isolation by Distance (IBD) model, Kimura and Weiss's Stepping Stone model, and Wright's Island model (Hamrick, 1987; Lande and Barrowclough, 1987). IBD applies to continuously distributed populations, whereas the latter two apply to spatially discrete subpopulations, such as those found in forest fragments. The Island and Stepping Stone models differ principally in the assumed mode of gene exchange between subpopulations. The former assumes that migration occurs at the same rate between all subpopulations. As such, it represents 'the extreme in long distance gene flow' (Hamrick, 1987). The Stepping Stone model, by contrast, assumes that gene flow occurs predominantly between adjacent populations; as pointed out by Wright (1969) himself, it is therefore more realistic. However, in the case of a two-dimensional spatial configuration of subpopulations, theoretical investigation of both models leads to a similar conclusion, *i.e.* that 'remarkably' little gene flow between subpopulations is required to prevent genetic drift (Hartl and Clark, 1989). Under the Island Model, numbers of immigrants as low as $mN_e > 2$ severely limits genetic divergence (Hartl, 2000), whereas, in the case of the Stepping Stone Model, Crow and Aoki's (1984) work demonstrated that only one to two migrants per generation are required to maintain overall panmixia. The difference between the two models is more significant when spatial configuration approximates the one-dimensional case, as, for example, in ridge-tops or fragmented riparian habitat. In this case, the more realistic Stepping Stone model produces markedly different results to the spatially 'blind' Island Model: considerably more gene flow per generation is required to maintain overall panmixia, because the correlation between neighbouring demes is higher (Hartl and Clark, 1989; Wright, 1969). For example, in a Stepping Stone model with migration rates of 0.1 and $(2)(10^{-5})$ respectively for adjacent and

long-distance gene flow, 'considerable' local differentiation will occur if $N_e < 100$ (Kimura and Weiss, 1964).

The main findings of the Island and Stepping Stone models in two-dimensional spatial configurations have received practical expression in the 'one-migrant-per-generation' (OMPG) rule, as applied 'in the USA [in] nearly every recovery plan that considers genetic issues and insularization' (Mills and Allendorf, 1996). The OMPG rule seeks to attain a balance between subpopulation and total diversity, *i.e.* avoiding both loss of alleles within subpopulations and the uniformity associated implied by complete panmixia. An equilibrium fixation index value of $F_{st}=0.2$ is considered to achieve this balance, corresponding to $mN_e=1$.

Many studies of forest fragmentation have tended to assume tacitly or implicitly that spatial isolation by fragmentation would also lead to reproductive isolation, or at least to some reduction in numbers of immigrants. There appear to be good reasons for such an assumption. Firstly, in many species, fragmentation necessarily involves the removal of one source of immigrant pollen, *i.e.* those trees formerly located in the deforested matrix. Secondly, as outlined above, fragmentation may lead to reductions in pollinator abundance, due both to lower populations of resident pollinators and reduced visitation. In themselves, both factors will tend to reduce the amount of incoming pollen or seed to a given fragment. Unless fragmentation leads also to at least equivalent reductions in pollination events involving 'home' pollen, this will lead to decreases in migration rate and, consequently, reductions in mN , the gene flow parameter. However, it is worth stressing that, as suggested above, such reductions in 'home' pollination are certainly an expected consequence of fragmentation, and consequently it is questionable whether an assumption of increased reproductive isolation is justified. It should also be mentioned that pollinator populations themselves may show relatively short-term evolutionary responses to fragmentation, *e.g.* adaptations for increased mobility (Van Dyck and Matthysen, 1999).

Recent studies of pollen dispersal in fragmented landscapes, which have demonstrated that long-range pollen flow is common in tropical trees, lend force to this conclusion. For example, Apsit *et al.* (2001) reported mean immigration over three years of >70 per cent to a *Enterolobium cyclocarpum* population, in spite of the fragmented surrounding landscape, and

commented that 'pollen dispersal distances of 1500m may be relatively common in *E. cyclocarpum*'. Dick (2001) reported *Apis*-mediated pollen flow of up to 3.2km to *Dinizia excelsa* (Fabaceae) fragments. White and Boshier (2000) reported that 70 per cent of pollen flow to an isolated *Svietenia humilis* tree came from a stand located 4.5km away, whilst Nason and Hamrick (1997) reported that, in small island populations of *Spondias mombin*, 90-100 per cent of progeny produced was the product of pollen flow from 80-1000m away. Although such extensive post-fragmentation gene flow does not necessarily imply either reduced, increased or static levels of gene flow relative to pre-fragmentation conditions, it does imply that, in such fragments, genetic drift is less likely to threaten population viability. At the same time, for a number of reasons, it would be unwarranted to conclude that migration will in general remove the possible threat of genetic drift in fragmented populations. Firstly, migration rates are not necessarily independent of population size. At the extreme, for population size $N=1$ of any self-incompatible species, migration rate cannot be less (or more) than 50 per cent, whilst for any small fragment population, extrafragment trees may greatly outnumber population members, leading to the possibility of high expected gene flow, when this is not precluded by pollinator foraging capability. Such relations may not apply in larger populations. Secondly, migration numbers well in excess of 'OMPG' may be insufficient to halt centrifugal tendencies. Mills and Allendorf (1996), in their critique of unquestioning application of the OMPG rule, suggest that failure in the assumptions of the Island model (Whitlock and McCauley, 1999) imply that a more suitable general rule of thumb would be 'a minimum of 1 and a maximum of 10 migrants per generation'. This, or even higher numbers, would apply particularly in the case of gene flow in spatial configurations tending more to the one-dimensional than two-dimensional case. Thirdly, the expected mitigatory effect of gene flow applies principally to the equilibrium situation. Insofar as prefragmentation populations are in equilibrium, a sudden reduction in genetic diversity, but with unchanged gene flow, constitutes a disturbance of equilibrium, *i.e.* genetic variation will be less than expected for prevailing gene flow. Over time, gene flow is expected to restore genetic variation until the point of equilibrium is reached. However, in the interim, which can last many generations (Whitlock and McCauley, 1999) populations will still be subject to reduced genetic variation, with potentially serious results (for example, loss of S-alleles).

Evidently, the genetic effects of forest fragmentation are not easily predictable based on theory, even when reinforced by empirical observations of processes with genetic implications. Evaluation of the gravity of the genetic threat posed by fragmentation can only be satisfactorily evaluated by taking into account empirical studies.

GENETIC VARIATION IN FRAGMENTED PLANT POPULATIONS

Two broad approaches of detecting effects of fragmentation on genetic variation have been used. Firstly, comparison of fragmented populations with unfragmented populations, *i.e.* either comparable populations in unfragmented landscapes, or prefragmentation cohorts of the same populations. Secondly, relationships between measures of genetic variation and indices of intensity of fragmentation (*i.e.* typically, population size or, occasionally, isolation) have been studied. Studies of these sorts have produced diverse results. However, in several cases, important genetic effects have been detected, whilst, when this has not been the case, the lack of effect is generally explicable in terms of probable values of the factors outlined above.

Fragmentation has been associated with declines in allelic richness in a number of cases. For example, in 17 fragmented populations (N from 1 to 430) of the perennial *Swainsona recta*, Buza *et al.* (2000) reported significant effect of log population size ($R^2=0.70$, $p<0.001$), due to absence of rare alleles from small populations. Similar relationships were reported by Van Treuren *et al.* (1991) (respectively 14 and 12 populations of the bee-pollinated perennials *Salvia pratensis* and *Scabiosa columbaria*, $N=5-1500$, $14-100,000$, $r=0.57$, 0.65) and Prober and Brown (1994) (25 fragments of *Eucalyptus albens*, N from 14 to >10000 , $R^2=0.39$, $p=0.007$). Rajimann *et al.* (1994) (25 populations of *Gentiana pneumonanthe*, N from 6-100,000) reported A_s rather than A . They found positive correlations between log population size and both this parameter and P , the proportion of polymorphic loci, although the relationships with A_s was only marginally significant (A_s : $r=0.33$, $p=0.015$; P : $r=.49$, $p=.013$). The case of the daisy *Rutidosis leptorrhynchoides* (Young *et al.*, 2000) is of special interest. In this case, not only was overall allelic diversity in 17 fragment populations ($N=5-95,000$) strongly positively related to log population size, but fragmentation also led to loss of incompatibility alleles. The consequences of this extend beyond simple loss of fertility. In the authors' words: 'firstly...reduced S-allele richness lowers the effective population size, further exposing

populations to genetic drift. Secondly...low mate availability may reduce population-level seed-set while increasing interplant variance in fecundity...Thirdly, severe mate limitation will also favour self-compatible plants that occur at low frequencies'. Effects of fragmentation on diversity of S-alleles has also been implicated in local extinctions of the Great Lakes lakeside daisy, *Hymenoxys acaulis* var. *glabra* (Demauro, 1993).

In all the above instances of strong relationships between population size and allelic richness, the former parameter ranges from very small (*i.e.* <10) to several orders of magnitude greater. In studies in which no relationship has been detected, the range of population sizes has tended to be less or, perhaps more importantly, very small populations were not included (e.g. *Festuca ovina* (*N* of 25-980, *Lychnis viscaria* (10-680), *Arabis thaliana* (*N* of 2 to >100) (all Berge *et al.*, 1998), *Acacia anomala* (*N*=3-50) (Coates, 1988), *Arnica montana* (*N*=20-1500) (Kahmen and Poschlod, 2000), *Microseris lanceolata* (*N*=87-140,000) (Prober *et al.*, 1998).

In many cases, loss of allelic richness has been attributed to founder effects. Although gene flow between fragments might restore lost alleles relatively quickly (conceivably, in the first reproductive events after fragmentation), loss of non-neutral variation is also likely to affect fitness relatively quickly. Furthermore, alleles can only be restored if present in the post-fragmentation population as a whole. Young *et al.*'s (1993) work with *Acer saccharum* illustrates this point. Although fragmentation appeared to increase gene flow between fragments and, consequently, individual fragments tended to be more similar than the controls (tracts in continuous forest), the total number of alleles over all fragments was less than in the tracts of continuous forest, apparently because founder effects led to the loss from the fragmented area of the alleles in question.

As outlined above, gene diversity is not expected to be as sensitive to founder effects. Nevertheless, decline in values of this parameter have been reported. In Prober and Brown's (1994) study of *E. albens* (1994), log population size was more strongly related with gene diversity than with allelic richness ($R^2=0.49$, $p=0.001$). A relatively high correlation were also reported by Raijmann *et al.* (1994) for 25 populations of *Gentiana pneumonanthe* ($r=0.34$, $p=0.09$). However, in a number of other studies of relatively-recently fragmented populations, there were no clear relationships between gene diversity and log population

size (Berge *et al.*, 1998; Foré *et al.*, 1992; Young *et al.*, 1993, 1999, 2000; Buza *et al.*, 2000; Van Treuren *et al.*, 1991; Coates, 1998; Kahmen and Poschold, 2000). These results contrast with those from long-isolated, naturally-fragmented species. For example, Sampson *et al.* (1998) found low genetic diversity and high population differentiation ($G_{st}=0.24$) in isolated populations of the granite-outcrop species *Eucalyptus crucis*, whilst in populations of the dioecious conifer *Halocarpus bidwillii*, estimated to have been naturally fragmented for about 8000 years (*i.e.* ≈ 100 generations), gene diversity was significantly related to log population size ($p=0.004$, correlation coefficients not reported) (Billington (1991).

Both *E. albens* and *G. pneumonanthe* are self-fertile, and both of the mentioned studies reported evidence of increased selfing in fragmented populations. At least in the case of *G. pneumonanthe*, there is also evidence of higher selfing in small populations (see below). Such effects would also enhance relationships between population size and H_e (*i.e.* beyond the expectation for drift alone), as observed by Young *et al.* (1996) in connection with these two studies. This suggestion finds some support in the other mentioned studies. *F. ovina* (Berge *et al.*, 1988), *R. leptorhynchoides* (Young *et al.*, 1999) and *A. montana* (Kahmen and Poschold, 2000) are self-incompatible, whilst *A. thaliensis* (Berge *et al.*, 1998) is predominantly selfing. In these cases, there is no reason to expect correlation of selfing rate with population size, and therefore no reason why inbreeding should contribute to the gene diversity - population size relationship. Furthermore, in the cases of (possibly) self-compatible or partially self-compatible species such as *Acer saccharum* (Foré *et al.*, 1992), *Lychnis viscaria* (Berge *et al.*, 1998) and *Swainsona recta* (Buza *et al.*, 2000), such relationships are likely to be mediated by pollinator behaviour, and, rather than being axiomatic, are conditional on size and distribution of fragments.

A number of studies has examined the relationships between genetic variation and fragment isolation. For example, in *A. saccharum* patches (mean isolation of 58.9m), Foré *et al.* (1992) reported a significant positive relationship between number of canopy trees and observed heterozygosity of juveniles in highly-isolated patches (*i.e.* ≥ 65 m), suggesting that, in more isolated patches, gene flow may be insufficient to counteract factors responsible for higher observed homozygosity. Juveniles in highly isolated patches also had a significantly greater proportion of monomorphic loci. Prober and Brown (1994) found that the combination of

log population size and distance to the nearest large stand of *E. albens* explained 48 per cent of variation in allelic diversity, suggesting the presence of 'isolation thresholds' beyond which fragmentation effects might occur. Similarly, Hall *et al.* (1996) in their report on genetic diversity and differentiation among nine populations of *Pithecellobium elegans* in Costa Rica found that genetic variation in the small fragment closest to a large population was greater than that in the small fragments further away, suggesting the presence of 'important genetic links', and implying that in the other fragments the degree of isolation has eroded genetic variation, or at least not permitted its postbottleneck restoration. Dayanandan *et al.* (1999) found that genetic distance between adult and seedling cohorts in fragment populations of *Carapa guianensis* (Meliaceae) in Costa Rica was greatest in the most isolated population, which was also the only one in which allelic diversity was higher in the adult cohort. In other studies, relationships between isolation and diversity have been absent (Buza *et al.*, 2000; Young *et al.*, 1999).

Finally, several studies have reported inbreeding depression in fragmented populations. In *Swainsona recta*, Buza *et al.* (2000) found negative correlations between population size and fixation indices, and negative correlations between the latter and germination percentages. Fischer and Matthies (1998) found that remnant population size in *Gentianella germanica* was correlated with plant survival in a common garden experiment (Spearman $r=0.50$, $p<0.05$). Gigord *et al.* (1998) documented inbreeding depression on selfing in the Madagascan endemic tree *Dombeya acutangula* ssp. *acutangula*: outcrosses produced more fruits per flower, more seeds per fruit and lower proportions of flattened, lighter seeds. In a comparison of seven populations, population size was strongly positively correlated with one of these variables (mean number of seeds per fruit). Although between-population variation in proportion of flattened seeds was not significant, mean proportion in the three smallest populations ($N\leq 10$, mean 15.3% flattened) were notably higher than in the largest population ($N=81$, 5.4% flattened). In seven small populations of the self-compatible *Gentiana pneumonanthe*, Raijmann *et al.* (1994) reported a significant correlation between selfing-rate and log population size, apparently caused by decreased visitation of the relatively non-mobile bumblebee pollinator. As heterozygosity is known to affect fitness in this species (Oostermeijer *et al.*, 1994), this result implies inbreeding depression in small populations of this species.

CONCLUSIONS

Clearly, fragmentation effects on population genetics of forest trees and other plants are complex and difficult to predict. In particular, theoretical considerations are perhaps more useful in understanding and rationalizing, *i.e. a posteriori*, empirical results, rather than predicting them. However, it is worth emphasizing that both expectations and findings suggest that fragmentation can exert a rather rapid effect on genetics of fragmented populations, both through effects on pollinators and founder effects. Although the theoretical formulation of random genetic drift in subdivided populations appears to correspond closely to fragmentation, it does not and is not intended to describe all consequences of population subdivision. Consequently, genetic response to fragmentation will not necessarily occur only in long-term, and neither will it necessarily be easily mitigated by gene flow.

3. HISTORICAL AND SPATIAL PATTERNS OF DEFORESTATION AND FRAGMENTATION IN GUANACASTE PROVINCE, COSTA RICA: CHARACTERIZATION AND GENETIC IMPLICATIONS

INTRODUCTION

The seasonally dry tropics have been heavily deforested (Mooney *et al.*, 1995). In Mesoamerica, this trend is particularly marked, due largely to high human population concentration (Murphy and Lugo, 1995). According to Janzen (1986), only around 2 per cent of the original forest cover remains. However, even in such heavily disturbed zones, deforestation of a given area is rarely complete (Schelhas, 1996). Rather, one or more tracts within the original continuous forest remain. If viable, communities and populations in such fragments have the potential to mitigate the impact of deforestation on biodiversity. However, such viability is not assured, because small populations are subject to a number of pressures which are absent or of less importance in large populations (Soulé, 1987). These include a number of factors which, if present, have the potential to reduce genetic diversity in fragmented populations (see Chapter Two). This is of concern because genetic erosion may lead to decline in fitness and evolutionary potential (Brook *et al.*, 2002; Frankham and Ralls, 1998; Young *et al.*, 1996).

The genetic impact of fragmentation depends in part on its spatial and temporal patterns. Both partially define the fragmentation process and state, and thereby determine the intensity of fragmentation. Because the latter is a spatial phenomenon, this is most clearly seen in the case of spatial patterns. However, temporal patterns are also of great interest, as the effect of evolutionary processes depends not only on their intensity but also on the number of generations for which they act.

In the present document, historical and spatial patterns of forest fragmentation in a zone of seasonally dry, northwestern Costa Rica are characterized. After a description of the study area, forest history since the Spanish conquest is traced, based on published sources, maps and aerial photos, culminating in a quantitative analysis of forest change and state since the

1940s. Finally, the biological implications of the findings are discussed, with particular emphasis on genetic aspects.

THE STUDY ZONE

The study zone, an approximately 350km² rectangular section of the Pacific slope of Costa Rica (Figures 3-1, 3-2), lies between latitude 10°35' and 10°17'N and longitude 85°05' and 85°12'W. It rises from sea level at its southern limit, some 12km north of the Gulf of Nicoya, to 250m a.s.l. just north of the village of Palmira, approaching the foothills of the Miravalles and Tenorio volcanoes. Around 95 per cent of the mean annual rainfall of 1693.4mm ($\sigma = 459.4$) falls between May and November (San Luis, Cañas meteorological station, 1921-1978, MIRENEM, 1988). Altitude varies from 0-200m a.s.l.; mean annual temperature at 95m a.s.l. is 27.5°C (Jiménez *et al.*, 1987). The dry season is characterized by strong (up to 90km hr⁻¹) northerly winds (Coen, 1983) and temperatures up to 37°C. Soils in the study zone fall into two broad categories: south and immediately north of the Interamerican Highway are mollisols, alfisols, vertisols and alluvial inceptisols of generally medium to high agricultural potential, although with some drainage problems. An escarpment three to four kilometres north of the Highway marks a transition to shallower inceptisols, some derived from deep volcanic ash deposits (Oficina de Planificación Sectorial Agropecuaria, 1987). Within both areas, topography is generally flat to undulating. Two main land-uses predominate in the study zone, reflecting the soil differences mentioned above: the southern zone is dominated by latifundist industrial agriculture (flooded rice and sugar-cane) whilst the northern zone is dominated by beef-cattle ranching, with mixed land tenure. The rivers Cañas, Corobicí, Tenorio, Tenorito and their numerous tributaries flow in a predominantly southwest direction across the study zone, draining into the River Bebedero and thence to the Gulf of Nicoya. They form deep ravines and canyons in the tuff overlaying parts of the northern sector of the study zone. Forest is the natural vegetation of the entire study zone, with the possible exception of some parts of the poorly drained vertisols in the southern sector. Naturally occurring species include a number of high commercial value, e.g. mahogany (*Swietenia macrophylla*), Spanish cedar (*Cedrela odorata*), rosewood (*Dalbergia retusa*) (Jiménez *et al.*, 1987; see also Appendix 3). The city of Cañas, population 18,798 (INEC, 2002), is the only relatively large centre of population in the study

zone. In addition, there are two larger villages: Bebedero (pop. 2123) and Palmira (pop. 916) (INEC, 2002). The former, situated at the confluence of the Corobicí and Tenorio rivers (Figure 3-2), is a long-established river port and has been settled since at least 1687 (IGN, 1972).



Figure 3-1. Costa Rica: major cities, relief and location of the study zone.

topography reflect common features of the zone, e.g. industrial agriculture on high quality soils, prominence of cattle ranching and the canyon rivers of the northern sector (*c.f.* Vargas Ulate, 2001).

The zone is known to have been inhabited at the time of the Spanish conquest in the early 16th century; Ibarra (1990) places it within the Zapandí and Corobicí tribal areas. Consequently, it is unlikely that it was undisturbed during pre-Columbian times. In particular, clearance of lands located close to permanent watercourses, e.g. on the fertile mollisols close to the River Tenorio (Figure 3-2), seems likely. According to Meléndez (1955), such locations were preferred sites for indigenous habitation; indeed, the same author mentions the discovery of petroglyphs in the Paso Hondo farm, which is located in this area. In spite of this, there is no reason to believe that human activity in the pre-Columbian period led to extensive deforestation or forest fragmentation. There appear to be no records of major population centres within the study zone, whilst descriptions of the subsequent colonial period correspond to those of a very largely forested (pre-Columbian) landscape in the process of conversion to the largely deforested one of today.

This deforestation process is traced here based on two principal sources of evidence: general accounts of the agrarian history of Guanacaste province, and specific references to the study zone. Although the latter are less complete than the former, joint consideration of both sources of evidence is informative. Reconstruction is also facilitated because many of the general references apply specifically to the area known historically as the Bagaces Valley, of which the study zone forms a part.

GENERAL PATTERNS OF DEFORESTATION IN LOWLAND GUANACASTE

Deforestation in lowland Guanacaste can be described in three main phases, all associated with particular historical-agricultural and land-use developments, as described below.

Phase One: feral cattle ranching (1560-to late 19th century)

Cattle - equine and bovine - was first introduced to Guanacaste in 1561 (Fournier, 1992). However, the result was not the birth of an intensive cattle-based agriculture, but rather the founding of what would become a semi-feral cattle population that would be 'managed'

essentially as a renewable natural resource for more than three centuries. In large measure, this was because low human population densities (the 1688 population in the Bagaces Valley, which includes the study zone, was 297) and large distances to markets precluded more intensive husbandry (Edelman, 1992). Cattle roamed in unfenced, wooded areas known as *sitios*, owners having rights to a certain number of cattle thought to be grazing a particular *sitio*, plus a proportion of the 'natural increase' (Edelman, 1992).

Phase Two: Timber exploitation, introduction of exotic pasture and cattle (1880-1930s)

Between 1880 and 1920, there was a marked expansion (approximately tenfold) in timber production from Guanacaste (Edelman, 1992). As well as leading directly to forest destruction, much of the proceeds from timber exploitation were invested in pasture improvement (Acuña y Molina, 1991), *i.e.* sowing of exotic varieties which permitted year-round grazing and up to 300 per cent increase in stocking rate. Improved pasture was sown preferentially on fertile sites close to rivers, and seed was also broadcast in the *sitios* (Edelman, 1992). In 1909 Guanacaste was legally declared a 'livestock zone', *i.e.* agriculturists became responsible for 'fencing out' cattle in order to protect their crops, rather than the onus being on cattle-owners to 'fence in' (Edelman, 1992). Both timber extraction and cattle expansion must have contributed to further forest destruction and attrition. By 1940, according to Becker (1943), the extractive timber phase was 'complete on all privately owned land with easy access to the main rivers and their tributaries'. However, it should be noted that selective logging of commercially valuable species such as mahogany (*Swietenia* spp) and Spanish cedar (*Cedrela odorata*), which tend to occur at low densities, is not likely in itself to have led to outright forest destruction.

Phase Three: Deforestation in the 1950s and after

Between the 1950s and the 1970s, the area under pastureland in Guanacaste more than doubled as a result of expansion in the beef trade (Edelman, 1992); the latter author stresses that, in comparison with this phase, deforestation in the 1880-1930 period was relatively insignificant.

DEFORESTATION IN THE STUDY ZONE

Phases One and Two

There is no doubt that the wider changes outlined above were manifested also in the study zone. This is particularly well-established for the southern sector of the study zone. Edelman (1992) reported that concessions for grazing *sitios* were made in the Bagaces Valley as early as the 1560s (that is, coinciding with the introduction of cattle). Much of the southern part of the study zone is occupied by one large holding, the Taboga (formerly Higuerón) *hacienda*. Taboga, together with the contiguous Paso Hondo property (now mostly broken up by land reform), are amongst the oldest established properties in Guanacaste, founded no later than 1712 and 1787, respectively (Gudmundson, 1983). According to Meléndez (1955), in 1792 both were amongst the main farms of the region. Although, in the period to 1611, cattle production was mostly equine (mules) rather than bovine (Quirós, 1990), by the early 18th century (at latest) this had changed: the traveller John Cockburn recorded seeing 'great herds of cattle' on the east bank of the Tempisque (Meléndez, 1974), whilst other documents record similar observations as early as 1719 (Meléndez, 1974, citing Fernández Bonilla, 1884, p. 315). Edelman (1992) comments that 'in the colonial period [*i.e.* to 1821], the lowland plain along the eastern bank of the Tempisque, between the towns of Cañas and Liberia, was the principal area of cattle production within the Costa Rican jurisdiction'. A number of *sitios*, now mostly no longer forested or wooded, are marked on the 20th Century 1:25000 and 1:50000 topographic maps of the study zone, e.g. Sitios de Paso Hondo (IGCR, 1956a), Sitio Paraiso (IGCR, 1956b), Sitios de la Uvita (IGCR 1956c), Sitio Cascante (IGN 1973). They testify eloquently to the destruction of the zone's forest, as well as its previous use for extensive grazing.

Taken as a whole, the above information indicates that much of the southern part of the study zone was subject to grazing pressure from an early date. The effect of such pressure was recorded by the traveller von Seebach, who visited the area in 1864 (Meléndez, 1974). He refers specifically to the crossing of the Duquesa and Reventado rivers (Figure 3-2), and described the countryside as 'not very dense and ... interspersed with small pastures' (Meléndez, 1974).

It is also very likely that the zone was affected by the phase two changes. There was year-round grazing (*i.e.* based on exotic pasture grasses) in Hacienda Mojica (bordering the River

Tenorio) by 1885 (Gudmundson, 1983), whilst the same author reports introduction of exotic pastures in the Paso Hondo farm in the 1930s. Edelman (1992) reports that between the 1920s and the 1930s Julio Sánchez L. had 13,624 ha dedicated to (Nelore) cattle in the Taboga farm. With regard to the timber expansion, León (1952) wrote that Bebedero, effectively an enclave between the Mojica and Paso Hondo *haciendas*, was the most important river port in the timber trade. It would be odd if the timber exported did not include the product of the neighbouring properties themselves, a conjecture supported by Glander and Nisbett's (1996) comment that mahogany and other species were being logged at Paso Hondo during the 1920s. Independent of the causes - pasture conversion or logging - the major point is that there is evidence that large sections of the southern sector of the study zone was already deforested before the major cattle expansion of the 1950s. Céspedes Marín travelled the road from Bebedero to Cañas in 1923, and describes the wide, open plains ('planicie tan extenso') interspersed with occasional ('uno que otro') trees (Meléndez, 1974); given his route, this description is almost certainly of the Mojica, Paso Hondo and Taboga properties. More clear still, although recording a later date, is the evidence of the 1:25000 map series, published in the 1950s but based on aerial photos from 1945, which shows that by this date much of the area of these two properties was already deforested (IGCR 1955, 1956a, 1956b, 1956c, 1956d).

Much of the northern sector of the study zone was formerly occupied by the Hacienda Tenorio property; according to Sequeira's (1985) map, by 1880 this included the lands between the Corobicí and Tenorio Rivers north to the upper slopes of the Tenorio and Miravalles volcanoes. The *hacienda* was founded by 1770 (Gudmundson, 1983), when it was described in state records as the *sitio* Tenorio, a designation that implies grazing activity. By 1794, at least 1000 cattle were located in the property (Gudmundson, 1983). Colegial (1989) reported that, when taken over by the United Fruit Company in 1949 (for production of mules for use on banana plantations), only a few patches of the original forest had been cleared by small farmers for pastures, coffee and cane (the presence of smallholders has also been recorded by Edelman (1992)). However, records cited by Gudmundson, indicating that 7000 of the total 17000 ha had been cleared by 1935, suggest that Colegial's more anecdotal account underestimated the amount of land clearance prior to acquisition by the United Fruit Company. For present purposes, the situation is confused further because

much of the Hacienda Tenorio lies outside the study zone, i.e. the medium to upper slopes of the volcanoes. It seems likely that historic grazing would have been concentrated in the southern sectors, which are closer to the historic cattle-drivers' road and contiguous with the other known cattle ranches. This is supported by the designation on the 1956 map series (IGCR 1956d) of a large wooded area of what is now the La Pacífica farm as 'Potrero [i.e., pasture] La Pacífica', suggesting a traditional wooded *sitio*. The 1945-based map series offers no guidance with respect to the northern zone itself, as this area was not covered by this map series.

Quantitative analysis

Methods

For much of the study zone, forest cover before the cattle-boom of the 1950s is documented in the 1:25000 map series published in 1950-1951, based on aerial photos taken in 1945 (IGCR 1955, IGCR 1956a, 1956b, 1956c, 1956d). However, the zone to the north of 10°30'N and east of 85°07'W, corresponding roughly with the northern sector of the study zone, was not covered by this series. However, this zone is included in the first 1:50000 map series, based on photos taken from 1945-1961. The state of forest cover in the study zone in 1945 (southern zone) and 1961 (northern zone) is characterized here based on random point sampling of these map series. Each sample point was classified as forest, swamp, deforested land (i.e. non-swamp land no longer under forest), and other land (roads, rivers, lakes, etc.) and for its location relative to the nearest watercourse (i.e. whether within 100m distance). Estimated proportional land cover by class was estimated as the proportion of all points falling in each class. Two-by-two contingency tables for presence of forest in relation to distance from nearest watercourse (i.e. whether within 100m) were compiled. Equality of ratios (forested/deforested) in the two distance classes was tested using Fisher's exact test; the odds ratio for forest cover in the two watercourse-distance classes and the associated confidence intervals were also estimated (Ramsey and Schafer, 1997). SPSS was used for both procedures (SPSS, 1996).

Current forest cover was characterized based on a set of 1:10000 aerial photographs taken in 1998. A sheet with 20 systematically spaced punch-holes was placed over each photo (each photo corresponds to an area of approximately 5.3 km²), and four of these were randomly

selected for location of random point samples at their centres. At each sample point, land cover was classified as either riparian forest, non-riparian forest, pastureland, agricultural land, scrub, swamp and other use (aquatic, public utilities, roads, urban zones and areas which could not be classified). In addition, for sample points which fell on unforested areas, the presence of individual trees in a circle of approximate area 0.25ha centred on the sample point was scored (1=one or more trees present, 0=absent). This area, although arbitrary in that it was derived from the scaled-up area of the punch-circles drawn on the photos, proved satisfactory in order to detect differences in tree stocking between matrix types. In addition, the presence of continuous forest within 100m of the sample point was scored. Sample points were also scored for their location relative to the nearest watercourse, as for the map data. The same analysis approach was used as for the map data. As field observations suggested that pastureland and proximity to water-courses both had positive effects on forest or tree presence, one-tailed probability values were used.

Results

In 1945, the southern sector of the study zone was close to 30 per cent forested (Figure 3-3a). By 1998, this proportion had fallen to 13.4 per cent (Figure 3-3c), of which almost half (6.7 per cent of total) consisted of riparian remnants. In 1945, points within 100m of watercourses were estimated to be almost three times more likely to be forested (Table 3-1). In 1998, points within 100m of watercourses were estimated to be more than 5 times as likely to be forested (Table 3-2). In addition, locations in pastureland are significantly more likely to have individual trees within 0.25ha surrounding area and to be within 100m of forest (Tables 3-3, 3-4).

In the case of the northern sector of the study zone, forest cover estimate for the 1945-1961 period (15.0 per cent) was lower than the 1998 estimate (24.2 per cent). In the case of the earlier period, there was no significant effect of proximity to watercourse on forest cover (Table 3-1). In the case of the 1998 data, locations within 100m of a watercourse were almost three times more likely to be forested (Table 3-2). Meaningful comparisons between pastureland and agricultural land were not possible for this sector because of the low proportion (2.0 per cent) of agricultural land.




Figure 3-3. Land use in the study zone, 1945-1998: estimated percentage cover (standard error) by use category.

Discussion

The recent forest history of the study zone appears to diverge in several ways from the general trends for Guanacaste. Although there was considerable deforestation in the southern part of the study zone in the post-1945 period, this cannot be considered to more significant than previous deforestation, as the zone was already substantially deforested by this time. Deforestation was certainly rapid during this period, and probably more rapid than at any other time, but was not concentrated during this period. The southern sector also appears to be atypical, in that post-1950s deforestation was associated at least as much with industrial agriculture as with cattle expansion. A large area of the study zone around the low hills known as Las Lomas, on the Hacienda Taboga, was deforested sometime after 1956, apparently to feed the new sugar-mill established on the farm in 1958 (Edelman, 1992)

(Figure 3-4). The La Pacífica farm appears to have been deforested in the same period: as mentioned above, much of it appears as the wooded 'Potrero La Pacífica' on the 1945 map series. In aerial photos dating from 1956, it is comprised of mixed open and closed woodland, but by 1986 had been largely deforested. Currently, the farm is a mosaic of pastureland, forest remnants, shelterbelts and agriculture (Jiménez *et al.*, 1987).




Figure 3-4. Las Lomas and Llanos de San Pedro area, Hacienda Taboga, in 1956 (left) and 1986 (right)

In the case of the northern section of the study zone, the foregoing analysis suggests an increase in forest cover following the 1945-61 period. Although it is possible that this is partially associated with mapping criteria discrepancies, (e.g. similar to those mentioned by Sader and Joyce, 1988) it probably also reflects abandonment or partial abandonment of marginal farms, e.g. in the hilly zone between Cañas and Palmira (Figure 3-2), where there are relatively large areas of young secondary forest (*tacotal*).

The implications of deforestation patterns as revealed both by the foregoing analysis and suggested by historical trends are considered below.

GENERAL DISCUSSION

Evidently, forest in the study zone has been profoundly affected by human intervention since the Spanish conquest. Although the most marked changes are likely to have occurred during the acceleration of deforestation from the late 19th century, it should not be assumed that the forest grazing practiced before this time was harmless. Judging from the appearance of traditional *sittas* such as that pictured in Becker (1943), it seems to have culminated in a radical transformation of the forest. Grazing within forest has been identified as one of the forest practices expected to affect genetic processes (Namkoong *et al.*, 1996). Specifically, the latter authors comment that 'grazing would be expected to have a thinning effect on regenerating vegetation and hence could affect genetic drift. It could also affect the understorey vegetation that may also be directly grazed and hence exert a selection effect and changes in the population density of those species, affecting their selection and drift effects. Since grazing may also compact soils and alter stand structure, it may also induce selectively significant environmental changes'.

Such effects would not be expected to impact all species in the same way. More palatable species, and those with seeds less able to survive ingestion by cattle, would be expected to be most subject to reduction of population density. At one extreme, population fragmentation, as well as density reduction, may have occurred, as susceptible species disappeared from sites with severe grazing pressure. Recent studies suggest that probable consequences of population density reduction go beyond the selection and drift effects identified above, as reduction in population density of forest trees may affect pollinator behaviour. For example, in the tropical tree *Shorea siamensis*, Ghazoul *et al.* (1998) found that the proportion of pollinator flights between (as opposed to within) plants was negatively correlated with plant density. In fully or partially self-compatible species, such changes in pollinator behaviour may lead to increases in selfing rates. Thus, Murawski and Hamrick (1992) established that outcrossing rate in *Cavanillesia platanifolia* is density-dependent: isolated trees had high selfing, whereas trees in groups were predominantly outcrossed. Increased selfing rates are likely to lead to inbreeding depression, as in the case of the Madagascan endemic *Dombeya acutangula* ssp. *acutangula*. (Gigord *et al.*, 1998). In self-incompatible species, decreased interplant movements may result in selection pressure in

favour of increased self-compatibility (Murcia, 1996), with similar long-term consequences as for self-compatible species.

From the late nineteenth century, forest change in the study zone seems to have been typified by removal and fragmentation rather than density reduction, as traditional *sitios* were converted to improved pastureland or to agriculture. The earlier intensification of deforestation - i.e., as compared to the post-1950s increase in other parts of the Province - implies an earlier onset of problems associated with habitat fragmentation.

The pattern of deforestation in the study zone, as revealed by the foregoing quantitative analysis, has implications for the seriousness of such problems. Firstly, land conversion to grazing - both by attrition in the long (post-)colonial phase, and in subsequent change to 'improved' pastureland - has been less destructive of trees and forest than conversion to industrial agriculture. The latter is associated with 'classic' matrix-island landscape (e.g. Figure 3-4), whereas grazing lands in the study zone have a far higher density of individual trees and a higher tendency to be close to forests, reflecting a more diverse landscape structure. As presence of intervening trees and lower mean distance to forest fragments is likely to reduce reproductive isolation, this implies that forest fragments located in pastureland are likely to be less reproductively isolated than forest fragments located in agricultural lands. This is of relevance to viability of forest fragments, as the genetically homogenizing effects of random genetic drift can be mitigated by relatively low amounts of immigration (Hartl and Clark, 1989; Mills and Allendorf, 1996). Furthermore, small, isolated fragments may be subject to pollinator limitation of reproductive output (Murcia, 1996; Groom 1998; Jennersten, 1998). However, it is worth noting that the apparent relative benignancy of the pastureland matrix is likely to be a transient phenomenon without active human intervention to maintain current stocking levels. Current pasture trees are remnants of previous forests or more heavily wooded *sitios* and recruitment in pastureland appears to be virtually nil.

Secondly, deforestation and fragmentation in the study zone has clearly been a spatially non-random process. In both sectors of the study zone, proximity to watercourses is associated with significantly and notably higher probabilities of forest persistence. Both non-randomness as such and the particular 'bias' in question have important genetic

implications. A number of species native to the study zone occur in both riparian and non-riparian situations, e.g. *Pithecellobium saman*, *Anacardium excelsum*, *Astronium graveolens*, *Bombacopsis quinata*, *Enterolobium cyclocarpum*, *Spondias mombin*, *Swietenia macrophylla*. It is possible that the marked environmental differences between riparian and non-riparian forest in the study zone (Janzen, 1976) could have given rise to local adaptation, implying greater than expected loss in genetic diversity than under random deforestation (Young *et al.*, 1996). It is also possible that deforestation may have led to reduction in gene flow between riparian and non-riparian remnants, thereby favouring local genetic differentiation. This outcome might lead to higher overall diversity, but at the cost of loss of some populations and increased isolation of remnants.

A 'bias' in forest retention in favour of riparian types implies a change in the spatial characteristics of forest in the study zone, i.e. a redimensioning from two to one dimensions. This is of particular relevance to genetic impacts of forest fragmentation. Although, as indicated above, theory predicts that only a small number of immigrants are needed to offset the genetically homogenizing effects of random genetic drift, this conclusion applies to a two-dimensional configuration of populations. The same body of theory predicts that configurations tending more to unidimensionality, such as population fragments located on essentially linear landscape elements such as ridge-tops and watercourses, will be more prone to population subdivision and associated inbreeding effects than two dimensional configurations. Under such conditions, considerable more gene flow per generation is required to avoid accumulation of relatedness in subpopulations (Hartl and Clark, 1989; Wright, 1969). For example, under the Stepping Stone migration model (which assumes that gene flow occurs predominantly between adjacent populations) and with migration rates of 0.1 and $(2)(10^{-5})$ respectively for adjacent and long-distance gene flow, 'considerable' local differentiation will occur if $N_e < 100$ (Kimura and Weiss, 1964). Consequently, such populations are expected to be more susceptible to population fragmentation and isolation than those located in two-dimensional configurations. The prevalence and special characteristics of riparian remnants suggest that they should be taken into account in studies of genetic and other effects of forest fragmentation.

The higher probability of forest persistence close to watercourses may to some extent be attributable to legal protection. Current Costa Rican law prohibits logging within 15m of

rural watercourses, or within 50m if on sloping ground. (Costa Rica Legislative Assembly 1996). However, this law has been widely violated, with virtual impunity (Porras and Villareal, 1993; Campos *et al.*, 2001) and, in the study zone, has manifestly been ignored in many cases. Furthermore, that many riparian strips extends beyond even the higher 50m legal minimum suggests that other, non-legal factors are of at least equal importance. Riparian forest is likely to be intrinsically less susceptible to deforestation, either because it is less susceptible to fire, or (in some cases) because steep topography may make it useless for agriculture or inaccessible and dangerous to livestock. Additionally, landowners may themselves recognize the usefulness of riparian forest in protection of watercourses and, in some cases, provision of dry season grazing which, at least in the medium-term, may be compatible with forest persistence.

In itself, the continued presence and greater robustness of riparian forest is encouraging, as the essentially permanent nature of the causal factors mentioned above suggest that such forest may persist as a long-term landscape element. However, such persistence is also dependent on biological viability. Although the occurrence of ancient, naturally isolated gallery forest remnants, e.g. in the *llanos* of Colombia and upland Belize (Kellman *et al.*, 1994), indicates that that these can be viable in the long-term, this conclusion is not necessarily applicable to narrower, anthropogenic fragments.

The present study suggests Guanacastecan forest cover notably higher than reported in other studies. For example, according to Sader and Joyce (1988), virtually the whole of lowland Guanacaste had been deforested by 1940, whereas Sánchez-Azofeifa *et al.* (2001) reported 3 per cent forest cover (i.e. 284km²) for the Province. These discrepancies reflect various factors, particularly the resolution of the imagery used and stated differences in approach. Sader and Joyce's methods did not permit detection of forest fragments smaller than 55ha. Sánchez-Azofeifa *et al.*'s methodology permitted much higher resolution (3ha), but, according to the authors, was unsuited to detection of tropical dry forest. However, even non-detection only of fragments <3ha, as well as linear fragments of small width but large total areas, is likely to result in forest cover total estimates which fail to take into account biologically significant and visually prominent landscape elements. These considerations suggest that the study zone is less atypical of the rest of Guanacaste than casual comparison with these other studies might suggest. Rather, the landscape elements

and features of the study zone seem rather similar to much of lowland Guanacaste, and suggest that many of the conclusions drawn above are likely to have applicability beyond the study zone. Riparian remnants, particularly, persist throughout seasonally dry Central America, even in some of the most highly deforested zones. Furthermore, this tendency seems to extend beyond the region: Kellman *et al.* (1996) remark that 'gallery forests are by far the most frequent form of natural forest patch in subhumid tropical landscapes', a phenomenon probably largely explicable in many zones by the same combination of factors mentioned above. Such remnants would appear to have an important role to play in biodiversity conservation, particularly outside protected areas, where forest may be otherwise scarce.

4. INHERITANCE, LINKAGE AND NEUTRALITY OF ALLOZYMES OF THE NEOTROPICAL TREE *ANACARDIUM EXCELSUM* (BERTERO & BALBIS EX KUNTH) SKEELS (ANACARDIACEAE)

INTRODUCTION

Anacardium excelsum is a large, evergreen tree native from Honduras in northern Central America, south to Ecuador and the Guyanas (Hartshorn and Gentry, 1991). In seasonally-dry zones, as in the Pacific watershed of Central America, *A. excelsum* is generally found in moist sites less subject to seasonal drought, for example riparian forests. The population genetics of *A. excelsum* is of interest for two main reasons. Firstly, gallery forest appears to represent a large proportion of the remaining closed forest of this largely deforested zone (Chapter Three; Janzen, 1986) As a dominant and common species of such forest (Glander and Nisbett, 1996), the 'genetic health' of *A. excelsum* populations is linked to the conservation status of these habitats. Secondly, information about the population genetics of *A. excelsum* may also generate insights into conservation genetics of other taxa with similar life histories and similar patchy or fragmented distributions.

The suitability of allozymes as genetic markers for population genetics studies is partly due to their codominant inheritance. However, as various factors may cause apparent or real departures from codominance (Gillet and Hattemer, 1989), it should be demonstrated rather than simply assumed. In the present article, enzyme polymorphism and its genetic basis in six enzyme systems of *A. excelsum* are described. Linkage relationships and neutrality of these enzymes are also reported, as these characteristics affect to some degree their usefulness in population genetics studies, e.g. Ritland and Jain's (1981) algorithm for outcrossing rate estimation assumes that loci are independent and selectively neutral between gametic union and point of census. The use of five of these polymorphic loci in studies of population genetics of the species is reported elsewhere (Chapter Five).

MATERIALS AND METHODS

In April 1999, open-pollinated seeds were collected from individual trees in 12 forest fragments located in the *cantón* of Cañas, Guanacaste Province, Costa Rica (see Chapter Five). *A. excelsum* seeds are both dispersed by bats, such as *Artibeus* spp. (Janzen *et al.*, 1976) and fall under gravity when ripe (seeds weigh up to 3g each). The collections were made from the ground below inflorescence-bearing branches of each tree, avoiding areas beneath overlapping crowns of neighbouring trees.

There appear to be no published enzyme extraction or electrophoresis protocols for *A. excelsum*. For this reason, combinations of different extraction buffers, electrophoresis buffer systems and enzymes were screened in order to identify potentially useful markers. The systems AK (adenylate kinase, E.C. number 2.7.4.3), LAP (leucine aminopeptidase, 3.4.11.1), MDH (malate dehydrogenase) (E.C. number 1.1.1.37), PGD (phosphogluconate dehydrogenase, 1.1.1.43), PGM (phosphoglucomutase, 5.3.1.9) and UGPU (UTP-glucose-1-phosphate uridylyltransferase, 2.7.7.9) were selected based on their resolution or consistency with expected quaternary structure. Other enzyme systems were rejected for various reasons: because no activity was detected (alcohol dehydrogenase leaf tissue; β -galactosidase, α -glycerophosphate dehydrogenase, glycerate-2-dehydrogenase, glutamic dehydrogenase, sorbitol dehydrogenase, succinate dehydrogenase, xanthine dehydrogenase), because staining resolution or intensity was unsatisfactory on all tested buffer combinations (aconitase, acid phosphatase, aldolase, diaphorase, fumarase, glucose dehydrogenase, glyceraldehyde-3-phosphate dehydrogenase, isocitric dehydrogenase, malic enzyme, mannose-6-phosphate isomerase, peroxidase), because acceptable resolution or intensity was only inconsistently achieved (esterase), because of interpretation difficulties (superoxide dismutase) or because polymorphy was not detected (alcohol dehydrogenase radicular tissue, catalase, glucose-6-phosphate-dehydrogenase, hexokinase, glucose-phosphate isomerase, shikimic kinase, meniadone reductionase).

The seeds were germinated in greenhouse facilities of the University of Alberta and enzyme extracts prepared by grinding fresh, flash-frozen, leaf disks in Liengsiri *et al.*'s (1990) extraction buffer #9. The homogenate was absorbed on Whatman #3 filter

paper wicks, and stored at -80°C until needed. Horizontal starch (Connaught Laboratories, Ontario) gel electrophoreses were carried out using a pH5.7 histidine-citrate buffer system (Wendel and Weeden, 1990). Allozymes were visualized using staining protocols from Hodgkiss (2001), Liengsiri *et al.* (1990) and Wendel and Weeden (1990). Extraction, grinding and electrophoretic techniques are described in detail elsewhere (Appendix One). The isozymatically invariable *Pinus resinosa* Ait. (Fowler and Morris, 1977) was used as a control. Loci were named in ascending numerical order according to their mobility, whereas alleles of each locus were assigned alphabetic codes, 'A' representing the most common allele.

Analysis of segregation ratios followed the methodology outlined by Gillet and Hattemer (1989). Two specific null hypotheses were tested: that of 1:1 ratios of heterozygous and homozygous progeny of heterozygous mother-trees (e.g. $n_{AB} = n_{AA} + n_{BB}$ for mother-tree AB) and that of 1:1 ratios of progeny genotypes AC and BC in progeny of AB-heterozygous mother-trees (LAP only). The binomial probabilities of observed ratios were calculated. Subsequently, the significance of departure from 1:1 of individual arrays was examined using the Dunn-Šidák method of sequential Bonferroni testing (Sokal and Rohlf, 1995). Under this procedure, the ascendingly ranked probabilities are compared sequentially to a steadily increasing critical value $1-(1-\alpha)^{1/n}$, where α is the chosen probability value (0.05 in this case) and n is the number of arrays not already tested; testing continues until the first nonsignificant array is detected. The significance of departure from 1:1 of the pooled segregation ratios for each enzyme was tested using the G -test with William's adjustment (Sokal and Rohlf, 1995).

Formally, Gillet and Hattemer's methodology requires independent parental genotyping, that is, use of parental germplasm. When parental material is unavailable, as in the present case, putatively heterozygous mother-trees must first be identified on the basis of the presence of two homozygote types in the progeny. As, under dominant inheritance, the heterozygote is indistinguishable from one of the homozygotes, this means of identifying heterozygous parents in itself implicitly assumes codominant inheritance. Consequently, it is possible to test for departure from codominant inheritance only if a logical circularity is admitted. Therefore, the strongest formal

conclusion that can be made is that the segregation ratios of the observed phenotypes are consistent with their proposed genetic interpretations. However, particularly when the phenotypes are consistent with the quaternary structure of the enzyme in question, the degree of confidence attached to the assertion of codominant inheritance does not appear to be critically less than when parental material is available, particularly as the use of adult material for identification of mother-trees heterozygous for alleles expressed in the progeny in itself requires an assumption not necessary in the present case, *i.e.* that of ontogenetic stability.

Linkage disequilibrium

As neither haploid material nor progeny-test data were available, linkage disequilibrium coefficients based on frequencies of coupling and repulsion heterozygotes were not estimable. Therefore, values of Burrows's composite linkage disequilibrium coefficient Δ_j and the corresponding correlation coefficient (Weir, 1979) were estimated for each of the 12 populations. Significant values of Δ_j imply that there are differential frequencies of coupling and repulsion heterozygotes and/or non-random union of gametes (Roberds and Brotschol, 1985). Ohta's (1982) multiple population linkage disequilibrium (D) coefficients were also estimated. Ohta's partitioning of the variance of linkage disequilibrium is useful in elucidating the causes of observed disequilibria: when D_{IS}^2 (the variance component of disequilibrium within a population) is less than D_{ST}^2 (the variance of correlation between different gametes of one subpopulation relative to that of the total population) and $D'_{IS}{}^2$ (variance of within gamete correlation in a subpopulation relative to the total population) is greater than $D'_{ST}{}^2$ (variance of disequilibrium of the total population), then population subdivision (*i.e.* genetic drift) rather than epistatic selection is likely to be the main cause of linkage disequilibria (Kremer and Zanetto, 1997; Ohta, 1982). In the case of the triallelic locus LAP, for estimation of both Burrows's and Ohta's coefficients all alleles except the most common were pooled to a synthetic allele (Kremer and Zanetto, 1997). Missing values were eliminated from the data set as these add no information on non-gametic or gametic correlations in allele frequencies. POPGENE (Yeh and Boyle, 1997) was used for estimation of all the above disequilibrium parameters.

MDH and AK1 were omitted from the above linkage analysis due to interpretation difficulties (see below). However, in the case of AK1, an apparent linkage with PGD was tested for by chi-square analysis of a two-by-three contingency table of AK1 phenotypes (*i.e.* presence or absence of most frequent band, see inheritance results, below) against PGD putative genotypes.

Neutrality

POPGENE was also used to test for selective neutrality. It uses Stewart's algorithm for the Ewens-Watterson neutrality test, as detailed in Manly (1985). One thousand iterations were employed for generation of simulated distributions of the *F* statistic (sum of squared allele frequencies) under the null hypothesis. AK1 and MDH were omitted from the neutrality analysis, for the same reasons indicated above.

RESULTS

Inheritance

Adenylate kinase

The simplest zymograms were characterized by the presence of two invariable bands of unequal intensity, suggesting the presence of two loci (Figure 4-1). As neither band was always present, it was concluded that both loci are polymorphic. When the putative allele AK2-A was absent, there was always a band cathodal to putative allele AK1-A, indicating an overlap in the range of activity of the two loci (Figure 4-2). Other putative AK2 alleles could not be identified, although the possibility of a 'hidden' allele of equal mobility to AK1-A cannot be discounted. The pooled segregation ratio for AK2 (Table 4-1) shows a highly significant homozygote excess. One array (El Cepo 1554) had a probability value smaller than its Dunn-Šidák critical value of 0.004 ($n=26$) (Table 4-2).

Scoring of AK1 was impeded both by inconsistent resolution and the apparent overlap with AK2. Consequently, segregation ratios could not be determined.




Figure 4-1. Adenylate kinase zymogram of *Anacardium occidentale* (3 families), showing putative loci (marked '1' and '2'). 'C' indicates *Pinus resinosa* control.




Figure 4-2. Adenylate kinase zymogram of *Anacardium occidentale* (2 families). 'X' indicates lane with most common AK2 allele absent and intense staining cathodal of most common AK1 band (see text). 'C' indicates *Pinus resinosa* control.

Leucine aminopeptidase

One locus was detected. It followed the expected (May, 1998) monomeric banding pattern. Three putative alleles were observed (Figure 4-3). Overall segregation ratios were not significantly different from expectations for any of the pooled LAP data sets (Table 4-1). The binomial probability of the segregation ratio of one array (Marcela-419) was less than its corresponding Dunn-Šidák value of 0.0014 (Table 4-3). Two progeny arrays (El Ojoche-19, Toronja-107) each contained all three homozygous types.




Figure 4-3. Leucine aminopeptidase zymogram of *Anacardium occidentale*, showing putative genotypes AA (marked '1'), BB (2), AB (3), CC (4), AC (5) in four arrays, and *Pinus resinosa* controls (C).

Malate dehydrogenase

Malate dehydrogenase zymograms generally consisted of an invariant pattern with five clear bands. An occasional variant four-banded phenotype was observed (Figure 4-4). As there was no clear genetic interpretation (see discussion), no segregation analysis was made.




Figure 4-4. Malate dehydrogenase zymogram of *Anacardium occidentale*, showing variant phenotypes (marked V) in two progeny arrays and *Pinus resinosa* controls (C).

Phosphogluconate dehydrogenase

PGD zymograms showed two loci, one of them (PGD2) too indistinct to score. PGD1 displayed the expected dimeric banding pattern, with two loci (Figure 4-5). The putative BB homozygote in general showed two additional faint bands anodal of the main band (Figure 4-5). Although the origin of these bands is not clear, they are easily distinguished from the heterodimer and AA homodimer bands of the putative heterozygote on the basis of their more anodal positions and lower intensities.

The pooled segregation ratio for PGD1 (Table 4-1) shows a highly significant homozygote excess. Two arrays (El Cepo 1511, El Cepo 1566) had probability values smaller than their respective Dunn-Šidák critical values of 0.0031 and 0.0033 (Table 4-4).




Figure 4-5. Phosphogluconate dehydrogenase zymogram of *Anacardium occidentale*, showing putative genotypes AA (marked '1'), BB (2), AB (3) in two progeny arrays, and *Pinus resinosa* controls (C).

Phosphoglucosmutase

PGM is a monomeric enzyme (May, 1998), typically with 2 isozymes (Weeden and Wendel, 1990). Consistent with expectations, the PGM zymograms show two loci, one of them (PGM1) clearly resolved. This locus showed two single-band phenotypes and one double-banded phenotype, scored as two homozygotic types and the corresponding heterozygote (Figure 4-6). Faint, presumably artefactual, banding in the putative B-allele position was generally observed in putative AA genotypes (Figure 4-6). The pooled number of homozygotes was not significantly different from the number of heterozygotes (Table 4-1). The binomial probability of the most divergent ratio (El Cepo-1549, $p = 0.01$) was greater than its corresponding Dunn-Šidák critical value of 0.0013 (Table 4-5).




Figure 4-6. Phosphoglucosmutase zymogram of *A. coelestem*, showing putative genotypes AA (marked '1'), BB (2), AB (3) in three arrays, faint artefactual banding in PGM-B position (BB genotypes marked '2' and those adjacent (right)) and *Pinus resinosa* controls (C)

UTP-glucose-1-phosphate uridylyltransferase

The zymograms show two loci, one of which (UGPU2) was polymorphic (Figure 4-7). The observed banding patterns correspond to expectations for this reportedly monomeric (Chase *et al.*, 1995), *i.e.* one or two bands per lane (Figure 4-7). The number of homozygotes was not significantly different from the number of heterozygotes (Table 4-1). The probability of the most divergent ratio (R.S.R.-5, $p=0.01$) was greater than its corresponding Dunn-Šidák critical value of 0.0013 (Table 4-6).




Figure 4-7. UTP-glucose-1-phosphate uridylyltransferase zymogram of *Anacardium occidentale*, showing putative genotypes AA (marked '1'), BB (2), AB (3) in two progeny arrays, and *Pinus resinosa* controls (C).

Linkage

There were significant ($p < 0.05$) linkage disequilibria in 7 of the 120 permutations of 12 populations and 5 loci (Table 4-7). In three of the populations, there were significant positive disequilibria between AK2-A and UGPU-A; overall, correlations (regardless of significance) between the same pair were positive in ten of the 12 populations, and in no case negative. There were also three significant disequilibria between AK2 and PGD; one of these was negative, but in general correlations between AK2-A and PGD-A were positive. Each of the other significant disequilibria were observed in one population only, with no evidence of consistent trends. There was a strong, highly significant linkage between PGD and AK1 (Figure 4-8, Table 4-8). For all pairwise combinations, D_{IS}^2 was less than D_{ST}^2 and D'_{IS} was greater than D'_{ST} (Table 4-9).

Neutrality

Of the 60 locus and population combinations, observed F was equal to or outside the 95 per cent confidence limits in six cases (AK2 in El Rodeo, LAP in Ojoche and Palmira, PGD and PGM in E.J.N., UGPU in Paso Hondo) (Table 4-10). No locus had observed F consistently close to upper or lower limits.




Figure 4-8. Adenylate kinase (AK) (above) and phosphogluconate dehydrogenase (PGD) (below) zymograms for *A. excelsum* (family Marcela 418), showing apparent linkage (putative PGD genotypes: 1=AA, 2=AB, 3=BB. C indicates *Pinus resinosa* control.

DISCUSSION

Inheritance

With the exception of MDH, the loci studied show banding patterns consistent with their expected quaternary structure. In particular, given this structure, and the putative homozygous genotypes observed for each, they exhibited those heterozygote patterns expected under codominant inheritance. There is no obvious interpretation for the MDH zymograms. *A priori*, a rare variant in otherwise invariable zymograms would be predicted to correspond to a heterozygous genotype. However, in this case the rare phenotype is simpler than the variant, *i.e.* has fewer bands, and even a tentative scoring as heterozygous seems unjustified.

In spite of the evidence of codominant inheritance, there appears nevertheless to be a tendency to homozygote excess, strongest in AK2 and PGD. The presence of three homozygotes in two of the LAP arrays is also unexpected. Although many *A. excelsum* seeds fall directly from the tree, the possibility exists that seed collected from beneath any tree may include extraneous seed dropped by bats or, less commonly, by howler monkeys (Glander, 1979). This could account for the presence of three LAP

homozygotes in two of the progeny arrays. Inclusion of extraneous seed in individual progeny arrays would have two additional consequences, both of which could contribute to the observed homozygote excess. Firstly, >1 homozygote types could occur in progeny arrays of homozygote mothers, leading to incorrect assignment of heterozygous maternal genotypes. This, in turn, is likely to produce apparent segregation distortion. For example, for a diallelic locus with allele frequencies $p_A=0.8$ and $q_B=0.2$, the expected proportions of AA and AB genotypes in progeny of AA homozygotes are 0.8 and 0.2, respectively. In such a case, the inclusion of an extraneous BB 'progeny', whilst leading to redesignation of maternal genotype to AB, would have little effect on the ratio of homozygotes to heterozygotes. Consequently, segregation ratio would be approximately 4:1 instead of the 'expected' 1:1. Secondly, segregation ratios of genuinely heterozygous individuals will depart from 1:1, the magnitude and direction of distortion depending on the admixture rate and the genotype frequencies in admixed seed.

As some applications, e.g. mating system analysis, are dependent on the legitimacy of progeny arrays, the possibility that arrays may contain extraneous seed is of interest and merits further investigation. When numbers of homozygotes exceed numbers of heterozygotes in the population as a whole, inclusion of extraneous seed is expected to lead to homozygote excess in progeny of heterozygous individuals, whereas when overall numbers of heterozygotes exceed overall numbers of homozygotes, admixture is expected to lead to heterozygote excess. If the actual progeny arrays of heterozygous trees have, as expected, equal proportions of heterozygotes and homozygotes, and if admixed seed is drawn randomly from the population as a whole, then, for diallelic loci, the proportion of extraneous seed is:

$$P_e = \frac{f(XX_{(XT)}) - 0.5}{f(XX_{(pop)}) - 0.5},$$

where $f(XX_{(XT)})$ = frequency of homozygotes (or heterozygotes, in case of greater numbers of heterozygotes in the population as a whole) in progeny arrays of heterozygotes and $f(XX_{(pop)})$ = frequency of homozygotes (heterozygotes) in the population as a whole.

Estimates of this parameter for AK2, LAP, PGD, PGM and UGPU for five populations are detailed in Table 4-11. The triallelic locus LAP was included as, in the five populations, observed frequencies of AC, BC and CC genotypes were sufficiently small as not to cause appreciable bias in the use of the above equation. As the equation applies to progeny of heterozygous trees only, individual arrays in which homozygote excess appeared most likely to be caused by incorrect designation of maternal homozygotes as heterozygotes were not included (exceptions noted in Table 4-11). In order to ensure adequate sample size, populations in which there were fewer than three putatively heterozygous trees were excluded, as were cases in which total number of trees sampled was ≤ 5 , as in these cases the ≥ 3 putatively heterozygous trees would themselves largely determine the estimated populational genotypic frequencies.

In seven of the 16 cases (Table 4-11), ratios of homozygotes to heterozygotes in the general population and the collections from putative heterozygotes showed opposite tendencies, implying that the observed imbalances in putative heterozygote collections were not caused by seed admixture. Six of the nine positive estimates of the proportion of admixed seed were between 0.11 and 0.23. There was one estimate > 1 , found in the only case where both ratios showed heterozygote excess. The other two relatively high estimates of proportion of extraneous seed derive from the Congojas population, but estimates for two other loci for the same population were negative. Generalized seed admixture would be expected to be reflected in a positive relationship between denominator and numerator of the above equation. However, the correlation between the two variables is of questionable significance ($r=0.32$, one-tailed probability = 0.12). The evidence for seed admixture appears to be equivocal, suggesting that, if occurring at all, admixture is likely to be sporadic and concentrated on particular trees, *e.g.* trees on bat flight paths and/or close to feeding roosts. This hypothesis accords well with the pattern of homozygote excess in LAP, PGD and UGPU, which appears to be concentrated in specific arrays rather than generalized. In these loci, when arrays with nominally significant departures from 1:1 (*i.e.* probability of ≤ 0.05 under the null hypothesis of 1:1 segregation) are disregarded, the remaining progeny show similar numbers of homozygotes and heterozygotes (204:210, 237:212, 154:146, respectively) and the number of arrays with insignificant homozygote excess is similar to the number

of arrays with insignificant heterozygote excess (13:15, 14:10, 9:7). Under generalized seed admixture, homozygote excess would be expected to predominate even after excluding individually significant arrays, as homozygous genotypes tend to be more common than heterozygous genotypes (*i.e.*, under panmixia, $p^2+q^2 \geq 2pq$). In the case of AK2, on removal of individual nominally significant arrays homozygote excess is no longer significant, but remains marked (205:168) and there are more than twice as many positive as negative departures from 1:1 (14:6). As indicated above, in the case of AK2, the presence of an additional intermediate allele, masked by AK1, cannot be ruled out, and could be responsible for the observed homozygote excess. It should be noted, however, that there are other possible explanations for the homozygote excess, in these and the other loci, *e.g.* meiotic drive (Finkeldey, 1998) or gametic selection.

The effect of inclusion of extraneous seed in progeny arrays on estimation of mating systems parameters can be simulated relatively easily. To do so, sample data sets with known allele frequencies (3 loci), multilocus outcrossing rate (t_m) and maternal inbreeding coefficient F were generated using Ritland's 'datagen' and 'convdata' programmes (Ritland, 1996). Each data set consisted of 20 progeny of each 20 parents. A 'seed pool' data set was also formed by pooling all family arrays and randomizing by multilocus genotype or 'seed'. A programme was then written in Visual Basic for Excel in order to simulate 'seed' movement into the 'sample' arrays from the 'seed pool', as follows. Firstly, the number of sample trees (N) to be affected by the inward migration, and a fixed number (e) of extraneous seed per array were specified. For each run, the sample trees to be affected were randomly chosen and, for each of them, e of their 'real' progeny were overwritten by e multilocus genotypes randomly chosen from the 'seed pool' data set. Values of maternal inbreeding coefficient (F) and t_m (multilocus outcrossing rate) were then recalculated using Ritland's MLTR programme (Ritland, 1996). The simulation programme was run for two initial t_m settings ($t_m \approx 1.0$ and $t_m \approx 0.7$). Within each t_m runs were made for three allele frequency scenarios, and for each allele frequency scenario 5 runs for each of four seed movement scenarios (Table 4-12). Only general, intense seed movement (*i.e.* all 20 trees affected, 50 per cent extraneous composition in each array), particularly for $t_m \approx 0.7$, led to marked overestimation of t_m and gross underestimation of inbreeding coefficients (see cases 2.1-2.3, Table 4.12).

Other movement scenarios (general but moderate movement ($N=20$, $\epsilon=2$), sporadic but intense ($N=2$, $\epsilon=0.5$), and sporadic and moderate ($N=2$, $\epsilon=2$)) had little effect on t_m estimates and intermediate effect on maternal F . In several cases, there were three homozygote types for the triallelic locus in one or more arrays, leading to aborted MLTR execution.

Linkage disequilibrium

With 120 permutations of population and locus-pair, chance (*i.e.* type two error) would be expected to result in six estimates with associated probabilities of 0.05, and would appear to be the most parsimonious explanation for the observed disequilibria involving LAP, which have relatively high probabilities under the null hypothesis and/or inconsistency of sign in different populations (Table 4-7).

The low probability values and high consistency of sign associated with the disequilibria between AK2-UGPU, AK1-PGD and, less convincingly, AK2-and PGD, make the above explanation less likely, and suggest appreciable linkage disequilibria in this tropical angiosperm, a result consistent with findings in temperate broadleaves (Granger, 1996; Huang *et al.*, 1996; Roberds and Brotschol, 1985; Zanetto *et al.*, 1996) and temperate and boreal conifer species (Cheliak and Pitel, 1985; Strauss and Conkle, 1986; Xie *et al.*, 1991; Yang and Yeh, 1993; Yeh *et al.*, 1994).

The lowest reported ploidy number for the *Anacardium* genus is $2n=24$ for *A. occidentale* (cashew) (Mitchell and Mori, 1987; Anonymous, 2002); there appear to be no specific reports for *A. excelsum* (Anonymous, 2002). Assuming that the loci on different chromosomes have equal chances of being selected for study, and assuming that *A. excelsum* has $2n=24$, then the probability of any two of the six loci (including AK1) belonging to the same linkage group is $p= 1-[(12)(11)(10)(9)(8)(7) / 12^6] = 0.77$, indicating that, *a priori*, it is relatively likely that physical linkage is responsible for at least one of the observed disequilibria.

Whether linkage is physical (*i.e.* location on the same chromosome) or non-physical, disequilibrium can be caused either by epistatic selection or genetic drift. The values of Ohta's coefficients (*i.e.* $D_{IS}^2 < D_{ST}^2$, $D'_{IS} > D'_{ST}$) (Table 4-7) suggest that selective forces are not responsible for the observed disequilibria. Non-directional forces such as

founder effects, population subdivision and parental sampling effects (Yeh and Morgan, 1987) all represent possible causal factors. However, the consistency of the direction of the relationship between AK2 and UGPU appears to suggest directional forces, *i.e.* epistatic selection. It should also be noted that, in partially-selfing species such as *A. excelsum*, (see Chapter 5; Ghazoul and McLeish, 2001), the expected or identity disequilibrium is non-zero (Hartl and Clark, 1989), and selfing itself tends to retard decay of disequilibrium (Falconer, 1989, Hartl and Clark, 1989).

For many applications, *e.g.* mating systems studies, the causes of observed disequilibria are of less interest than their magnitude. Close linkage, structural or otherwise, leads to underestimates of outcrossing rates (Brown *et al.*, 1985; Yeh and Morgan, 1987). In the present case, the consistency and magnitude of the AK2-UGPU correlation suggest that only one of each of these locus pairs should be employed in mating system studies.

Neutrality

The results of the Ewens-Watterson test provide no evidence for the presence of single-locus diversifying or balancing selection, suggesting that neither are important at the sampled life stage. The absence of any tendency to heterozygote excess in progeny arrays of putatively heterozygous mothers also suggests absence of balancing selection.

Applications

The loci LAP, PGD, PGM and UGPU appear to be unlinked, selectively neutral and codominantly inherited. As such, they are suitable for applications based on these assumptions. AK2 appears to be unsuitable for mating systems applications, both because of strong linkage with UGPU and the interpretation difficulties noted above. The latter also imply that estimates of measures of genetic variation based on AK2 could be biased downwards, *i.e.* because of possible masking of an additional allele. Although the foregoing analysis suggests that admixture of extraneous seed in progeny arrays, if occurring at all, is sporadic and therefore unlikely to cause large errors in estimates of mating system parameters, caution dictates that such estimates should nevertheless be regarded as tentative.

5. THE EFFECTS OF FOREST FRAGMENTATION ON GENETICS AND REPRODUCTION OF THE TREE *ANACARDIUM EXCELSUM* (BERTERO & BALBIS) SKEELS (ANACARDIACEAE) IN NORTHWESTERN COSTA RICA

INTRODUCTION

Tropical forests were destroyed at a rate of around 15.2 million ha year⁻¹ in the decade 1990-2000 (FAO, 2001). However, deforestation is often not complete; rather, in many cases, one or more tracts within the formerly continuous tree cover remain forested, and are converted by deforestation into fragments set in an unforested matrix (Riitters *et al.*; 2000, Schelhas, 1996). If biologically viable, such fragments may mitigate some negative consequences of deforestation. It follows that the biological implications, including genetic aspects, of the conversion of forest tracts to forest fragments are of considerable relevance to the management and conservation of forests and biodiversity.

The possible effects of forest fragmentation on genetic diversity are complex and interacting. The most immediate of these effects occurs at fragmentation, which, for species formerly present in deforested matrices, leads to reduction in population size. Depending on their size and allele frequencies, reduced populations may not contain all alleles formerly present, *i.e.* they may show founder effects (Meffe and Carroll, 1994; Yeh 2000). Continued low population size is expected to lead to further loss of variation due to random genetic drift (Hartl and Clark, 1989), a process which may be exacerbated by changed post-fragmentation environment. The latter has the potential to cause additional reductions in population size and higher variation in fertility (*e.g.* Aldrich and Hamrick, 1998; Kelly *et al.*, 2000), thereby reducing the ratio N_e/N (effective to census population sizes) (Nunney, 1993; Falconer, 1989). The effects of fragmentation on inter- and intra-population gene flow may exacerbate or mitigate such responses. For example, disturbance-mediated declines in density of tree populations may, due to decline in inter-tree pollinator movements (Karron *et al.*, 1995; Ghazoul *et al.* 1998), lead to increased geitonogamous selfing, which may also be caused by disturbance-mediated changes in pollinator assemblages (Aizen and Feinsinger, 1994). Increased selfing may have an immediate negative impact on fitness, *e.g.* by causing inbreeding depression (Gigord *et al.* 1998), and also

increases susceptibility to drift by further reducing N_e (Yeh 2000). Maintenance of prefragmentation levels of gene flow may, with time, restore variation lost in founder events and may also prevent cumulative drift. Although the proportion of immigrant seed and pollen in a given fragment x may be lower than when the fragment was a tract in continuous forest, at the same time it may originate to a higher degree than previously from fragments located further from x than those extinct pollen and seed sources once present in the matrix. When genetic distance correlates with physical distance, such migrants may be more effective, because of their greater genetic divergence (Mills and Allendorf, 1996). Furthermore, as suggested above, fragmentation and concomitant disturbance may reduce the number of pollination events involving 'home' pollen, implying higher migration rate for a given amount of incoming pollen.

Clearly, the effect of fragmentation on genetic diversity is not easy to predict, particularly given the relatively limited empirical information available on tree responses to fragmentation (see Chapter Two). In the present document, the effects of forest fragmentation on population genetics, flowering, seed size and seedling growth rates of *Anacardium excelsum* are reported. *A. excelsum* is a large, evergreen tree native from Honduras to Ecuador and the Guyanas (Hartshorn and Gentry, 1991). The small, andromonoecious (Mitchell and Mori, 1987) flowers, which Ghazoul and McLeish (2001) reported as partially self-incompatible, are borne on large (up to 50cm length) panicles. In the study zone, trees flower annually between January and April (Cornelius, personal observation); large individuals may produce several hundred panicles. The most common floral visitors tend to be small native bees (*Trigona* spp.) (Ghazoul and McLeish, unpublished). After fertilization, the pedicel expands to a fleshy, twisted hypocarp, which bears a hard drupe at its distal end (Hartshorn and Gentry, 1991; Mitchell and Mori, 1987), which typically matures by early to mid-April (Cornelius, J.P., personal observation). The sweet hypocarps are consumed by bats and howler monkeys (Glander, 1979; Hartshorn and Gentry, 1991; Mitchell and Mori, 1987), which discard or drop the toxic drupes. Many also drop under gravity to the forest floor beneath fruiting trees. Seeds will germinate immediately on moist ground, e.g. on stream sides. Frequently, however, seeds lie in the dry litter until the wet season sets in, i.e. typically no earlier than mid-May. *A. excelsum* seeds are recalcitrant and do not form a long-term soil seed bank (Cornelius, personal observation).

In seasonally-dry zones, as in the Pacific watershed of Central America, *A. excelsum* is generally found in moist sites less subject to seasonal drought, e.g. riparian forests. Its conservation ecology in such sites is of interest for two reasons. Firstly, as a locally common species of naturally clumped distribution, information about the population genetics of *A. excelsum* may generate insights of relevance to other taxa with similar life histories and similarly patchy or fragmented distributions. Secondly, gallery forest appears to represent a large proportion of the remaining closed forest of this largely deforested zone (Chapter Three); as a dominant and common species of such forest, the 'genetic health' of *A. excelsum* populations is linked to the conservation status of these habitats.

METHODS

Study zone and fragments

The study zone is an area of approximately 350 km² located between 10°33' and 10°18' N, 85°02' and 85°12'W in the *cantones* of Cañas and Bagaces, Guanacaste Province, northwestern Costa Rica (Chapter Three; Figure 5-1). Around 95% of the mean annual rainfall of 1693.4 mm (s 459.4) falls between May and November (San Luis, Cañas meteorological station, 1921-1978, MIRENEM, 1988). Altitude varies from 20-200m a.s.l.; mean annual temperature at 95m a.s.l. is 27.5°C (Jiménez *et al.*, 1987). The dry season is characterized by strong (up to 90km hr⁻¹) northerly winds (Coen, 1983) and temperatures up to 37°C. The main land uses in the study zone are industrial agriculture (sugar-cane and rice) and beef cattle ranching. The former predominates on the mollisols, alfisols, vertisols and alluvial inceptisols found south and immediately north of the Interamerican Highway (Figure 5.1a), which cuts through the study zone, whilst ranching predominates on the shallower soils to the north of the highway (Chapter Three). The deforestation of the study zone appears to have occurred mostly during the last 80 years, as a consequence of three main factors: the replacement by exotic pasture species of the semi-open woodland ('sitios') formerly used for grazing, the conversion of closed forest to grassland, and the conversion of woodland to sugar-cane and rice production (Chapter Three). Within the zone, 30 forest remnants containing *A. excelsum* were located using maps, aerial photographs and field exploration. They include non-linear (non-riparian) and linear (mostly riparian) fragments, with wide variation in population size, disturbance, isolation and matrix type (Table 5-1).




Figure 5-1. The study zone and populations in a study of genetic effects of forest fragmentation on *Anacardium excelsum* populations located near Cañas, Guanacaste province, Costa Rica. Top: a. northern sector; bottom: b. southern sector

Genetic variation, gene flow and mating systems

Field and laboratory procedures

Individual trees in each fragment were mapped and their dbh measured. Tree density was selected as a possible explanatory variable for variation in outcrossing rate, because plant density effects on mating systems and pollinator behaviour have been documented previously (Ghazoul *et al.*, 1998; Karron *et al.*, 1995; Murawski and Hamrick, 1991; Murawski and Hamrick, 1992). In the regression analysis of outcrossing rates on density, the following distance-weighted neighbourhood density index (*NDI*) was used:

$$NDI = \frac{\sum_{i=1}^n (t_{25} + 0.5t_{50} + 0.25t_{100})_i}{n}$$

n = number of sampled trees, $t_{25/50/100}$ = number of mature *A. excelsum* trees within 0-25m, 25-50m, 50-100m of each sample tree).

In April 1999, seeds were collected beneath inflorescence-bearing branches of individual trees in 12 fragments (Table 5-2). In larger populations, seed trees were randomly selected, whilst in smaller populations collections were made from all fruiting trees. The collections were made from the forest floor directly beneath the selected trees, avoiding areas beneath overlapping crowns of neighbouring trees. The seeds were germinated in July and August of the same year, and starch gel electrophoresis was carried out on enzyme extracts prepared from seedling leaf tissue. Gels were stained for five codominantly inherited (see Chapter Four) enzyme systems: AK (adenylate kinase, E.C. 2.7.4.3), LAP (leucine aminopeptidase, E.C.3.4.11.1), PGD (phosphogluconate dehydrogenase, E.C. 1.1.1.43), PGM (phosphoglucomutase, E.C. 5.3.1.9) and UGPU (UTP-glucose-1-phosphate uridylyltransferase, 2.7.7.9) (laboratory protocols are detailed in Appendix One). These were selected from a wider group of enzymes based on resolution, polymorphism and consistency with expected quaternary structure (Chapter Four).

Population genetic analysis

Genetic parameters were estimated for maternal and progeny generations. As maternal material was unavailable, maternal genotypes were inferred using the most-likely-parent method (Brown and Allard, 1970), as programmed in Ritland's MLTR program (DOS

version, Ritland, 1996). For both generations, allele frequencies and allelic richness (A) (mean number of alleles per locus) were calculated. As all the sampled loci were polymorphic, A in this case is equivalent to AP (mean number of alleles per polymorphic locus) (Berg and Hamrick, 1997). In order to permit comparison of allelic richness between generations, estimates of progeny allelic richness were then adjusted to the respective maternal sample size using Hurlbert's (1971) rarefaction method, implicitly assuming that population sizes would remain constant over generations. For example, if 10 mother-trees were sampled, progeny allelic richness was rarefied to expected values for 10 sampled progeny). Rarefaction was carried out using Brzustowski's (undated) on-line calculator. Nei's expected heterozygosity ($H_e = 1 - \sum_i p_i^2$, where p_i = frequency of allele i) (Weir, 1996) was also estimated and averaged over all assayed loci. The fixation index F_u ($1 - H_o/H_e$ where H_o = observed heterozygosity), was calculated as a measure of heterozygote deficit or excess with regard to Hardy-Weinberg equilibrium (HWE) expectations. The null hypothesis of HWE within each population was tested using G-tests. In the case of the triallelic locus LAP, all alleles but the most common were pooled to one synthetic class for both the significance testing (to maximize the number of observations per expected genotypic class) and calculation of F_u (for consistency between the measure of disequilibrium and its test statistics). However, due to the low population numbers, in many cases the number of observations in the least common class was still ≤ 5 , usually considered as the threshold for uncritical application of the G-test (Sokal and Rohlf, 1995).

Nine of the 12 sampled fragments fall into two geographic groups, termed here the 'Corobicí group' and the 'Taboga group' (Table 5-1, Figure 5-1). The latter is composed of a set of discrete linear and non-linear fragments set in the sugar-cane matrix of the Taboga estate. It includes the large 'Marcela' fragment and a number of smaller remnants. The Corobicí group fragments are located on the River Corobicí and its tributaries; the matrix is mostly pastureland (Chapter Three). For the populations as a whole, and by these groups, homogeneity tests were carried out to test the null hypothesis of no population differentiation. As for the tests of Hardy-Weinberg equilibrium, and for the same reasons, the less common LAP alleles were pooled to one synthetic allele. Wright's F-statistics (F_{is} , correlation between uniting gametes relative to the subpopulations; F_{it} , correlation between

uniting gametes relative to the population as a whole; F_{st} , correlation between two gametes drawn randomly within subpopulations relative to that of two gametes drawn randomly from the population as a whole) were estimated as:

$$F_{is} = \frac{H_s - H_i}{H_s},$$

$$F_{it} = \frac{H_t - H_i}{H_t},$$

$$F_{st} = \frac{H_t - H_s}{H_t},$$

where H_i = observed average heterozygosity of individuals within subpopulations; H_s =expected average heterozygosity; H_t = total expected heterozygosity (gene diversity) for population as a whole (pooled subpopulations) (Yeh, 2000).

Average historical gene-flow was estimated using the F_{st} method (*i.e.* $mN=0.25(1-F_{st})/F_{st}$), where N =effective population size and m = average migration rate) (Yeh, 2000). Pairwise estimates of Nei's adjusted genetic distance were also calculated and UPGMA dendrograms (Weir, 1996) based on these were generated. Significance of the relationship between pairwise genetic distance and pairwise geographical distance between fragments was tested using the Mantel test (Sokal and Rohlf, 1995) as programmed in Mantel Version 2.0 (Liedloff 1999).

Except where indicated, the above analyses were carried out using the software package POPGENE (Yeh and Boyle, 1997).

Current gene flow

The Paso Hondo population, which was completely sampled (sample size = population size) is monomorphic for UGPU and also lacks LAP-C (see results). In this case, gene flow is estimable as:

$$m = \frac{q_i}{\bar{q}},$$

where m =proportion of immigrant alleles, q_i =allele frequency in the progeny generation, \bar{q} = mean allele frequency in the source population (Hamrick and Nason, 2000). Paso Hondo is a highly isolated population (Figure 5-1, 5-2), with no single obvious source population. Therefore, \bar{q} was set to the average maternal allele frequency over all populations (\bar{q} =0.15, 0.07 for UGPU, LAP-C, respectively).

Gene flow from the outlying tree Bosque Duquesa 706 (Figure 5-2) to the other three sampled trees of this population was also examined (tree 706 is heterozygous for UGPU, whilst the remaining sampled trees are homozygous AA). In this case, gene flow estimates from tree 706 represent maxima, as the allele could be present in the remaining five unsampled trees of the population.

Estimation of mating system parameters

The multilocus outcrossing rate (t_m), average single-locus outcrossing rate (t_s) correlations of outcrossing rate (r_i) and outcrossed paternity (r_p) were estimated by fragment using Ritland's MLTR programme (DOS Version 1.1) (Ritland, 1996), which employs Ritland and Jain's (1981) mixed mating system model. Within MLTR, the likelihood equations were solved using the EM method, and standard errors computed with 1000 bootstraps. Estimates of t less than 1-2S.E. were considered to depart significantly from full outcrossing (Liengsiri *et al.*, 1998); analogous criteria were used for r_p , r_i , *i.e.* estimates greater than 2 SE were considered to differ significantly from zero. For all MLTR analyses, all available parameters were simultaneously estimated. The AK2 locus was omitted, because it appears to be linked with UGPU (Cornelius 2003b); linkage tends to lead to underestimation of outcrossing rate (Brown *et al.*, 1985, Yeh and Morgan, 1987). Two progeny arrays with three LAP homozygotes (see Chapter Four), indicative of more than one contributing maternal parent, were excluded from the data sets. Estimates of individual-tree outcrossing rates were not made because of the relatively small numbers of progeny per maternal family (Table 5-2).




Figure 5-2. Location of Bosque Duquesa tree 706 in relation to other sampled trees (scale approximately 1:7400) (photo: Aerofotos de Costa Rica S.A.)

The expected equilibrium inbreeding coefficient was calculated using the formula:

$$\hat{F}_e = \frac{1 - t_m}{1 + t_m}$$

(Fyfe and Bailey, 1951; Liengsiri *et al.*, 1998). The relative values of \hat{F}_e and observed F_b are of interest as they may help determine whether observed t_m values have shown recent changes, *e.g.* due to fragmentation. Therefore, the relationship between the two variables was examined graphically and by correlation analysis.

Flowering

Field observations of both flowering and fruits production during the 1997 seed collection season suggested that both these fertility parameters varied between fragments. In particular, many trees in fragments located in dry and/or exposed situations (*e.g.* Avispa, Duquesa Arriba, Isla, Salitral, Toronja) were not observed to produce flowers or fruits. This phenomenon is of relevance to the genetic impact of fragmentation, because the mean and variance of fertility are related to population viability both directly (mean), and through their

relationship with effective population size (variance) (Nunney, 1993; Falconer, 1989; Wright, 1938). Accordingly, in March and April 1998, the number of inflorescences visible with binoculars from the ground on trees in each of 24 fragments (Table 5-2) were counted. The large inflorescences of *A. excelsum* persist throughout the flowering and fruiting season. In smaller populations, counts were made on all trees. In large populations, 10 trees were selected randomly (sample sizes are listed in Table 5-2). Observations were not made on trees with dbh < 60 cm as, for purposes of characterizing within-population variation in fertility, variation due to age variation was considered of less interest than fragmentation-induced environmental variation. The mean number of inflorescences per fragment was calculated as an index of fertility. As an index of flowering variability, Shannon's equitability index (J) (Begon *et al.*, 1998) was used:

$$J = \frac{-\sum_{i=1}^S P_i \ln P_i}{\ln S},$$

where P_i = proportion of total number of inflorescences contributed by the i^{th} sample tree; S = number of sample trees. Equitability, which varies from zero to one, is maximal when all individuals produce the same number of inflorescences. Zero values cannot be used in the above formula. This characteristic is of no concern with the commonest ecological application of J , *i.e.* in species equitability. In the present context, however, omission of non-flowering trees would give a misleading impression of fertility variation. For example, if two trees each produce 100 inflorescences, whereas eight produce none, then $J=1$, although gamete production is actually strongly dominated by the two individuals. Because of this, the inflorescence count of each tree was increased by one in order to eliminate zero values.

As there was significant heterogeneity of variance for both the log-transformed and the untransformed mean flowering counts, the null hypothesis of no differences in number of inflorescences tree⁻¹ fragment⁻¹ was tested using the Kruskal-Wallis test for independent samples (Sokal and Rohlf, 1995). Differences between fragments were highly significant (see results). In order to attempt to elucidate the causal factors involved, the following linear regression model was used:

$$\hat{I} = \alpha + \beta \text{type} + \lambda \text{dbh},$$

where \hat{I} = expected mean number of inflorescences for specified values of variables *type* and *dbb*, *type* = 1 for fragments in well-sheltered locations on watercourses, and 0 otherwise (type 1 fragments are indicated in Table 5-2), *dbb* = mean diameter at breast height of sampled trees.

In the case of fertility equitability, interfragment differences cannot be tested because there is only one datum per fragment. Consequently, a regression approach was used with the same model as for the flowering counts. Additionally, a simple linear regression on flowering count itself was also tested. Except where otherwise stated, statistical analyses of these and all other variables (see below) were done with SAS (1999).

Seed weight and growth rate

There are a number of mechanisms by which fragmentation might negatively affect *in situ* or *ex situ* growth rate of seedlings. Firstly, changed levels of conditions and resources, e.g. due to edge effects, may affect growth of regeneration (*in situ*). Secondly, the same factors might affect seed development, which in turn may correlate with seedling growth rate, *ex situ* or *in situ*. Thirdly, increased homozygosity due to drift or inbreeding may result in inbreeding depression (Charlesworth and Charlesworth, 1987; Gigord *et al.*, 1998; Husband and Schemske, 1996; Hedrick and Kalinowski, 2000), again affecting both *in situ* and *ex situ* growth rate.

In order to test for such effects, two common garden experiments were carried out. In April and May 1997, seeds were collected from individual trees in the fragments El Cepo, Canateca, Marcela, Quebrada Reventado (South), Central/South Q. Salitral (bulked), Q. Duquesa/Bosque Duquesa (bulked) and Toronja. Random samples of 55 seed were taken for each by drawing equal numbers of seeds from the individual family lots available for each fragment. The seeds were weighed individually to the nearest centigram. In order to test for fragment effects on seed weight, the data were analyzed by analysis of variance under the following model:

$$Y_{ij} = \mu + \alpha_j + e_{ij},$$

where Y_{ij} = weight of the i^{th} seed of the j^{th} fragment; μ = overall mean; α_j = effect of the j^{th} fragment; e_{ij} = residual associated with the i^{th} seed of the j^{th} fragment. Least square means were estimated by fragment and pairwise testing carried out using Bonferroni-adjusted t -tests.

On 15th July 1997, the seeds were direct sown in a nursery bed in Turrialba, Costa Rica under 50 per cent shade. Plant spacing was 17cm by 17cm and experimental design was randomized complete blocks (five blocks, 11 plants per plot). The experiment was hand-watered as necessary. Shade netting was removed on 18th August 1997. As no border rows was used, and because hand-watering tends to favour central rows, blocks were laid out longitudinally along the 1m wide nursery bed. Germination date for each seed was noted. Total height was measured to the nearest cm on 17th November, *i.e.* four months after sowing. In order to test for fragment effects on four-month height and germination day, analysis of variance under the following mixed model was carried out:

$$Y_{ijk} = \mu + \alpha_j + \beta_k + e_{ijk},$$

where Y_{ijk} = height of the i^{th} seedling of the j^{th} fragment in the k^{th} block; μ = overall mean; α_j = fixed effect of the j^{th} fragment; β_k = random effect of the k^{th} block, e_{ijk} = residual associated with the i^{th} seedling of the j^{th} fragment in the k^{th} block. This reduced model was derived from an initial fuller model including block-by-provenance interaction; the latter effect was dropped because it was insignificant. Least square means were estimated by fragment and pairwise testing carried out using Bonferroni-adjusted t -tests. As there were significant fragment effects on seed weight (see results), an analysis of covariance was also carried out using the same model with the addition of the seed weight covariate. Additionally, germination date was used as covariate in a similar analysis of effects on height growth. Five abnormal seedlings (characterized by shrivelled first leaves, diseased cotyledons or failure to shed testa and release cotyledons and first true leaves) (one in each of the fragments Cepo, Duquesa, Reventado, Salitral and Toronja) were excluded from the analysis of height. Although it is possible that these problems might be genetic in origin, fungal attack or other deterioration after dispersal seem more likely explanations.

The presence of fragment effects on germination percentage was tested using ANOVA under the following model:

$$Y_{ij} = \mu + \alpha_i + \beta_j + e_{ij},$$

where Y_{ij} = germination percentage of seeds of the i th fragment in the j th block, transformed by arcsin; μ = overall mean; α_i = fixed effect of the i th fragment; β_j = random effect of the j th block, e_{ij} = residual associated with the estimate of germination percentage of the i th fragment in the j th block.

A second common-garden test was established on 12 August 1999 in University of Alberta greenhouse facilities. The same fragments assayed for the mating systems study were included (Table 5-2), with the exception of Paso Hondo, for which number of available maternal seedlots was less than the adopted minimum of four per fragment. A randomized complete block design with 10 blocks and single-tree family plots was used. Seedlots of each family were sown in germination trays and transplanted to 12.7cm diameter plastic pots filled with MetroMix™ (Monsanto) after emergence of the hypocotyl but before shedding of the testa. Excess germinants were also transplanted and held in reserve in the same greenhouse. The experimental pots were placed contiguously, *i.e.* without additional space between pots, giving 12.7cm (equivalent to pot diameter) between plants. In 169 of the experimental germinants, it became apparent after shedding of the testa that one or more cotyledons were damaged or destroyed, apparently due to fungal attack during germination or storage. Sixty-seven of these plants died during the three weeks after establishment, and were replaced by randomly selected germinants from the 'reserve'. The remaining 102 damaged plants survived but did not recover normal growth rate. They were therefore excluded from the subsequent analysis of the experiment. In addition, 126 plants had not shed testas by day 23 and were showing no signs of continuing development. Testas of these plants were cut off with nail scissors. Thirty of these plants had damaged or infected cotyledons, and are included in the total of 102 mentioned above, *i.e.* they were excluded from the analysis. The resulting final representation of families by block is detailed in Table 5-3. Total height (81 days, 167 days), diameter at 2cm above the root collar (167 days) and, as a measure of seed size, testa length (mm) after shedding were measured. Least squared

provenance means were estimated and significance of provenance differences tested by analysis of variance under the following mixed linear model:

$$Y_{y(k)l} = \mu + \alpha_{j(k)} + \beta_k + \lambda_l + \delta_{y(k)l},$$

where $Y_{y(k)l}$ value of the response variable on the i^{th} seedling of the j^{th} family of the k^{th} provenance in the l^{th} block; μ = experimental mean, $\alpha_{j(k)}$ = random effect of the j^{th} family in the k^{th} provenance; β_k = fixed effect of the k^{th} provenance; λ_l = random effect of the l^{th} block; $\delta_{y(k)l}$ = random error associated with the ijl^{th} observation. This reduced model was derived from an initial fuller model including block-by-provenance interaction; the latter effect was dropped because it was insignificant for all variables. Similarly to the 1997 study, analysis of covariance was also carried out with date of testa shedding or removal and seed coat length as covariables. The analyses were carried out using SAS under the GLM procedure and the Analyst tool (SAS, 1999). Expected mean squares are included in Table 5-23.

Albino seedlings were noted in three of the sown families (Cepo 1505, 1541 and Ojoche 21). These were replaced with normal seedlings. Additional seed of these families was sown on 2nd December 1999 in order to estimate the proportion of albino seedlings in each case.

RESULTS

Genetic variation and structure

Maternal generation

All alleles of all loci were present in sampled maternal genotypes of the El Cepo, El Ojoche, and River Santa Rosa populations (Table 5-4). Allele LAP-C was absent from the sampled maternal genotypes of populations Bosque Duquesa, Las Congojas, Marcela, Palmira and Paso Hondo. UGPU-A was fixed in samples from La Isla, Paso Hondo y Toronja. Gene diversity (H_j) varied from 0.30 (Paso Hondo) to 0.39 (La Isla), and showed no relationship with population size (Figure 5-3). F_{is} values tended strongly to be negative; significant heterozygote excess was noted in LAP (El Ojoche, Marcela, Palmira), PGD (Cepo), PGM (Cepo, Congojas) (Table 5-4).




Figure 5-3. Scatter plot of log population size (log N) and gene diversity of adult trees in ten populations of *Anacardium occidentale* located in forest fragments in northwestern Costa Rica.

The homogeneity tests (Table 5-5) indicated that the populations as a whole are heterogenous at all studied loci. The overall F_{st} estimate of 0.18 (Table 5-6) suggests moderate to large genetic differentiation (Yeh 2000). Genetic distances (Table 5-7) appeared to be related to geographic distances (Figure 5-4), although the one-tailed p -value of 0.09 for the Mantel test ($Z=142.2$, $G=1.04$) was suggestive rather than conclusive. Genetic distances between the Paso Hondo and other populations were much higher than the others (Table 5-7). On removing this population, the relationship between genetic and geographical distances became strongly significant ($Z=78.19$, $G=2.18$, $r=0.39$, one-tailed $p=0.015$). Homogeneity tests (Table 5-5) and F_{st} (Table 5-6) estimates by group suggest similar, low-to-moderate genetic differentiation within the Taboga and Corobicí groups. UPGMA dendrograms for the 10 populations and two groups are illustrated in Figure 5-5 to 5-7.




Figure 5-4. Scatter plot of pairwise relationship between geographic and Nei's unbiased genetic distances in 10 populations of *Anacardium excelsum* located in northwestern Costa Rica (based on inferred maternal genotypes)




Figure 5-5. UPGMA dendrogram based on Nei's unbiased genetic distance between maternal trees in ten populations of *A. excelsum* located in northwestern Costa Rica.




Figure 5-6. UPGMA dendrogram based on Nei's unbiased genetic distance between maternal trees in four populations of *A. excelsum* (Corobici group) located in northwestern Costa Rica.




Figure 5-7. UPGMA dendrogram based on Nei's unbiased genetic distance between maternal trees in three populations of *A. excelsum* (Taboga group) located in northwestern Costa Rica.

Progeny generation

All alleles of all loci were present in progeny genotypes of all populations except Bosque Duquesa (LAP-C), El Rodeo (LAP-C, UGPU-B) and EJN (LAP-C, UGPU-B) (Table 5-8). When rarefied to the maternal sample size, progeny allelic richness was significantly lower than maternal allelic richness (signed rank test: sum of positive ranks = 51.5, sum of negative ranks = 3.5, $p < 0.02$). Gene diversity (H_s) varied from 0.20 (El Rodeo) to 0.41 (Palmira) and was not linearly related to log population size (Figure 5-8). F_{is} values tended strongly to be positive; significant homozygote excess was noted in at least one locus of all populations, and all loci showed significant homozygote excess in at least one population. Mean values ranged from 0.06 (El Ojoche) to 0.27 (Paso Hondo) (Table 5-8).




Figure 5-8. Scatter plot of log population size (log N) and gene diversity of progeny in twelve populations of *Anacardium occidentale* located in forest fragments in northwestern Costa Rica.

The homogeneity tests indicate that the populations as a whole are heterogenous at all studied loci (Table 5-9). The overall F_{st} estimate of 0.18 (Table 5-10) suggests moderate to large genetic differentiation (Yeh 2000).

Genetic distances (Table 5-11) appeared to be related to geographic distances (Figure 5-9), although the Mantel test statistic (Figure 5-9) was not significant. Genetic distances between the Paso Hondo and other populations were unusually high, but the relationship between genetic and log geographic distance was not made stronger or more significant by removing this population. Homogeneity tests (Table 5-9) and F_{st} (Table 5-10) estimates by group suggest rather greater genetic differentiation within the Corobicí group than the Taboga group. UPGMA dendrograms for all 12 sampled populations and two groups are illustrated in Figures 5-10 to 5-12.




Figure 5-9. Scatter plot of pairwise relationship between geographic and Nei's unbiased genetic distances in 10 populations of *Anacardium occidentale* located in northwestern Costa Rica (progeny genotypes)




Figure 5-10. UPGMA dendrogram based on Nei's unbiased genetic distance between progeny in 12 populations of *Anacardium occidentale* located in northwestern Costa Rica.




Figure 5-11. UPGMA dendrogram based on Nei's unbiased genetic distance between progeny in four populations of *A. excelsum* (Corobicí group) located in northwestern Costa Rica.




Figure 5-12. UPGMA dendrogram based on Nei's unbiased genetic distance between progeny in three populations of *Anacardium excelsum* (Taboga group) located in northwestern Costa Rica.

Gene flow

For maternal and progeny data, mean estimates of gene flow based on the F_{st} method were around 1 individual generation⁻¹ for the populations as a whole, but from 2-4 individuals generation⁻¹ within the Corobicí and Taboga groups (Tables 5-6, 5-10).

The mean estimated migration rate into the Paso Hondo population was 0.17 (Table 5-12). There was zero estimated gene flow from Bosque Duquesa Tree 706 to the other three sampled trees, *i.e.* UGPU-B was absent from their progeny (Table 5-8).

Mating system parameters

Estimates of t_m (Table 5-13) ranged from 0.266 (Palmira) to 0.744 (El Ojoche). All estimates departed significantly from 100 per cent outcrossing. Estimates of t_s ranged from 0.258 (Palmira) to 0.726 (Marcela), but were generally higher than corresponding multilocus estimates. All estimates departed significantly from 100 per cent outcrossing, except for La Isla and Paso Hondo. Correlation of outcrossing rate (r_s) were low, ranging from 0.086

(Santa Rosa) to 0.230 (Palmira), but significant in six cases. Correlation of outcrossed paternity (r_p) ranged from 0.418 (Bosque Duquesa) to 0.980 (Toronja), and departed significantly from zero in all cases. \hat{F}_e ranged from 0.15 (Ojoche) to 0.58 (Palmira) and appeared to be positively linearly correlated with observed F_{is} (Figure 5-13). In seven cases \hat{F}_e was greater than F_{is} ; the latter was marginally greater in the three fragments where this was not the case.




Figure 5-13. Scatter plot of observed inbreeding coefficients (F_{is}) and equilibrium inbreeding coefficients (F_e) based on estimated outcrossing rates ($\hat{F}_e = (1-t_m)/(1+t_m)$) (one-tailed) in populations of *Anacardium occidentale* located in 10 forest fragments in northwestern Costa Rica (p -value is one-tailed)

Multilocus outcrossing rate was significantly and strongly positively related to population neighbourhood density mean ($t_m = 0.4182 + 0.0071ndi$, $F = 6.83$, $p = 0.0309$, $R^2 = 0.461$) (Figure 5-14).

Flowering

Fragment mean flowering indices varied from 1.9 tree⁻¹ to 211 tree⁻¹ (Table 5-14). There were highly significant differences between populations in tree flowering indices (Kruskal-Wallis test, chi-square = 83.4, df=23, $p < 0.0001$). Both mean dbh and fragment type had positive and significant effects on mean flowering index (Table 5-15).




Figure 5-14. Linear regression of multilocus population outcrossing rates on population mean neighborhood density index (see text) in ten populations of *Anacardium excelsum* located in northwestern Costa Rica

Population equitability (J) values varied from 0.17 to 0.85 (Table 5-14). In several cases, one tree produced >50% of the total inflorescence count. J was significantly positively affected by *type*, but not by dbh mean or standard deviation (Table 5-16). J was significantly correlated with mean flowering index (Figure 5-15).

There was a highly significant effect of origin (fragment) on seed weight (Table 5-17), mean weights ranging from 1.72g (Duquesa) to 2.59g (Cepo). (Table 5-18). Mean seed weights in the El Cepo and Canateca fragments were significantly higher than those of the other fragments. There were no significant differences between the Toronja, Salitral and Duquesa fragments, and these had significantly lower weights than all the other fragments apart from Quebrada Reventado (Table 5-18).




Figure 5-15. Scatter plot of flowering equitability and mean flowering index in 24 populations of *Anacardium occidentale* located in forest fragments in northwestern Costa Rica.

Common garden experiments

1997 experiment

The overall germination rate was 45 percent. There was a highly significant fragment effect on arcsin germination percentage (Table 5-19). Germination percentage varied from 22 percent (Salitral) to 78 percent (Marcela) (Table 5-20).

The mean height at four months was 27.0 cm ($s=7.31$ cm). ANOVA revealed highly significant fragment effects on four month height (Table 5-19); least square means varied from 22.1cm (Toronja) to 31.5cm (Marcela) (Table 5-20).

Mean time to germination was 15.5 days ($s=2.22$); germination day was significantly affected by fragment of origin (Table 5-19); least square mean germination day varied from 14.1 (Reventado) to 16.0 (Duquesa, Salitral) (Table 5-20). Seed weight and germination day were respectively significant and highly significant as covariates, but a strongly significant fragment effect on growth rate remained (Table 5-21). Least squares height means for the covariance analysis are included in Table 5-20.

1999 experiment

Analysis of variance revealed a significant effect of provenance (fragment) on height and diameter at both measurement ages (Table 5-22) and on testa length. The relationships of least square means of the three growth variables (Table 5-22) with outcrossing rate are illustrated in Figures 5-16 to 5-18.




Figure 5-16 Scatter plot of least squares mean height at 81 days (based on ANOVA) and estimated outcrossing rate in populations of *Anacardium excelsum* from nine forest fragments in northwestern Costa Rica (p-value is one-tailed)




Figure 5-17. Scatter plot of least square mean height at 167 days (based on ANOVA) and estimated outcrossing rate in populations of *Anacardium excelsum* from nine forest fragments in northwestern Costa Rica (p-value is one-tailed)




Figure 5-18. Scatter plot of least square mean diameter at 167 days (based on ANOVA) and estimated outcrossing rate in populations of *Anacardium occidentale* from nine forest fragments in northwestern Costa Rica (p-value is one-tailed)

In the analysis of covariance, both covariates were highly significant for all three variables (Table 5-24). Significance of fragment effects persisted, and least squares means generated under the covariance analysis (Table 5-25) appeared to be more strongly related to outcrossing rate (Figures 5-19 to 5-21).




Figure 5-19. Scatter plot of least square mean height at 81 days (based on ANCOVA) and estimated outcrossing rate in populations of *Anacardium occidentale* from nine forest fragments in northwestern Costa Rica (p-value is one-tailed)




Figure 5-20. Scatter plot of least square mean height at 167 days (based on ANCOVA) and estimated outcrossing rate in populations of *Anacardium occidentale* from nine forest fragments in northwestern Costa Rica (p-value is one-tailed)




Figure 5-21. Scatter plot of least square mean diameter at 167 days (based on ANCOVA) and estimated outcrossing rate in populations of *Anacardium occidentale* from nine forest fragments in northwestern Costa Rica (p-value is one-tailed)

The numbers of albinos produced by families 1505, 1541 and Ojoche 21 are detailed in Table 5-26. The segregation ratio for Ojoche 21 (31:13) was not significantly different from 3:1 ($G=0.41, G_{crit, p=0.05} = 3.84$). Albino seedlings from this seedlot are pictured in Figure 5-22.



Figure 5-22. Albino progeny of *Anacardium excelsum* tree 21, El Ojoche population, northwestern Costa Rica

DISCUSSION

Comparison of levels of within-population gene diversity shown by *A. excelsum* with those of published meta-analyses is not straightforward, as the latter (e.g. 0.13 ± 0.011 for native tropical woody taxa, Loveless, 1992) are inclusive of monomorphic loci. However, as Loveless reported percentage of loci polymorphic (P) for the same group of 39.0 per cent, this implies mean gene diversity of polymorphic loci for this group of $(0.13/0.39) = 0.33$, i.e. closely similar to the estimates (means of 0.34, 0.33 in maternal and progeny generations) reported here. Similarly, the estimates for *A. excelsum* are comparable to other estimates based on polymorphic allozyme loci of neotropical trees, e.g. for *Enterolobium cyclocarpum* ($H_e = 0.365$, five loci, Rocha and Lobo, 1996), *Pithecellobium elegans* ($H_e = 0.31$, recalculated for 6 polymorphic loci, Hall *et al.*, 1994a), *Pentaclethra macroloba* ($H_e = 0.21$, three loci of adult trees, Hall *et al.*, 1994b), *Carapa guianensis* ($H_e = 0.31$, 6 polymorphic loci, Hall *et al.* 1994c). There is, at least, no indication from the data - i.e. genetic variation at polymorphic loci - that *A. excelsum* shows any general tendency to low levels of within-population gene diversity.

Nevertheless, the results of the present study indicate clearly that forest fragmentation threatens this *status quo*, essentially due to the action of several mutually reinforcing processes, i.e. founder effects, declining and variable fertility, higher inbreeding and lower

growth rates. These are considered below. Subsequently, the potential of gene flow and selection to mitigate these factors is discussed.

Founder effects

The reduced allelic richness of the completely sampled Paso Hondo population (maternal generation) provides the clearest evidence of founder effects: two of the alleles present in the group of populations as a whole are absent in the four mother-trees which make up the population. Although records suggest that this population has been isolated since at least the first part of the last century (see Chapter Three), the large size of the trees (mean dbh 151.3cm) implies that all four trees derive from prefragmentation mating events, *i.e.* that cumulative drift is unlikely to be responsible. Founder effects also seem to be responsible for the notable genetic divergence of the population, as quantified in the genetic distance matrices.

In the case of the other populations (excepting Palmira, which was completely sampled, and from which an allele (LAP-C) has also been lost) progeny data are more informative, as they sample more completely the populations in question. Three populations (Bosque Duquesa, El Rodeo, and E.J.N.), exhibit progeny allelic richness lower than the maximum of $A=2.2$. In the case of Bosque Duquesa, it seems probable that the estimate of $A=2.0$ reflects loss of LAP-C from the population, in which >50 per cent of adult trees were sampled. Sampling intensity was lower in the case of E.J.N. and El Rodeo, suggesting that sampling effects, rather than founder effects, may be responsible in these cases.

Flowering and seed weight

The strong effect of fragment type on flowering fertility suggests resource limitation of flowering in some fragments. The three fragments with particularly low levels (COSTASEM, Las Avispas, Quebrada Salitral) are all located in the largely treeless and flat sugar-cane and rice matrix of the southern part of the study zone. These fragments lack protection from exposure; Las Avispas also appears unusually dry for *A. excelsum* habitat, presumably due to the drainage-works in the surrounding cane-fields.

In many cases, reduced flowering may imply lower seed production, independently of whether there are subsequent additional (pollinator or resource) limitations on the latter.

Although no formal study of seed production was made, the expectation of reduced seed production is born out by experience in seed collection in the southern part of the study zone. For example, in the 1997 collection season, mean numbers of seed per tree collected in the Avispa, Duquesa Arriba, Isla and Salitral Central fragments ranged from 2.0-4.4. By contrast, in the large riparian fragments north of the Interamerican highway, numbers of seed collected were in general an order of magnitude greater (in practice, a collection ceiling of 30-50 seeds tree⁻¹ was set. Ghazoul and McLeish's (2001) findings on seed production in nine of these fragments are consistent with these observations. They found notably reduced mature seed production in the Avispa and Paso Hondo ('Tres Espavels') fragments.

The results suggest that the same factors that give rise to declining flowering are also likely to increase variance in flowering fertility, *i.e.* deteriorating conditions and resource levels, under which only a few trees are able to produce normal numbers of seed and flowers, leading to dominance of reproduction by one or a few trees. A similar 'secondary bottleneck' in response to fragmentation has been reported by Aldrich and Hamrick (1998) in *Symphonia globulifera*. Both low mean fertility and such bottlenecks tend both to exacerbate initial founder effects and predispose populations to ongoing drift. Low mean fertility would tend to cause continuing population decline, or at least would inhibit population growth, whilst high variance in fertility tends to be the most important cause of N_e being less than N (Falconer (1989).

Fragment effects on seed weight and size appear to parallel the effects on flowering; with the single exception of the Marcela fragment (for seed weight), degraded, exposed and/or dry fragments have the lowest seed weights (Duquesa, Salitral, Toronja) and testa lengths (Bosque Duquesa, Toronja, Palmira). Clearly, this suggests that the same resource limitations that affect flowering may also have independent effects on seed development.

Inbreeding

The observed relationship between neighbourhood density and outcrossing implies that pollinator movement is limited by distance. This is indirectly supported by the high correlations of paternity of outcrossed progeny (suggesting that a few neighbouring trees may be responsible), by the current (zero) gene flow estimate for the Bosque Duquesa population, and in Ghazoul and McLeish's (2001) data on insect visitation to flowers in five

of these fragments. When all five fragments in common are considered, there is a positive but insignificant correlation between t_m and number of insect visits per unit time ($r=0.43$, $p=0.47$, $df=3$). However, the Paso Hondo (Tres Espavels) fragment appears to have a markedly different flower visitor assemblage to all the other fragments, with 17 per cent of visits by *Trigona* bees, against >50 percent in the other fragments. When this fragment is disregarded, there is a significant, near perfect correlation between insect visitation frequency and t_m ($r=0.96$, $p=0.04$, $df=2$).

The relationship between outcrossing rate and neighbourhood density has implications for the effect of forest fragmentation on genetics of *A. excelsum*, as reduction of neighbourhood density in *A. excelsum* is an expected consequence of fragmentation in three related scenarios. First, in riparian fragments, as the width of the gallery forest 'buffer' is reduced, the forest is effectively redimensioned from plane to line, with a corresponding reduction in neighbourhood density. This effect can be particularly strong when one river bank is completely deforested, as is common in the study zone. Secondly, conditions in the disturbed fragments may favour other species at the expense of *A. excelsum*, e.g. initially, successional earlier species, and, subsequently, species better suited to the drier, hotter conditions. Finally, as demonstrated here, post-fragmentation conditions may be inimicable to flowering, seed production, seed development and seedling growth (see below) and, therefore, for all four reasons, to recruitment. In effect, there is a conversion to marginal habitat, which is expected to offer fewer suitable microsites and higher probability of competitive exclusion (Yeh and Layton, 1979) and will therefore lead to lower stocking per unit area.

The segregation ratio for albinism in tree Ojoche 21 is strongly suggestive of selfing, or near selfing, of an individual heterozygous for the locus controlling this trait. Sporadic occurrence of albinism in other families may be caused either by outcrossing events between heterozygous individuals or lower degrees of self-fertilization of heterozygous individuals. The presence of this lethal allele in two of the populations demonstrates that inbreeding, particularly selfing, in *A. excelsum* can have an adverse effect on fitness. Ghazoul and McLeish (2001) also found that selfed seed of the species had higher abortion rates. It

follows that the higher selfing rates likely to be caused by fragmentation-mediated density reduction may directly to reduced viability fitness.

The observed positive relationship between F_{is} and \hat{F}_s suggests that increased selfing is already affecting genetic variation in these fragments. At the same time, the tendency for \hat{F}_s to be higher than F_{is} indicates that equilibrium values have not yet been reached, and therefore that the most serious effects of increased inbreeding have yet to be manifested.

It is worth noting that at least some of the outcrossing estimates are surprisingly low, and are suggestive of complete self-compatibility, as reported by Freitas and Paxton (1996) for cashew (*A. occidentale*), rather than the partial self-incompatibility reported by Ghazoul and McLeish (2001). Roubik (1995) also reported self-compatibility in *A. giganteum*.

Seedling growth rate

The analyses suggest that the observed variation in growth rate between seedlings from different fragments is partly due to variation in seed weight and size, implying that the probable causal factors implicated in the latter (*i.e.* resource limitation) also affect on seedling growth rate. This factor may be a sufficient explanation for the similarity in the results of the two trials, *i.e.* the poor performance in both of the Toronja and Duquesa sources. The superiority in both trials of plants from the Marcela fragment cannot be explained in these terms. It is possible that lower inbreeding depression might be one cause of the superiority of the Marcela source, which is the largest of the non-riparian populations and has relatively high t_m and relatively low r_p (suggesting a low tendency to biparental inbreeding). However, as the relationship between outcrossing rate and growth in the second trial is, at strongest (Figure 5-19), no more than suggestive, this explanation can be no more than tentative. The lack of any clear relationship between growth rate and outcrossing rate may imply that inbreeding depression is expressed predominantly at earlier life stages, *e.g.* immediately after fertilization, as observations by Ghazoul and McLeish (2001) suggest.

In summary, *A. excelsum* populations in the study fragments show, variously, evidence of founder effects, excess homozygosity in progeny, lower (rarefied) allelic richness in progeny than adults, 'secondary bottlenecks' related to fragment conditions and higher selfing rates due to lower density. The possible roles of gene flow and selection in mitigating these problems are now considered.

Gene flow

Gene flow between populations tends to result in increased intra-population genetic variation. It has the potential to restore genetic variation lost in founder effects, to counteract cumulative drift, and to reduce selfing (because seed produced from immigrant pollen is necessarily outcrossed). One such effect can be seen directly in the present study, *i.e.* in the 'reintroduction' of LAP-C and both UGPU-B and LAP-C to, respectively, Palmira and Paso Hondo (these alleles are absent in maternal generations, but present at low frequency in the progeny, due to gene flow). However, without substantial increases in population sizes, these alleles are unlikely to be present in actual progeny generations (consistent with this, they are not reflected in rarefied allelic richness).

Theoretically, the low-to-moderate degree of subpopulation-within-group differentiation suggests that gene flow at this level has been substantial, *i.e.* 3-4 migrants generation⁻¹. However, there are grounds for caution in concluding from these data that gene flow will be sufficient to mitigate the negative impacts mentioned above. Firstly, gene flow between the populations as a whole appears to have been notably less (as evinced by the high overall F_{st} values). The estimate of $F_{st}=0.18$ is higher than average G_{st} values for comparable species, *e.g.* long-lived woody species in general (0.084 ± 0.008 , Hamrick *et al.*, 1992), and, within this group, mixed breeding system, animal-pollinated species (0.122 ± 0.038), gravity-attached seed dispersal (0.099 ± 0.024), and much higher than Loveless's (1992) mean value for biotically dispersed tropical species (0.050 ± 0.008). The plots of genetic *v.* geographic distance reveal obvious relationships, disrupted by individual exceptional pairwise combinations, *i.e.* those due to the Paso Hondo population in the case of the maternal generation, and, in the case of the progeny, the Ojoche-Marcela pairwise observation, which has geographic and genetic distances of 21km and 0.008, respectively (Figure 5-6). Correlations between genetic and geographic distance imply isolation-by distance (Yeh, 2000). This may reflect both that, as

expected, pollinators should forage no further afield than necessary to meet their energy requirements, and inability of a relatively weak-flying pollinator (*i.e.* *Trigona* spp.) to traverse larger distances. Either way, increasing distance between *A. excelsum* fragments would tend to promote shifts to foraging on alternative species rather than energy-expensive moves between widely separated fragments. Low levels of flowering in some fragments, as observed here, would also be expected to reduce probability of visitation by non-resident pollinators. Ghazoul and McLeish's (2001) finding of greatly reduced *Trigona* visitation in the isolated Paso Hondo fragment is consistent with this expectation. The presence of IBD suggests also that diaspore vectors are subject to similar constraints. Field studies of bat foraging in the study zone support this, e.g. Heithaus *et al.*'s (1975) found that although individuals of *Artibeus* and other taxa ranged relatively widely along the River Corobicí, none were recaptured in sample points 4km northwest, on the River Tenorito. Clearly, individuals also carrying fruits with seeds still attached would be still rarer.

Secondly, even within the two groups, the UPGMA grouping is consistent with geographical expectations, suggesting presence of IBD also at this smaller geographic scale. The fragments Marcela and Duquesa group separately from the Toronja fragment, from which they are separated by a range of low hills (Las Lomas). The Cepo and Santa Rosa populations, which were connected by continuous riparian forest until the construction of the dam on the River Santa Rosa, group separately from the two non-linear fragments Ojoche and Congojas.

Thirdly, findings on current gene flow suggest that immigration rate may be idiosyncratic to particular fragments. The low-flowering Bosque Duquesa fragment shows no evidence of inward gene flow. The Paso Hondo fragment, by contrast, appears to be subject to substantial current gene flow; the migration rate estimate of 0.17 implies that 34% of pollen is of immigrant origin. However, even in this case, mN would still be relatively small, even if $N_e=N$.

Selection

F_{IS} estimates in maternal generations, which tend to be negative, are in marked contrast to those for the progeny generations, which tend to show homozygote excess. It is possible that this may reflect either heterozygote advantage or selection against deleterious

recessives. Both the latter and, in the case of asymmetrical overdominance, the former (Young *et al.*, 1996), may result in purging of genetic load. Such purging, although in itself tending to increase homozygosity, would tend to mitigate immediate effects on fitness of increased inbreeding. However, studies suggest that purging does not act consistently in natural populations, possibly because slightly deleterious alleles may principally be responsible for genetic load (Byers and Waller, 1999). Furthermore, purging would not prevent loss of currently neutral variation, which in changed environmental conditions may become adaptive.

As a whole, the results of this study suggest a species rather ill-equipped to deal with the threat posed by habitat fragmentation. This is well exemplified by the case of the Bosque Duquesa fragment, which shows loss of variation due to founder effects, combined with at least five other factors expected to lead to continuing decline in census or effective population sizes (reduced flowering, low seed weight, slower seedling growth rate, high variance in fertility, low outcrossing), whilst there is little or no mitigating immigration. Although, as suggested by current levels of gene diversity, the impact of forest fragmentation on genetic variation of *A. excelsum* appears to be largely incipient rather than yet realized, it is not clear that either gene flow or selection will be able to obviate the likely adverse consequences.

A. excelsum shows fewest signs of fragmentation effects under the least disturbed conditions, and it seems probable that in large, relatively undisturbed riparian fragments the species will continue to persist indefinitely. However, the risk posed by disturbance of such habitat, whether caused by fragmentation or other factors such as grazing or damming (as in the River Santa Rosa fragment), remains clear.

6. INHERITANCE, LINKAGE AND NEUTRALITY OF ALLOZYMES OF THE NEOTROPICAL TREE *PLUMERIA RUBRA* L. (APOCYNACEAE)

INTRODUCTION

Plumeria rubra L. (calichuche, flor blanca, frangipani, sacuanjoche) is a small to medium sized deciduous tree. It occurs naturally from Mexico to Panama (Standley and Williams, 1969; Woodson, 1938), where it is found principally on sea-cliffs, limestone outcrops, and canyon sides within seasonally dry zones (Haber, 1984; Pittier, 1978). The hermaphroditic flowers are pollinated by hawkmoths (Haber and Frankie, 1989). The winged seeds appear to be selected primarily for wind dispersal, although, as trees often overhang watercourses, dispersal in streamflow also seems inevitable.

The natural range of *P. rubra* includes much of the Pacific watershed of Central America, where cattle-raising and agriculture have replaced all but approximately 8 per cent of the original dry forest vegetation (Janzen, 1986). Outside protected areas, a large proportion of the remaining forest appears to be concentrated in linear riparian remnants of various types and degrees of degradation (see Chapter Three). The population genetics of the constituent species of these strips, such as *P. rubra*, are of interest for two principal reasons. Firstly, the 'genetic health' of such populations partly determines their ecological viability. Secondly, information on the population genetics of such species may generate insights into conservation genetics of other taxa with similar life histories and similarly patchy or fragmented distributions.

The suitability of allozymes as genetic markers for population genetics studies is partly due to their codominant inheritance. However, as various factors may cause apparent or real departures from codominance (Gillet and Hattemer, 1989; Strauss and Conkle, 1986), it should be established rather than simply assumed. In the present article, enzyme polymorphism and its genetic basis in five enzyme systems of *P. rubra* are described. Linkage relationships and neutrality of these enzymes are also reported, as these characteristics affect to some degree their usefulness in population genetics studies, e.g. Ritland and Jain's (1981) algorithm for outcrossing rate estimation assumes

that loci are independent and selectively neutral between gametic union and point of census. The use of the selected markers in the characterization of the impact of forest fragmentation on seven *P. rubra* populations is described separately (see Chapter Seven).

MATERIALS AND METHODS

Field and laboratory procedures

In February to April of 1997, 1998 and 1999 open-pollinated seeds were collected from individual trees in 7 forest remnants located in the *canton* of Cañas, Guanacaste Province, Costa Rica (see Chapter Seven). Ripe capsules were collected directly from the trees using a pruning pole and attached basket. Seeds were extracted manually and stored until needed.

There appear to be no published enzyme extraction or electrophoresis protocols for *P. rubra*. For this reason, combinations of different extraction buffers, electrophoresis buffer systems and enzymes were screened in order to identify potentially useful markers. The systems reported (aspartate aminotransferase (AAT, E.C. 2.6.1.1), alcohol dehydrogenase (ADH, E.C. 1.1.1.1), glucose-phosphate isomerase (PGI, E.C. 5.3.1.9), phosphoglucomutase (PGM, E.C. 5.4.2.2) and phosphogluconate dehydrogenase, (PGD, E.C. 1.1.1.43)) were selected based on their clarity of resolution or consistency with expected quaternary structure. Other enzyme systems were rejected for various reasons: because staining resolution or intensity was unsatisfactory on all tested buffer combinations (glutamic dehydrogenase, glucose dehydrogenase, isocitric dehydrogenase), because acceptable resolution or intensity was only inconsistently achieved (adenylate kinase, esterase, leucine aminopeptidase, malic enzyme, shikimic kinase), because of interpretation difficulties (malate dehydrogenase) or because polymorphy was not detected (diaphorase, meniadone reductionase).

Extracts of the five enzymes were prepared by crushing recently emerged radicles in Liengsiri *et al.*'s (1990) extraction buffer #9. The homogenate was absorbed to Whatman #3 filter paper wicks, and stored at -80°C until needed. Horizontal starch (Connaught Laboratories, Ontario) gel electrophoreses were carried out using pH7.0 histidine-tris (Pitel and Cheliak, 1984)) (ADH, PGM, PGD) and pH8.1 lithium borate / tris citrate

(Ridgeway *et al.*, 1970) (AAT, PGI) buffer systems. Allozymes were visualized using staining protocols from Liengsiri *et al.* (1990) and Wendel and Weeden (1990). Laboratory procedures are described in detail elsewhere (see Appendix One). The isozymatically invariable *Pinus resinosa* Ait. (Fowler and Morris, 1977) was used as a control. Loci were named in ascending numerical order according to their mobility, whereas alleles of each locus were assigned alphabetic codes, 'A' representing the most common allele.

Inheritance

Segregation analysis followed the methodology outlined by Gillet and Hattemer (1989). The following specific null hypotheses, both corresponding to codominant allelic action, were tested here on progeny arrays of putatively heterozygous mother-trees: (i) for maternal genotype XY ($X \neq Y$), $n_{XY} = n_{XX} + n_{YY}$; where n =numbers of progeny; (ii) for maternal genotype XY, $n_{XZ} = n_{YZ}$, ($Z \neq X, Y$) (AAT, PGM). The binomial probabilities of observed ratios were calculated. Subsequently, the significance of departure from 1:1 of individual arrays was examined using the Dunn-Šidák method of sequential Bonferroni testing (Sokal and Rohlf, 1995). Under this procedure, the ascendingly ranked probabilities are compared sequentially to a steadily increasing critical value $1-(1-\alpha)^{1/n}$, where α is the chosen probability value (0.05 in this case) and n is the number of arrays not already tested; testing continues until the first nonsignificant array is detected. The significance of departure from 1:1 of the pooled segregation ratios for each enzyme was tested using the G-test with Williams's adjustment (Sokal and Rohlf, 1995).

Formally, Gillet and Hattemer's methodology requires independent parental genotyping, that is, use of parental germplasm. When parental material is unavailable, as in the present case, putatively heterozygous mother-trees must first be identified on the basis of the presence of two homozygote types in the progeny. As, under dominant inheritance, the heterozygote is indistinguishable from one of the homozygotes, this means of identifying heterozygous parents in itself implicitly assumes codominant inheritance. Consequently, it is possible to test for departure from codominant inheritance only if a logical circularity is admitted. It follows that the strongest formal conclusion that can be made is that the segregation ratios of the observed phenotypes

are consistent with their proposed genetic interpretations. However, particularly when the phenotypes are consistent with those expected for the quaternary structure of the enzyme in question, the degree of confidence attached to the assertion of codominant inheritance does not appear to be critically less than when parental material is available, particularly as the use of adult material for identification of mother-trees heterozygous for alleles expressed in the progeny in itself requires an assumption not necessary in the present case, *i.e.* of that ontogenetic stability.

Linkage disequilibrium

As neither haploid material nor progeny-test data were available, linkage disequilibrium coefficients based on frequencies of coupling and repulsion heterozygotes were not estimable. Therefore, values of Burrows's composite linkage disequilibrium coefficient Δ_i and the corresponding correlation coefficient (Weir, 1979) were estimated for each of the seven populations. Significant values of Δ_i imply that there are differential frequencies of coupling and repulsion heterozygotes and/or non-random union of gametes (Roberds and Brotschol, 1985). Ohta's (1982) multiple population linkage disequilibrium (D) coefficients were also estimated. Ohta's partitioning of the variance of linkage disequilibrium is useful in elucidating the causes of observed disequilibria: when D_{IS}^2 (the variance component of disequilibrium within a population) is less than D_{ST}^2 (the variance of correlation between different gametes of one subpopulation relative to that of the total population) and D'_{IS}^2 (variance of within gamete correlation in a subpopulation relative to the total population) is greater than D'_{ST}^2 (variance of disequilibrium of the total population), then population subdivision (*i.e.* genetic drift) rather than epistatic selection is likely to be the main cause of linkage disequilibria (Kremer and Zanetto, 1997; Ohta, 1982). For estimation of both Burrows's and Ohta's coefficients, all alleles except the most common were pooled to a synthetic allele (Kremer and Zanetto, 1997). Missing values were eliminated from the data set as these add no information on non-gametic or gametic correlations in allele frequencies. Data from all collection years were pooled. POPGENE (Yeh and Boyle, 1997) was used for estimation of all the above disequilibrium parameters.

Neutrality

POPGENE was also used to test for selective neutrality. It uses Stewart's algorithm for the Ewens-Matterson neutrality test, as detailed in Manly (1985). One thousand iterations were employed for generation of simulated distributions of the F statistic (sum of squared allele frequencies) under the null hypothesis.

RESULTS

Inheritance

Aspartate aminotransferase

There was one clearly resolved locus, with three putative alleles. The banding patterns were those expected for a dimeric enzyme, although putative BB, BC, and CC genotypes frequently showed indistinct banding at the A locus (Figure 6-1). On some gels the heterodimer band of the putative AC genotype was slightly anodal of the expected intermediate position (Figure 6-2). None of the pooled segregation ratios of the progeny of putatively heterozygous (AB or AC) mother-trees was significantly different from the 1:1 expectation (Table 6-1). The lowest probability associated with individual progeny arrays, i.e. $p=0.07$ (family 4014) (Table 6-2) is greater than the corresponding Dunn-Šidák critical value of $1-(0.9)^{0.57} = 0.004$ ($n=27$).




Figure 6-1. Aspartate aminotransferase zymogram of *Pinus rubra*, showing (1) faint banding in A-locus (most cathodal) position in BC and BB genotypes (marked respectively 1 and 2); C indicates the control (*Pinus resinosa*).




Figure 6-2. Aspartate aminotransferase zymogram of *Pinus resinosa*, showing asymmetry of AC heterodimers in two families. 'C' indicates the control (*Pinus resinosa*).

Alcohol dehydrogenase

Three variable loci were detected. The most cathodal locus was too poorly and inconsistently resolved to permit interpretation. The most anodal locus (ADH β) showed the expected (May, 1998) dimeric banding pattern. Two putative alleles were detected (Figure 6-3). Segregation of the intermediate-mobility locus (ADH2) coincided completely with that of ADH β , but its banding pattern was characteristic of a monomeric enzyme (Figure 6-3). The pooled segregation ratios were not significantly different from expectations (Table 6-1). The lowest probability associated with individual progeny arrays, i.e. $p=0.04$ (family 1007) (Table 6-3) is greater than the corresponding Dunn-Šidák critical value of $1-(0.9)^{0.23}=0.002$ ($n=44$).




Figure 6-3. Alcohol dehydrogenase zymogram of *Pinus resinosa*. ADH2 (intermediate mobility) shows an apparently monomeric banding pattern and apparent close linkage with the dimeric, most anodal locus. 'C' indicates the control (*Pinus resinosa*).

Glucose-phosphate isomerase

The PGI zymograms were generally complex. However, the most simple of them reveal four zones of activity (Figure 6-4). The most anodal of these was too poorly resolved to permit scoring, whereas the complexity of the zymograms precluded genetic interpretation of the two most cathodal loci.




Figure 6-4. Example of less complex glucose-phosphate isomerase zymogram of *Pinus rubra*, showing four zones of activity (putative loci, labelled 1 to 4). 'C' indicates the control (*Pinus resinosa*).

PGL3 appeared to present banding patterns consistent with those expected for a dimeric enzyme with three alleles (Figures 6-5 and 6-6). However, the putative allele PGL3-B occurs within the apparent range of mobility of the putative locus PGL2. Therefore, PGL3-B and its possible combinations with allele PGL3-A could be confused with polymorphisms correctly attributable to locus PGL2. For this reason, only segregation ratios of apparent AC mothers were assessed (putative allele PGL3-C occurs anodal of the most common PGL3 allele, and there is therefore little risk of confusion with alleles of loci 1 and 2), (Figures 6-6 and 6-7). The pooled segregation ratio did not depart from the expectation under codominant inheritance (Table 6-1). None of the individual arrays departed significantly from expectations (Table 6-4).




Figure 6-5. Glucose-phosphate isomerase zymogram of *Pinus rubra*, showing putative genotypes of locus 3: 1=AA, 2=AB, 3=BB; C indicates control (*Pinus resinosa*).




Figure 6-6. Glucose-phosphate isomerase zymogram of *Pinus rubra*, showing putative genotypes of locus 3: 2=AC, 1=AA; C indicates control (*Pinus resinosa*); families 2026 (first lane), 2034 (final two lanes), 2027 (other lanes).




Figure 6-7. Glucose-phosphate isomerase zymogram of *Pinus rubra*, showing putative genotypes of locus 3: 1=AA; 2=AC; 3=CC. C indicates control (*Pinus resinosa*). Family QP12.

Phosphoglucomutase (PGM; EC 5.4.2.2)

PGM1 shows the monomeric banding pattern characteristic of this enzyme system (May, 1998), with three alleles (Figures 6-8, 6-9). None of the pooled PGM1 segregation ratios showed departures from expectations under codominance (Table 6-1). None of the individual progeny arrays showed nominally significant departures from the 1:1 expectation (Table 6-5). Allele PGM1-B showed the same relative migration as the least

common putative allele of the anodal PGM2 locus. Potentially ambiguous cases were scored based on band intensity and unambiguous occurrence of the putative alleles elsewhere in the progeny array in question. Resolution of PGM2 was inconsistent and did not permit genetic interpretation, although on the clearest gels (e.g. Figure 6-8) a polymorphic, monomeric banding pattern was evident.




Figure 6-8. Phosphoglucosylase zymogram of *Phloxieris rubra*, showing putative genotypes at locus PGM1: AA (marked 1), AB (marked 4).




Figure 6-9. Phosphoglucosylase zymogram of *Phloxieris rubra*, showing putative genotypes at locus PGM1: AA (marked 1), CC (marked 2), AC (marked 3).

Phosphoglucosylase dehydrogenase (PGD, EC 1.1.1.43)

Two single-banded phenotypes, i.e. putative homozygotes, and a three-banded putative heterozygote, consistent with expectations for this typically dimeric locus (May, 1998), were observed. However, genetic interpretation was complicated by the occurrence of two-banded phenotypes (Figures 6-10, 6-11). These consisted of one band of the same mobility as the heterodimer band, and one band of similar intensity at one of the two putative homodimer / homozygote positions. Initial segregation analysis without these phenotypes revealed a consistent homozygote excess in progeny of heterozygous

mother-trees. It was therefore hypothesized that the double-banded phenotypes represent additional AB heterozygotes. The pooled segregation ratio calculated on this basis showed a non-significant heterozygote excess (Table 6-1). The lowest probability associated with individual progeny arrays, i.e. $p=0.01$ (family Pachanga 7) (Table 6-6) is greater than the corresponding Dunn-Šidák critical value of $1-(0.9)^{0.192} = 0.002$ ($n=52$).




Figure 6-10. Phosphogluconate dehydrogenase zymogram of *Plumeria rubra*, showing putative genotypes: 1=AA, 2=BB, 3=AB, 4=double-banded phenotypes scored as AB (see text); 'C' indicates control (*Pinus resinosa*).




Figure 6-11. Phosphogluconate dehydrogenase zymogram of *Plumeria rubra*, showing putative genotypes: 1=AA, 2=BB, 3=AB, 4=double-banded phenotypes scored as AB (see text) (family 1004); 'C' indicates control (*Pinus resinosa*).

Linkage disequilibrium

There were significant ($p < 0.05$) linkage disequilibria in three of the 70 permutations of 7 populations and 10 locus-pairs (Table 6-7). Three of the populations each showed one disequilibrium, but not for the same pair of loci. Of the 10 possible pairwise locus combinations, three evinced linkage disequilibria. All estimates of correlations between loci were $< |0.13|$. For all pairwise combinations, D_{IS}^2 was less than D_{ST}^2 and D'_{IS}^2 was greater than D'_{ST}^2 (Table 6-8).

Selective neutrality

Observed values of AAT, PGI and PGM were within the 95 per cent confidence limits for selectively neutral loci. In the case of ADH and PGD, observed values of F were

generally close to, and in one case (ADH, Corobici) less than, the lower 95 per cent confidence limits (Table 6-9).

DISCUSSION

Inheritance

The segregation ratios of these loci are consistent with the hypothesis of codominant inheritance, suggesting that, in these populations of this species, factors that may lead to departure or apparent departure from codominance, such as meiotic drive, null allelic action, and pre- or immediate postzygotic selection (Strauss and Conkle, 1986; Xie *et al.*, 1991) are not important for these loci.

With the partial exception of *PGD1*, banding patterns accorded with expected quaternary structure. As the foregoing interpretation of the double-banded phenotypes in *PDG1* is itself based partly on observed segregation ratios, the non-significance of departure from expected segregation ratios in this locus represents less conclusive evidence of codominance than in the other loci. However, alternative explanations appear to be untenable. Interpretation of these phenotypes as double-banded homozygotes would imply a large excess of homozygotes within progeny of heterozygotes and a similar excess of homozygosity in the progeny as a whole, whilst, by contrast, the maternal genotypes would be inferred as largely heterozygous, due to the higher prevalence under this interpretation of arrays with both putative homozygotes. The present interpretation implies partial null action of these alleles when in the heterozygous condition. The similar staining intensity of the two bands suggests that, whilst a heterodimer of intermediate polarity is formed, the active component is contributed by only one of the alleles, i.e. the one inactive in the homodimer state is also inactive in the heterodimer. The occurrence of both double-banded and 'typical' heterozygotes within the same families suggests that this polymorphism may be artefactual, rather than caused by defective forms of the alleles themselves.

Linkage disequilibrium

With 70 permutations of population and locus-pair, chance (i.e. type two error) would be expected to result in three to four estimates with associated probabilities of 0.05. The

probability associated with one of the estimates (ADH/PGI, Palmira population) approaches this level; chance would appear to be the most parsimonious explanation for this observed disequilibrium.

The low probability values associated with the other disequilibria make the above explanation less likely, and suggest some linkage disequilibria in this tropical angiosperm, a result consistent with findings in temperate broadleaves (Granger, 1996; Huang *et al.*, 1996; Roberds and Brostchol, 1985; Zanetto *et al.*, 1996) and temperate and boreal conifer species (Cheliak and Pitel, 1985; Strauss and Conkle, 1986; Xie *et al.*, 1991; Yang and Yeh, 1993; Yeh *et al.*, 1994). There is insufficient information to distinguish between structural (i.e. location on the same chromosome) and non-structural causes of the observed linkage disequilibria. As haploid chromosome number of *P. rubra* appears to be $n=18$ (Bawa, 1973; van der Laan and Arends, 1985), and assuming that the loci on different chromosomes have equal chances of being selected for study, then the probability of any two of the five loci belonging to the same linkage group is $p = 1 - [(18)(17)(16)(15)(14) / 18^5] = 0.45$, indicating that, *a priori*, it is moderately likely that physical linkage is responsible for one of the observed disequilibria.

Linkage disequilibrium of either structural or non-structural origin can be caused either by epistatic selection or genetic drift. The values of Ohta's coefficients, together with the lack of consistency in the direction (sign) of the estimated significant and non-significant disequilibria (Table 6-8), suggest that selective forces are not responsible for the observed disequilibria. Non-directional forces such as founder effects, population subdivision and parental sampling effects (Yeh and Morgan, 1987) all represent possible causal factors. For many applications, e.g. mating systems studies, the causes of observed disequilibria are of less interest than their magnitude. Close linkage, structural or otherwise, leads to underestimates of outcrossing rates (Brown *et al.*, 1985; Yeh and Morgan, 1987). In the present case, correlations between loci are relatively weak (i.e. $< |0.13|$) and found only in three populations. As such, it is unlikely that they will appreciably affect estimates of mating system and other parameters.

Neutrality

The results of the Ewens-Watterson test suggest non-neutrality in both ADH and PGD. Observed F greater than or equal to the lower confidence limit suggests greater heterozygosity than expected under the infinite (neutral) alleles model, i.e. heterozygote superiority. However, the lack of departure from expected segregation ratios under codominance suggest that the selective forces responsible for maintaining the higher than expected levels of heterozygosity are absent (ADH) or relatively weak (PGD) at the sampled life-history stage (i.e. young progeny). It follows that the non-neutrality detected here is unlikely to influence parameter estimates, including mating parameters, based on young progeny material.

7. THE EFFECTS OF FOREST FRAGMENTATION ON GENETICS AND REPRODUCTION OF THE TREE *PLUMERIA RUBRA* L. IN NORTHWESTERN COSTA RICA

INTRODUCTION

Tropical forests were destroyed at a rate of around 15.2 million ha year⁻¹ in the decade 1990-2000 (FAO, 2001). However, deforestation is often not complete. Rather, in many cases, one or more tracts within the formerly continuous tree cover remain forested, and are converted by deforestation into fragments set in an unforested matrix. If biologically viable, such fragments may mitigate some negative consequences of deforestation. It follows that the biological implications, including genetic aspects, of the conversion of forest tracts to forest fragments are of considerable relevance to the management and conservation of forests and biodiversity.

The possible effects of forest fragmentation on genetic diversity are complex and interacting. The most immediate of these effects occurs at fragmentation, which, for species formerly present in deforested matrices, leads to reduction in population size. Depending on their size and allele frequencies, reduced populations may not contain all alleles formerly present, *i.e.* they may show founder effects (Meffe and Carroll, 1994; Yeh 2000). Continued low population size is expected to lead to further loss of variation due to random genetic drift (Hartl and Clark, 1989), a process which may be exacerbated by changed post-fragmentation environment. The latter has the potential to cause additional reductions in population size and higher variation in fertility (*e.g.* Aldrich and Hamrick, 1998; Kelly *et al.*, 2000), thereby reducing the ratio N_e/N (effective to census population sizes) (Nunney, 1993; Falconer, 1989). The effects of fragmentation on inter- and intra-population gene flow may exacerbate or mitigate such responses. For example, disturbance-mediated declines in density of tree populations may, due to decline in inter-tree pollinator movements (Karron *et al.*, 1995; Ghazoul *et al.* 1998), lead to increased geitonogamous selfing, which may also be caused by disturbance-mediated changes in pollinator assemblages (Aizen and Feinsinger, 1994). Increased selfing may have an immediate negative impact on fitness, *e.g.* by causing inbreeding depression (Gigord *et al.* 1998), and also

increases susceptibility to drift by further reducing N_e (Yeh 2000). Maintenance of prefragmentation levels of gene flow may, with time, restore variation lost in founder events and may also prevent cumulative drift. Although the proportion of immigrant seed and pollen in a given fragment x may be lower than when the fragment was a tract in continuous forest, at the same time it may originate to a higher degree than previously from fragments located further from x than those extinct pollen and seed sources once present in the matrix. When genetic distance correlates with physical distance, such migrants may be more effective, because of their greater genetic divergence (Mills and Allendorf, 1996). Furthermore, as suggested above, fragmentation and concomitant disturbance may reduce the number of pollination events involving 'home' pollen, implying higher migration rate for a given amount of incoming pollen.

Clearly, the effect of fragmentation on genetic diversity is not easy to predict, particularly given the relatively limited empirical information available on tree responses to fragmentation (Cornelius, 2003a). In the present document, the effects of forest fragmentation on the reproduction and population genetics of *Plumeria rubra* (calichuche, flor blanca, frangipani, sacuanjoche) are reported. *P. rubra* is a small to medium sized deciduous tree native from Mexico to Panama (Standley and Williams, 1969; Woodson, 1938), where it is found principally on sea-cliffs, limestone outcrops, and canyon sides within seasonally dry zones (Haber, 1984; Pittier, 1978). Pollination system is based on 'deceit'; the hawkmoth pollinator is attracted by the general conformity of the flowers to the 'hawkmoth syndrome', but there is no nectar reward (Haber, 1984). The latter author considers the absence of nectar in the case of *P. rubra* to be 'one of the great mysteries of pollination biology'. There is no published information on mating systems, although Haber (1984) remarks that the species's low ratio of fruit to flower production, given the close proximity of stigma and stamens, indicates self-incompatibility. The winged seeds appear to be selected primarily for wind dispersal, although dispersal in streamflow also seems inevitable where trees overhang watercourses. Individual trees can flower at two years age in favourable environments (Cornelius, personal observation). However, growth and development can be relatively slow in typical natural habitat. The largest trees in the present study zone are probably at least 30 years old.

In the study zone, *P. rubra* is found principally on river canyon-sides and rock outcrops. As a locally common species of naturally clumped distribution, information about the population genetics of *P. rubra* may generate insights of relevance to other taxa with similar life histories and similarly patchy or fragmented distributions. Additionally, as riparian forest appears to represent a large proportion of the remaining closed forest of this largely deforested zone, the 'genetic health' of *P. rubra* populations is linked to the conservation status of these habitats.

METHODS

Study zone and fragments

The study zone (Figure 7-1) is an area of approximately 100 km² located between 10°33' and 10°27' N, 85°05' and 85°10'W in the *canton* of Cañas, Guanacaste Province, northwestern Costa Rica. It is located in the northern sector of the wider study zone described in Chapter Three. Around 95% of the mean annual rainfall of 1693.4 mm ($s=459.4$) falls between May and November (San Luis, Cañas meteorological station, 1921-1978, MIRENEM, 1988). Altitude varies from 40-200m a.s.l.; mean annual temperature at 95m a.s.l. is 27.5°C (Jiménez *et al.*, 1987). The dry season is characterized by strong (up to 90km hr⁻¹) northerly winds (Coen, 1983) and temperatures up to 37°C. The deforestation of the study zone occurred mostly during the last 80 years, as a consequence of three main factors: the replacement by exotic pasture species of the semi-open woodland ('sitios') formerly used for grazing, the conversion of closed forest to grassland, and the conversion of woodland to agriculture (see Chapter Three). Currently, the dominant land use is beef cattle ranching, and the study zone consists in the main part of pastureland dissected by the predominantly northeast-southwest orientated tributaries of the Tenorio River. Forest in the study zone is concentrated along these watercourses (Chapter Three). Non-riparian forest is essentially confined to young secondary (*e.g.* abandoned pastureland) or degraded primary remnants.




Figure 7-1. Study zone and location of populations in a study of the effects of forest fragmentation on *Plumeria rubra* in northwestern Costa Rica

Within the zone, 7 populations were located using maps, aerial photographs and field exploration (Figure 7-1, Table 7-1). All populations except Pachanga and Sandillal are located in linear riparian forest. The Pachanga trees are drawn from two foci of *P. rubra* within an area of mixed pastureland, semi-abandoned pastureland, young secondary forest and canyon forest between the Magdalena and Corobicí Rivers and the hamlet of

Correntadas. In this area, the species occurs as widely-spaced small groups and individual trees. The Sandillal population is located on a ridge-edge overlooking the River Santa Rosa.

The study populations are not drawn from wholly discrete forest fragments set in a treeless matrix. Rather, they represent components of a degraded dendritic forest network. The degree of discontinuity and attrition of forest cover within this network varies greatly, *i.e.* from zero in stretches of intact primary forest, through sectors reduced to narrow bands of forest on both sides, to stream reaches bordered by isolated trees, to complete absence of forest on one or both river banks. Trees (although only very rarely *P. rubra*) are also relatively common in the pastureland matrix (Chapter Three). The distribution of *P. rubra* in the study zone is characterized by the occurrence of population foci or 'swarms' in sites where the species has a strong competitive advantage, *i.e.* canyon-sides and rocky-banked river reaches. In general, such forest has been less affected by deforestation, agricultural encroachment and grazing damage than more accessible sites on deeper and more productive soils. Consequently, these processes have increased the degree of mutual isolation of *P. rubra* populations within the study zone.

Field and laboratory procedures

Individual trees in each fragment were mapped and measured for dbh (except Praderas population). Neighbourhood density index (*NDI*) of each tree was calculated as:

$$NDI = t_{0,25} + t_{25,50} + t_{50,100}$$

where $t_{0,25}$ = number of conspecifics within 25m, (etc.). The index is a distance-weighted measure of the number of conspecifics within 100m of the tree in question, *i.e.* trees at 50-100m and 25-50m are given respectively one quarter and one half of the weighting of trees located within 25m. Population mean *NDI* and dbh were calculated (Table 7-1).

Seed collections were made from February to April of 1997 (all except Pachanga, Sandillal, Quebrada Palmira), 1998 (all except La Pachanga) and 1999 (all except San Isidro and Praderas). All trees with ripe fruit were collected in each case, except in those few cases where trees were inaccessible. Ripe capsules were collected directly from the trees using a pruning pole and attached basket. Seeds were extracted manually and stored until needed.

Starch gel electrophoresis was carried out on enzyme extracts prepared from recently emerged radicles. Five codominant isozyme loci (Cornelius 2003b) were assayed: aspartate aminotransferase (AAT, E.C. 2.6.1.1), alcohol dehydrogenase (ADH, E.C. 1.1.1.1), glucose-phosphate isomerase (PGI, E.C. 5.3.1.9), phosphoglucomutase (PGM, E.C. 5.4.2.2), phosphogluconate dehydrogenase (PGD, E.C. 1.1.1.43). Extraction and electrophoresis protocols are documented fully elsewhere (see Appendix One). These loci were selected from a wider group of 16 allozymes based on resolution, ease of interpretation and polymorphism (Chapter Six). ADH and PGD appear to be non-neutral, but segregation ratios indicate that selection is absent or weak between fertilization and the census age used (*i.e.* recent germinants) (Chapter Six). The sampling schedule is detailed in Table 7-2.

Specific research questions

A priori, there are several reasons to expect that *P. rubra* might tend towards higher subpopulation differentiation and lower within-population variation than many other tropical broadleaves. Firstly, its preferred discontinuous habitat results in a naturally-aggregated or fragmented population distribution. The aggregated distribution of *P. rubra* should also be contrasted with such distributions that result from temporary niches, *e.g.* due to catastrophic disturbance. Preferred *P. rubra* habitat, *e.g.* exposed rock on canyon sides, is not temporary in nature and appears to favour long persistence of small populations in the same place for long periods of time. In the absence of migration, such populations will be subject to loss of variation due to genetic drift. Secondly, abiotically-dispersed species are in general expected to show greater population differentiation than species whose diaspores are dispersed by highly mobile birds and mammals (Loveless, 1992). Although the occurrence of isolated single trees of *P. rubra* seems to demonstrate the possibility of long-distance dispersal of its samaras, these are nevertheless likely to be less mobile than seeds dispersed by animal vectors, which may actively seek neighbouring populations. Furthermore, although the dry season winds coincide with the seed dispersal period of *P. rubra*, the species habitat is often substantially sheltered from these winds, which would also tend to favour within-river, *i.e.* broadly north-south, movement. Thirdly, apparent population size tends to be relatively small in *P. rubra*, *i.e.* often in tens rather than hundreds. Genetic drift is more likely to be an important evolutionary factor in these circumstances. Fourthly, the population shape of *Plumeria rubra* is characteristically linear. Under both the

isolation-by-distance and stepping stone migration models, population differentiation is likely to be increased by such unidimensionality (Hartl and Clark, 1989; Wright, 1969; Kimura and Weiss, 1964).

Given these considerations, the present study addresses the following issues: firstly, does the species show any sign of reduced genetic variation and high spatial genetic structure? Secondly, how has *P. rubra* in the study zone responded to forest fragmentation and its concomitant effects?

Parental generation

Genetic parameters were estimated for maternal and progeny generations. As maternal material was unavailable, maternal genotypes were inferred using the most-likely-parent method (Brown and Allard, 1970), as programmed in Ritland's MLTR program (DOS version, Ritland, 1996). Initially, inferences were made on pooled (across collection years and capsules) arrays. However, as there is correlated mating within capsules of *P. rubra* (see results), pooling of capsular arrays within progenies could lead to information loss and consequent incorrect maternal genotype inferences. For example, the maternal parent of an array of 40 progeny consisting of equal numbers of XX and XY ($X \neq Y$) could (depending on allele frequencies) be inferred as XY, whereas if the array was known to be made up two capsular arrays, one exclusively XX and the other exclusively XY, the probability of maternal heterozygosity would be remote (*i.e.* 0.5^{20}) (in this case the genotypic composition of the array would most likely be attributable to fertilization by two homozygous genotypes XX and YY). Accordingly, inferred maternal genotypes were cross-checked against their corresponding capsular arrays and reassigned maternal genotype (*i.e.*, heterozygote to homozygote) where appropriate (see Appenndix Two for details). In the case of arrays which include two homozygous types, this procedure is unnecessary, as maternal genotype is known.

For the maternal generation and each of the progeny cohorts, allele frequencies and allelic richness (\mathcal{A}) (mean number of alleles per locus) were calculated. As all the sampled loci were polymorphic, \mathcal{A} in this case is equivalent to AP (mean number of alleles per polymorphic locus) (Berg and Hamrick, 1997). Nei's expected heterozygosity ($H_e = 1 - \sum_i p_i^2$, where

p =frequency of allele i) (Weir, 1996) was also estimated and averaged over all assayed loci. The fixation index F_{is} ($1-H_o/H_e$, where H_o =observed heterozygosity) was calculated as a measure of heterozygote deficit or excess with regard to HWE expectations. The null hypothesis of HWE within each population was tested using G-tests. All alleles but the most common were pooled to one synthetic class for both the significance testing (to maximize the number of observations per expected genotypic class) and calculation of F_{is} (for consistency between the measure of disequilibrium and its test statistics). However, due to the low population numbers, in many cases the number of observations was still ≤ 5 , usually considered as the threshold for uncritical application of the G-test (Sokal and Rohlf, 1995).

Homogeneity tests were carried out to test the null hypothesis of no population differentiation. As for the tests of Hardy-Weinberg equilibrium, and for the same reasons, the less common alleles were pooled to one synthetic allele. Wright's F-statistics (F_{is} , correlation between uniting gametes relative to the subpopulations; F_{it} , correlation between uniting gametes relative to the population as a whole; F_{st} , correlation between two gametes drawn randomly within subpopulations relative to that of two gametes drawn randomly from the population as a whole) were estimated as:

$$F_{is} = \frac{H_s - H_I}{H_s},$$

$$F_{it} = \frac{H_T - H_I}{H_T},$$

$$F_{st} = \frac{H_T - H_s}{H_T},$$

where H_I = observed average heterozygosity of individuals within subpopulations; H_s =expected average heterozygosity; H_T = total expected heterozygosity (gene diversity) for population as a whole (pooled subpopulations) (Yeh, 2000).

Average historical gene-flow was estimated using the F_{st} method (*i.e.* $mN=0.25(1-F_{st})/F_{st}$), where N =effective population size and m = average migration rate) (Yeh, 2000). Pairwise estimates of Nei's adjusted genetic distance were also calculated and UPGMA dendrograms (Weir, 1996) based on these were estimated. Significance of the relationship between pairwise genetic distance and pairwise geographical distance between fragments was tested using the Mantel test (Sokal and Rohlf, 1995) as programmed in Mantel Version 2.0 (Liedloff 1999).

Except where indicated, the above analyses were carried out using the software package POPGENE (Yeh and Boyle, 1997).

Current gene flow

The Sandillal population, which was completely sampled (*i.e.* sample size = population size) is monomorphic at three loci (see results). In this case, gene flow is estimable as:

$$m = \frac{q_i}{\bar{q}}$$

where m =proportion of immigrant alleles, q_i =allele frequency in the progeny generation, \bar{q} = mean allele frequency in the source population (Hamrick and Nason, 2000). As the results demonstrate correlation between genetic and geographic distance, the nearest population (*i.e.* River Magdalena, see Figure 7-1) was assumed to be the pollen source. Allele frequencies in the Magdalena population were based on average frequencies over the one to three collection years. These estimates may be biased by pollen immigration to the source, but they were preferred to inferred maternal genotypes or ovule allele frequencies (estimable with MLTR (Ritland, 1996)) both because they allow both for differences in male fertility and for pollen donors not included in the sample of mother-trees.

Mating system

The multilocus outcrossing rate (t_m), average single-locus outcrossing rate (t_s) correlations of outcrossing rate (r_s) and outcrossed paternity (r_p) were estimated by fragment using Ritland's MLTR programme (DOS Version 1.1) (Ritland, 1996), which employs Ritland and Jain's (1981) mixed mating system model. Within MLTR, the likelihood equations were

solved using the EM method, and standard errors computed with 1000 bootstraps. Estimates of t less than 1-2S.E. were considered to depart significantly from full outcrossing (Liengsiri *et al.*, 1998); analogous criteria were used for r_p , r_i , *i.e.* estimates greater than 2 SE were considered to differ significantly from zero. MLTR was also used for estimation of individual outcrossing rates and pollen allele frequencies. For these estimations, progeny arrays were pooled across collection years in order to maximize sample size. The significance of between fragment differences in mean individual-tree outcrossing rate was tested with the Kruskal-Wallis test (Sokal and Rohlf, 1989) using SAS (Version 8.0 for Windows, SAS Institute, Cary, NC, USA) (SAS, 1999). As there were many 'ties' (due to the prevalence of 100% outcrossing), Monte Carlo simulation of exact tests were used for generation of probability values for the K-W test (SAS, 1999). For all MLTR analyses, all available parameters were simultaneously estimated.

Fruit production

Total number of capsules per sample tree was recorded during each seed collection (open capsules persist for some time, so capsule counts are unbiased by any differences in maturation date). A positive relationship between capsule production and neighbourhood density would indicate pollinator limitation, *i.e.* less visitation. This relationship was tested for using multiple linear regression of the natural log of individual capsule count (averaged over collection years for each tree) on both dbh and neighbourhood density index. Dbh was used because of the evident and practically axiomatic relationship between tree size and flower production. The null hypothesis of no between-fragment differences in log capsule production was tested using analysis of covariance (dbh as covariate). Analyses were carried out using SAS.

RESULTS

Genetic variation

Maternal generation

All alleles of all loci were present only in the Quebrada Palmira and Río Corobici populations (Table 7-3). Fixation of one or more loci occurred in maternal trees sampled in the San Isidro, Praderas (both for PGI), Pachanga (PGM) and Sandillal (AAT, ADH, PGM) populations. For all loci except ADH, the most common allele was the same in each

population. For ADH, $p_A > q_B$ in one population, whilst $q > p$ in four populations and $p = q$ in two populations. The frequency of ADH-A varied from $p=0$ (Sandillal) to $p=0.53$ (San Isidro). Gene diversity (H_e) varied from 0.12 to 0.38. There were two nominally significant Hardy-Weinberg disequilibria (heterozygote excess in Río Corobici (PGD), and Praderas (ADH)). Twenty-five of the F_{is} values were negative, three were positive, one equal to zero, and seven were undefined ($p=1$).

The homogeneity tests (Table 7-4) indicate that the populations are heterogenous for ADH and PGI. The overall F_{st} estimate of 0.07 (0.03 without the smallest populations) (Table 7-5) suggests low to moderate genetic differentiation (Yeh 2000). Pairwise genetic distances (Table 6) were significantly positively related to pairwise geographic distances (Mantel test, $G=1.60$, $Z=7.14$, $r=0.275$, one-tailed $p=0.036$). Populations from Corobici and its tributaries group together, as do the two River Tenorito populations (Figure 7-2).



Figure 7-2. UPGMA dendrogram based on Nei's unbiased genetic distance between maternal trees in seven population of *P. rubra* located in northwestern Costa Rica.

Progeny: 1997

All alleles of all loci were present in the Corobici, Magdalena and San Isidro populations (Table 7-7). PGI-B was absent from Praderas. For all loci except ADH, the same allele was the most common in each population. For ADH, ADH-B was commonest in Corobici and

Magdalena, whilst $p_A \approx q_B$ or $p_A = q_B$ in San Isidro and Praderas. Gene diversity varied from 0.27 (Magdalena) to 0.36 (San Isidro). There were two significant HW disequilibria, both associated with homozygote excess. Most F_{is} estimates were close to zero; five were positive, 15 were negative and one undefined (Table 7-7).

The homogeneity tests (Table 7-8) indicate that sampled populations differ at all loci. The overall F_{st} estimate of 0.07 (0.03 without the smallest populations) (Table 7-9) suggests low to moderate genetic differentiation (Yeh 2000). The relationship between pairwise genetic and geographic distances (Table 7-10) was weak and non-significant (Mantel test: $g=0.0531$, $Z=1.166$, $r=0.0286$). The River Magdalena population grouped separately from the other populations (Figure 7-3).




Figure 7-3. UPGMA dendrogram based on Nei's unbiased genetic distance between four population of *P. rubra* located in northwestern Costa Rica (1997 progeny)

Progeny: 1998

All alleles were present in all six sampled populations (Table 7-11), except AAT-C (absent in Sandillal) and PGM-C (Quebrada Palmira, Praderas, Sandillal). There was near loss of alleles ($p \leq 0.03$) in Quebrada Palmira (PGM-C), San Isidro (PGM-C), Praderas (PGI-B, PGM-B), Magdalena (AAT-C, PGM-B,C) and Sandillal (AAT-B, ADH-A). For all loci except ADH, the same allele was the most common in each population. For ADH, ADH-B was commonest in Quebrada Palmira, Corobicí, Praderas, Magdalena, Sandillal whilst $p_A = q_B$ in San Isidro. Gene diversity varied from 0.18 (Sandillal) to 0.39 (San Isidro). There were seven significant HW disequilibria, four with homozygote excess and three with heterozygote excess. Eighteen F_{is} estimates were negative and 12 were positive.

Results of the homogeneity tests (not shown) were similar to those for the 1997 data, *i.e.* indicate highly significant heterogeneity for all loci (all probabilities <0.001). The F_{st} estimates of 0.069 for these six populations (Table 7-12) suggest low to moderate genetic differentiation. The relationship between pairwise genetic and geographic distances (Table 7-13) was strongly positive (Mantel test: $G=1.858$, $Z=6.82$, $r=0.39$) and significant (one-tailed $p=0.02$). The Magdalena and Sandillal populations grouped separately from the remaining populations (Figure 7-4).




Figure 7-4. UPGMA dendrogram based on Nei's unbiased genetic distance between four population of *P. rubra* located in northwestern Costa Rica (1998 progeny)

Progeny: 1999

All alleles were present in all five sampled populations (Table 7-14), except AAT-C (Sandillal), PGM-C (Magdalena, Pachanga, Sandillal). There was near loss of alleles ($p \leq 0.03$) in Quebrada Palmira (PGM-B, C), Corobicí (PGI-B), Magdalena (AAT-C, PGI-B), Pachanga (PGI-B, PGM-B) and Sandillal (AAT-B, PGM-B). For all loci except ADH, the same allele was the most common in each population. ADH-B was more frequent in Magdalena, Pachanga, and Sandillal whilst $p_A \approx q_B$ in Quebrada Palmira and Corobicí. Gene diversity varied from 0.14 (Sandillal) to 0.35 (Quebrada Palmira, Corobicí). There were four significant HW disequilibria, two with homozygote excess and two with heterozygote excess. Seventeen F_{st} estimates were negative, eight were positive, and one was undefined (monomorphism).

The homogeneity tests (not shown) indicated highly significant heterogeneity for all loci (all probabilities <0.001). The F_{st} estimates of 0.09 for these five populations (Table 7-15) suggest moderate genetic differentiation. The relationship between pairwise genetic and geographic distances (Table 7-16) was positive ($g=1.18$, $Z=4.65$, $r=0.422$) and weakly significant (one-tailed $p=0.08$). The Magdalena and Sandillal populations grouped separately from the remaining populations (Figure 7-5).




Figure 7-5. UPGMA dendrogram based on Nei's unbiased genetic distance between four population of *P. rubra* located in northwestern Costa Rica (1999 progeny)

Gene flow

For maternal and progeny data, mean estimates of gene flow based on the F_{st} method range from 2-3 individuals generation⁻¹ when the Sandillal and Pachanga populations are included, and around 4-7 individuals generation⁻¹ when these are omitted (Tables 7-5, 7-9, 7-12, 7-15).

Mean migration rates estimates into the Sandillal population from the Magdalena population were 0.14 and 0.12 in 1998 and 1999, respectively (Table 7-17).

Mating system parameters


In general, estimates of population outcrossing rates were >0.9 (Table 7-18). There was one statistically significant departure (*i.e.* by the $(\hat{i} + 2 \text{ s.e.} < 1.0)$ criterion (Liengsiri *et al.*, 1998) from 100% outcrossing (Sandillal 1998, $\hat{i} = 0.986 \pm 0.006$). Estimates for Praderas (1998 and pooled data) and Magdalena (1997, 1998, and pooled data) departed more strongly from full outcrossing, but with higher standard errors.

Estimates of population outcrossing rate based on mean single locus values (t_s) were lower than t_m estimates in all cases except three, and in general departed significantly from unity. Estimates of biparental inbreeding based on the difference between these two estimates of the outcrossing rate varied from 1.3% to 6.0%. Estimates of r_s were in the range 0.1-0.25. Most estimates of r_p were close to 0.9, with a minimum of 0.593.

Estimates of individual outcrossing rates ranged from complete selfing to complete outcrossing, but for all populations except Magdalena and Quebrada Palmira, most estimates were >0.9 (Figure 7-6). There were no significant differences between fragments in individual-tree outcrossing rates (Kruskal-Wallis test, chi-square 7.12, $p_{df=6}=0.31$). The scatter plot of individual-tree outcrossing rate against individual neighbourhood density indices (Figure 7-7) did not suggest any clear relationship between the two variables.

Fruit production

The log of number of capsules tree⁻¹ was significantly positively related to both dbh and neighbourhood density index (Table 7-19; Figure 7-8) (dbh and density index were not related).






Figure 7-6. Frequency distributions of estimates of individual-tree outcrossing rates in seven populations of *Plumeria rubra* located in northwestern Costa Rica




Figure 7-6. Frequency distributions of estimates of individual-tree outcrossing rates in seven populations of *Plumeria rubra* located in northwestern Costa Rica (continued)




Figure 7-7. Scatter plot of relationship between neighbourhood density index and individual-tree outcrossing rate in trees of *Plumeria rubra* from 6 forest fragments in northwestern Costa Rica.




Figure 7-8. Scatter plots of log number of capsules tree⁻¹ against dbh and neighbourhood density index rate in trees of *Plumeria rubra* from 6 forest fragments in northwestern Costa Rica.

Fragment mean numbers of capsules tree⁻¹ varied from 23.0 (Magdalena) to 98.0 (Sandillal) (Table 7-20). After taking into account the effect of diameter variation, there was no significant effect of fragment on mean log numbers of capsules tree⁻¹ (Table 7-21), whereas the effect of the covariate was highly significant.

DISCUSSION

Population differentiation

The results indicate that, for these loci, and for the maternal generation, there is low to moderate genetic differentiation between these populations of *P. rubra*. The estimates, although based on relatively few loci, are similar to average G_{st} values for comparable species, e.g. long-lived woody species in general (0.084 ± 0.008 , Hamrick *et al.*, 1992), and, within this group, outcrossing, animal-pollinated species (0.099 ± 0.017), wind-dispersed species (0.076 ± 0.009), although it is worth noting that Loveless (1992) reports a higher mean value for abiotically dispersed tropical species (0.138 ± 0.026). When the small Sandillal population is omitted from the analysis, the estimate of population differentiation is notably lower ($F_{st}=0.035$ vs. $F_{st}=0.075$). The estimates of F_{st} in the 1997 and 1998 progeny cohorts are closely similar to those of the maternal generation. The slightly higher estimates for the 1999 cohort may reflect lower sample sizes.

The results indicate a positive relationship between genetic distance and geographic distance. In the maternal generation, populations from Corobicí and its tributaries group together, as do the two River Tenorito populations. The small, discrete Sandillal population forms a separate group. In the progeny cohorts, the Magdalena population groups separately from both the River Corobicí and Tenorito populations, and is found in the same branch as the geographically relatively close Sandillal population.

Within-population genetic variation

Comparison of levels of within-population gene diversity shown by *P. rubra* with those of published meta-analyses is not straightforward, as the latter (e.g. 0.13 ± 0.011 for native tropical woody taxa, Loveless, 1992) are inclusive of monomorphic loci. However, as Loveless reported percentage of loci polymorphic (P) for the same group of 39.0 per cent, this implies mean gene diversity of polymorphic loci for this group of $(0.13/0.39) = 0.33$, i.e. similar to the estimates ported in Tables 7-3, 7-7, 7-11 and 7-14. Similarly, the estimates for *P. rubra* are comparable to other estimates based on polymorphic allozyme loci of neotropical trees, e.g. for *Enterolobium cyclocarpum* ($H_e=0.365$, five loci, Rocha and Lobo, (1996)), *Pithecellobium elegans* ($H_e=0.31$, recalculated for 6 polymorphic loci, Hall *et al.* (1994c)), *Pentaclethra macroloba* ($H_e=0.21$, three loci of adult trees, Hall *et al.*, 1994b), *Carapa*

guianensis ($H_e=0.31$, 6 loci, Hall *et al.* 1994a). There is, at least, no reason to conclude from the present data that *P. rubra* shows any general tendency to low levels of genetic diversity.

F_{is} estimates for the maternal generation were generally negative. Heterozygote excess was significant in two cases, in both cases at the two loci found to be non-neutral (Chapter Six). However, the greater prevalence and magnitude of negative F_{is} estimates for the maternal than the progeny generations suggests that heterozygote superiority in later age classes may not be restricted to these two loci. If this is the case, then selection, as well as migration, may contribute to maintaining high genetic variation and low genetic structure in this species.

Although, in general, these *P. rubra* populations have maintained high genetic variation and little divergence, there is, nevertheless, clear evidence of small population effects. The most notable case is that of the Sandillal population, which has retained only about 36% of the gene diversity (H_e) of surrounding populations. In part, this may reflect founder effects at fragmentation. This hypothesis is supported to some degree by the allelic richness of the Sandillal population ($A=1.4$). The expected average number of alleles remaining after a founder event is:

$$E = m - \sum_j (1 - p_j)^{2N},$$

where m = the number of alleles before population reduction, p_j = frequency of the j th allele, N = population size after reduction (Meffe and Carroll, 1994). Assuming pre-bottleneck frequencies equivalent to average frequencies for the Magdalena population (*i.e.*, $p=0.85$, $q=0.13$, $r=0.02$ (AAT); 0.23, 0.77 (ADH); 0.95, 0.05 (PGI), 0.965, 0.03, 0.005 (PGM); 0.73, 0.27 (PGD)), by this formula expected mean allelic richness for a 'sample' of four trees is $\hat{A} = 1.64$, *i.e.* not very different from the observed value of 1.40.

However, in general founder effects on gene diversity are less intense (Frankel and Soulé, 1981). Expected change in gene diversity as a result of a population size reduction (Young *et al.* (1996)) is:

$$\Delta H_e = \frac{-H_e}{S},$$

where S = number of gametes. In the present case ($H_e \approx 0.32$, $S=8$), the expected loss of 0.04 is much less than the observed loss of around 0.2, implying that additional factors have contributed to the loss of variation. One possibility is subsequent genetic drift. As expected gene diversity after t generations is approximated by:

$$H_e = H_0 e^{-\left(\frac{t}{2N}\right)},$$

where N = effective population size and H_0 = initial gene diversity, the number of generations required for a given change ΔH_e can be estimated using:

$$-t = (2N) \ln \frac{H_e}{H_0} \quad (\text{Hart and Clark, 1989}).$$

Assuming constant effective population size of $N=N_e=4$, the proportion of gene diversity lost in excess of that predicted due to founder effects (*i.e.* 0.12/0.28) would require $(\ln 0.43) / (-0.43) \approx 0.8$ or 7 generations. Although, given the short generation time of *P. rubra*, this number of generations does not seem too high to explain our observations, there may also be other factors at work, *e.g.* $N_e < N$ due to greater than expected variation in fecundity, overlapping generations, biparental inbreeding and possibly clumped population structure due to restricted seed dispersal.

Loss of genetic variation is of particular interest in the case of the locus ADH, which appears to be subject to balancing selection (*i.e.* heterozygote advantage) in the adult or older juvenile phases (see Chapter Six). Such loci are of intrinsic interest, *i.e.* not solely as 'markers'. Fixation has occurred at ADH3 (and, therefore, the closely linked ADH2 (Chapter Six)). Depending on the nature and presence in this population of the environmental factor(s) responsible for the selection pressure, this may imply a loss in population fitness, which will be permanent unless restored by migration or, improbably, by mutation.

There is some additional evidence for genetic drift effects on genetic variation. In all progeny years, the Magdalena and Praderas populations show lower gene diversity values than the other populations (except Sandillal). Praderas is the most isolated of all the populations (Figure 7-1), whilst the Magdalena population is located in a semi-urban and highly-disturbed setting, and both have relatively low population sizes. Relatively low gene diversity for the non-neutral ADH was also observed in the case of the Magdalena population ($H_e=0.39$, compared to $H_e\approx 0.5$ for the larger populations).

Gene flow

Its significance notwithstanding, the low-to-moderate degree of subpopulation differentiation described here suggest that to date, gene flow between these populations of *P. rubra* has been substantial. Overall estimates of gene flow based on the Island Model suggest an average of 2-7 migrants generation⁻¹. Given the failures in assumptions of the Island Model in the present case, it is possible that this range underestimates true values.

The direct estimates of current gene flow provide partial support for this argument. If applicable to other populations, the observed migration rate of 0.13 into the Sandillal population would imply mN well in excess of numbers usually deemed necessary to prevent undesirable population divergence. However, in the case of the smaller populations, such as Magdalena ($N=20$), such levels of gene flow may nevertheless be inadequate to prevent population divergence. The difference in grouping of the Magdalena population in progeny and adult generations, as well as the skewed frequencies of ADH in this population, seems to support this conjecture. In the case of Sandillal, quite extensive gene flow has demonstrably been insufficient to prevent divergence. As 26 per cent pollen flow corresponds to 0.5 migrants per generation, this is in accordance with theoretical predictions of substantial divergence at $mN < 1$, even taking into account small population size (*i.e.* using Wright's (1969) full equation). The case of the Sandillal population is perhaps an extreme one, both in terms of isolation and small population size. However, slightly larger populations such as Magdalena are not uncommon. Given the expectation that $N_e < N$, such populations could be at risk from genetic drift effects.

Mating system parameters

The results support Haber's (1984) deduction that, like many tropical trees, *P. rubra* is self-incompatible. Only one population (Sandillal) showed significant departure from full outcrossing in any year, and the estimate in question is very close to 100 per cent, *i.e.* of questionable biological importance. However, significant biparental inbreeding, *i.e.* mating between relatives, was noted in at least one year in all populations except Praderas. A number of factors could promote biparental inbreeding in *P. rubra*, *e.g.* limited seed dispersal ability, clumped habitat, and high correlation of paternity, itself due to uniparental or close to uniparental paternity of individual capsules. However, there is little indication of accumulation of inbreeding in the F_{st} estimates, which tend to be strongly negative in adult populations. This may indicate that inbreeding depression acts to purge progeny of related individuals, *i.e.* of full sibs, as has been speculated for the colonizer *Centaurea solstitialis* (Sun and Ritland, 1998). Differences between fragments in biparental inbreeding tend to be small and non-significant (*i.e.* within the limits of the respective standard errors), and therefore preclude the direct establishment of relationships between degree of biparental inbreeding and fragment characteristics.

The estimates of correlation of paternity suggest strongly that individual capsules of *P. rubra* tend to be sired by one or few pollen parents. A similar finding was reported for the apocynaceous *Stemmadennia donnell-smithii* (James *et al.*, 1998), and is consistent with the observed low frequency of pollinator visits (Haber, 1984).

There were a number of cases of individual outcrossing estimates <100 per cent, including two apparent cases of complete selfing. Such apparent departures from full outcrossing in individuals of a self-incompatible (or any other) species occur when fertilizing gametes have haploid multilocus genotypes that could be produced by the tree under study. This could occur either as a result of self-fertilization or as a result of fertilization by another tree with the same multilocus genotype or, at least, with ≥ 1 allele in common at each locus. Normally, such events would be unusual, *i.e.* confined to individual zygotes within progeny arrays. However, if all seeds within capsules are full-sibs, then each capsular array is analogous to each element (zygote) of a non-correlated array, in the sense that each represents an observation of a single pollination event. However, they are distinct in the sense that there

is a higher probability of unambiguous designation of the pollination event as an outcross, as the detection of even one unambiguous outcross in a full-sib capsular array would permit designation of the entire array as outcrossed. Many, and perhaps all, of the cases of apparent departure from complete outcrossing reported here may be due to biparental inbreeding. For example, Tree QP2, one of the two individuals with $\hat{t} = 0$, has inferred multilocus genotype AAT-AA, ADH-BB, PGI-AA, PGM-AA, PGD-AB. Its 24 progeny, all derived from one capsule, are all homozygous for the first four loci. Clearly, in the Quebrada Palmira population, which has relatively high diversity at the first three loci (Table 7-3), this array must either be due to selfing or pollination by one, similar genotype. Tree QP 1-309 has the same multilocus genotype, whilst the only progeny collected from the nearby tree QP4 has genotype consistent with the same maternal genotype. Based on these considerations, and given the apparent presence of an incompatibility mechanism in *P. rubra*, it seems reasonable to suggest that biparental inbreeding, rather than breakdown of self-incompatibility, represents the most parsimonious explanation of the observed departures from 100 per cent outcrossing.

The individual outcrossing rates provide additional confirmation of abundant potential gene flow in *P. rubra*. The progeny arrays assayed derived from the most isolated mother-trees (Palmira cemetery tree (nearest neighbour 390m), River Corobicí Tree 1000 (nearest neighbour at 318m), and River Corobicí Tree 1011 (nearest neighbour 190m) had $\hat{t} = 1.0$, $\hat{t} = 1.0$ and $\hat{t} = 0.9$, respectively, in spite of their high degree of isolation.

Fruit production

The estimated regression coefficient for NDI (0.056) indicates that increases in 1 unit of NDI (equivalent, for example, to 1 additional tree within 25m, or 4 additional trees between 50m and 100m) is associated with a multiplicative change in the median number of capsules for the NDI in question of $e^{0.056}$, i.e. 5.8% (, with lower and upper 95% confidence limits of $e^{0.0002} \approx 0\%$ and $e^{0.101} = 10.6\%$ (Ramsey and Schafer, 1997). For dbh=30 cm and NDI=0.25, predicted number of capsules is 23.66, whilst at NDI 4.0, predicted number of capsules is 29.2. The estimated slope of the relationship between NDI and capsules is the same as for dbh, although with wider confidence limits.

As outlined above, high outcrossing rates even in highly isolated trees indicate that the hawkmoth pollinators of *P. rubra* reach even highly isolated trees. However, the relationship with NDI seems to suggest that the probability of them so doing is related to the degree of individual isolation. This has implications for the effects of fragmentation on genetics of the species. Isolated populations, insofar as these are located in fragments with low resident population of pollinating species, will suffer reduced population fitness due to lower fertility. The same will occur in the case of low density populations, e.g. as the result of fragmentation-mediated disturbance.

Implications

Hawkmoths are known to be strong fliers and effective pollinators. *P. rubra*, a 'naturally-fragmented', hawkmoth-pollinated tree species, shows little sign of exhibiting unusually low within-population variation or high inter-population differentiation at the scale examined here, i.e. that appropriate to the spatial scale of the forest fragmentation itself. The lack of strong population differentiation exhibited by the species is indicative of substantial between fragment gene flow. Given the relatively low probability of long-distance seed dispersal, it seems reasonable to conclude that the nature of its pollinator is in large measure responsible for the apparent robustness to small population effects of *Plumeria rubra*. The self-incompatibility of the species, especially given the presence of pollinator limitation, also seems to contribute to this robustness.

Fundamentally, species occupy fragmented habitat because they are able to compete, grow and persist in such habitat. It appears that, in the case of *P. rubra*, this ability is conferred partially by the pollination system. Such species characteristics could be considered as constituting a variety of 'preadaptation' to habitat fragmentation. However, the resilience of this system is not inexhaustible, and, in certain cases, thresholds may be crossed beyond which adverse consequences may ensue. Some such consequences are illustrated by the present study, e.g. loss of genetic variation and declines in population fitness, as evidenced both by fixation at non-neutral loci and by reduction in fertility of isolated trees.

P. rubra appears to have a rather finely balanced pollination system: some of its characteristics enable it to overcome apparent isolation, whilst others appear to potentially increase its vulnerability to fragmentation. The potent scent-production of its flowers

enables it to attract pollinators at distance, and its spectacular floral displays aids visual location even at night. Its nectarless flowers presumably promote inter-tree pollinator movement (a goal possibly secured in nectar-producing mass-flowering species by inter-tree variation in nectar production (Frankie and Haber, 1983)) and may discourage repetitive crossing patterns, as might occur in 'triplining' behaviour in linear habitat (Janzen, 1974). However, as Haber (1984) pointed out, the effectiveness of its deceitful, imitative pollination system may be partially dependent on the persistence in increasingly degraded habitat of other, nectar-producing species. Consequently, although, to date, the continued persistence of this species in its favoured habitat seems not be at risk due to deforestation, fragmentation and disturbance, there is no guarantee that this will continue to be the case.

8. GENERAL CONCLUSIONS: GENETIC IMPLICATIONS OF FOREST FRAGMENTATION FOR SPECIES OF LOWLAND GUANACASTE PROVINCE

INTRODUCTION

The foregoing chapters have documented variable but clear actual and incipient genetic impacts of forest fragmentation on both *Anacardium excelsum* and *Plumeria rubra*. As indicated in the introduction (Chapter 1) to the present document, one of the justifications for individual species studies of this type is the generation of insights into the possible impact of fragmentation on other species. Accordingly, in this closing chapter this higher level of inference is considered, with particular reference to the tree species of the study zone itself. I begin with a brief overview of forest trees of the study zone, focussing particularly on those characteristics of particular relevance to the outcome of fragmentation, i.e. pollination and seed vectors, breeding systems and demographic aspects.

SPECIES OF THE STUDY ZONE

Jiménez *et al.* (1987) listed tree species found within the La Pacífica *hacienda*. Although some species present in the study zone are not found at La Pacífica, the list is sufficiently complete for the purposes of the present chapter. The reproductive and demographic characteristics of these species are summarized in Appendix Three (Tables A3-1 to A3-3; methodological details and sources are detailed in the appendix text). This detailed information is summarized below.

Approximately 69.3 per cent of the listed species are hermaphroditic (Table 8-1). Of these, for those for which information exists, 74 per cent are self-incompatible, whilst the remaining 26 per cent show varying degrees of self-compatibility. When monoecious, androdioecious and gynodioecious species are included, 70.1 per cent of those for which compatibility data exist are self-incompatible. 18.4 per cent of the species are dioecious. As would be expected, the distribution of breeding systems is similar to that described by Bawa (1974) and Bawa and Opler (1975) for the COMELCO property, situated some 25km to the

west. Bawa and Opler reported 20 per cent dioecy, and 79 per cent of (hermaphroditic) species as self-incompatible, a marginally higher proportion than reported here.

Approximately 59 per cent of species have biotically dispersed diaspores (birds, bats, terrestrial mammals), whereas around 41 per cent are abiotically dispersed (mostly wind) (Table 8-2). These tendencies are consistent with the 53 per cent zoochory reported by Opler (1978, reproduced and cited in Bullock, 1995) for the Cañas area. Bats, hawkmoths and medium-to-large bees are the dominant pollinators for respectively 9.9, 11.1 and 26.3 per cent of the species, whereas small bees and other small insects account for 44 per cent of the total (Table 8-2).

The mean number of stems species⁻¹ hectare⁻¹ for riparian and 'upland' (i.e. dry, non-riparian) sites at La Pacífica, based on Glander and Nisbett's (1996) data, was 2.81 ha⁻¹ and 8.93 ha⁻¹, respectively (Table A3-3). The respective medians were 0.5 ha⁻¹ and 0.2ha⁻¹; the divergence between the two measures reflects the presence of a number of species with high densities, particularly in the upland forest, which was dominated by *Lonchocarpus minimiflorus*. The data, which are based on censi including all stems ≥1cm, are in general consistent with the expectation of low adult population densities for tropical dry forest trees (Hubbell, 1979). However, there are a number of species with relatively high population densities: *Guzuma ulmifolia*, *Lonchocarpus minimiflorus*, *Myrospermum frutescens* (both site types), *Anacardium excelsum*, *Swietenia macrophylla* (riparian only), *Albizia caribaea*, *Cordia colocolca*, *Cordia alliodora*, *Lysiloma divaricatum*, *Luehea candida*, *Trichilia americana*, *Casearia corymbosa*, *Machaerium biovulatum* and *Tabebuia ochraceae* (upland site) all have densities ≥10 stems ha⁻¹.

As reported in Chapter 3, in the study zone as a whole these species subsist in forest remnants of increasingly linear shape, interspersed with almost treeless agricultural land or relatively well (tree-)stocked pastureland. Morales and Kleinn (2001) report that the most common pastureland tree species in the Cañas area are (in order of frequency): *Guzuma ulmifolia*, *Enterolobium cyclocarpum*, *Pithecellobium saman*, *Tabebuia rosea*, *Cordia alliodora*, *Byrsonima crassifolia*, *Bombacopsis quinata*, *Acrocomia uinifera*, *Mangifera indica*, *Cedrela odorata*, *Tabebuia ochraceae*, *Dalbergia retusa*, *Spondias mombin*, *Chione costaricensis*, *Albizia caribaea*, *Cassia grandis* and *Acosmium panamense*.

Clearly, species of the study zone are characterized by a wide range of reproductive and other characteristics, each the product of many generations of evolutionary change, in which adaptive response to natural selection may be assumed to have played an important role. The impact of fragmentation on genetics of these species is likely to depend on the genetic implications of these characteristics, 'adapted' (Begon *et al.*, 1990) by past environments, in present-day, human-modified environments. Obviously, the implications of fragmentation for each species cannot be identified without detailed, individual studies. However, given the findings of the present study, the species characteristics outlined above, and *a priori* considerations, some general points may be made and some specific examples cited.

EFFECTS ON OUTCROSSING

The findings for *A. excelsum*, in conjunction with previous studies (Karron *et al.*, 1995; Ghazoul *et al.*, 1998; Murawski and Hamrick, 1992; Murawski *et al.* 1990; Prober and Brown 1994) confirm the potential of fragmentation-mediated reductions in tree density to lead to increased selfing rates. This would primarily affect self-compatible species, such as *Ardisia revoluta*, *A. excelsum*, *Ceiba pentandra*, *Curatella americana*, *Calycophyllum candidissimum*, *Malpighia glabra*, *Muntingia calabura*, *Prockia crucis*, *Sloanea terniflora*, although individual trees of generally otherwise self-incompatible trees may also be self-compatible (e.g. Bawa, 1974). As the effect is dependent on changes in pollinator behaviour, it seems likely to be strongest in the case of species with relatively weak-flying pollinators, i.e., from the above list, *A. revoluta*, *A. excelsum*, *C. candidissimum*, *P. crucis*, *S. terniflora*. (see Table A3-2). As outlined in connection with *A. excelsum*, density reduction may occur with habitat degradation, or because of increased linearization of habitat. In the case of species with lower population densities than *A. excelsum*, fragmentation may also lead to direct increases in nearest neighbour distance, i.e. through removal of matrix trees. Although reduced tree density cannot generally lead to increased selfing in self-incompatible species, it should be noted that it may nevertheless be associated with increased geitonogamous pollination (without fertilization). It follows that even in such species, decreased tree density may negatively affect tree fertility.

In extreme cases of fragmentation, self-compatibility offers clear advantages, as pointed out by Baker (1955) for colonization. Evidently, for $N=1$ and restricted migration, a self-

compatible species has a higher chance of persistence in a given fragment than a self-incompatible or dioecious species. Given the low population density of many species of the zone, the capacity of self-fertilization may be an important influence in determining future species composition of isolated fragments.

HIGH VARIANCE IN REPRODUCTIVE OUTPUT

High variance in reproductive output, as noted in the case of some *A. excelsum* populations, is likely to result from two causes, which may be termed negative and positive effects. The negative case, in which normal levels of fertility are depressed in the majority of individuals (as in many *A. excelsum* fragments in the present study) is most likely to occur in species susceptible to fragmentation-induced moisture limitations. Many mesophytic species in highly disturbed fragments may be subject to such effects, e.g. *Ardisia revoluta*, *Andira inermis*, *Inga vera*, *Licania arborea*, *Ocotea veraguensis*, *Pithecellobium longifolium*, *Terminalia oblonga*. The positive case is exemplified by Aldrich and Hamrick's (1998) discovery of reproductive dominance of pastureland trees in *Symphonia globulifera*. Several of the common pastureland species listed by Morales and Kleinn (2001) also occur in riparian and other forest fragments, e.g. *G. ulmifolia*, *E. cyclocarpum*, *Pithecellobium saman*, *Tabebuia rosea*, *T. ochraceae*, *B. quinata*. Given the large size of many of the pastureland trees, particularly *E. cyclocarpum* and *P. saman*, there is potential for 'secondary bottlenecks' similar to those described by Aldrich and Hamrick for *S. globulifera*. However, with the exception of *Byrsonima crassifolia*, itself rare in forest fragments in the study zone, none of the common pastureland species listed by Morales and Kleinn are self-compatible, thus reducing the risk of 'swamping' of remnants with selfed propagules, even supposing that diaspore dispersal vectors have the potential to cause such effects.

FOUNDER EFFECTS

Low population densities imply acute founder effects, particularly in smaller fragments. Hubbell (1979) found that distribution of dry forest tree species tended to be clumped rather than evenly or randomly distributed. In a sense, clumping might mitigate founder effects, in that, whilst reducing the chance of species representation in any given fragment, at the same time it implies that more than one individual may be found in surviving fragments. However, single or very few individuals nevertheless occur in many small

fragments. For example, the Toronja fragment (area approximately 1ha) contains one individual of *Spondias mombin* (Anacardiaceae) and two individuals of the endangered 'cannonball-tree', *Couropita nicaraguensis* (Lecythidaceae). Similarly, the Ojoche fragment contains one *Brosimum alicastrum* ('ojoche') (Moraceae) tree. Furthermore, clumps of individual species may frequently be the product of limited seed dispersal rather than, necessarily, clumped habitat, and may therefore be more highly related than would be expected by chance. For example, the Bosque Duquesa fragment contains a small group of trees of the wind-dispersed bombacaceous species *Pseudobombax septenatum*, which is rare in similar habitat throughout the study zone. Such relatedness within clumps will tend to increase founder effects relative to expectations based on population sizes alone.

GENE FLOW AND RANDOM GENETIC DRIFT

Given the combination of low population densities and clumping of conspecifics, the capacity for at least occasional long-distance dispersal of gametes is likely to be of adaptive value even in undisturbed tropical dry forest. However, fragmentation will, in many cases, lead to a more extreme degree of population aggregation than under natural conditions, and it is possible that under such changed conditions some species's capacity for long-term dispersal may fall short of that consistent with maintenance of adequate genetic diversity in fragmented landscapes. The respective findings for *A. excelsum* and *P. rubra* illustrate this point. Within the Corobicí group, *A. excelsum* shows notably higher subpopulation differentiation, with F_{st} values approximately three times those of *P. rubra* (disregarding the atypical Sandillal population), even though the area covered by the *P. rubra* populations is actually somewhat larger than that of the *A. excelsum* populations. *P. rubra* is a more naturally-fragmented and less abundant species than *A. excelsum*. *A priori*, it might be expected to exhibit lower levels of genetic diversity and higher levels of population subdivision. That this is not the case seems likely to be due at least in part to the greater mobility of *P. rubra*'s hawkmoth pollinator, which to some degree appears to preadapt the species to forest fragmentation: the same factor that permits its persistence in naturally fragmented populations equips it also to persist within a fragmented forest set in a deforested matrix.

In Table 8-3, the data presented in Table 8-2 is further reduced to two pollen vector classes: relatively long-distance vectors, i.e. bats (Bawa, 1990), hawkmoths (Haber and Frankie, 1989), medium to large bees (Janzen, 1971; Frankie et al., 1976), fig wasps (Nason and Hamrick, 1997) and wind, and relatively short distance (small bees, beetles, small diverse insects) (Bawa and Opler, 1975). As, in general, biotic diaspore vectors are expected to be associated with more frequent long distance dispersal, the resulting cross-classification yields four (gametic) dispersal groups, which may be broadly characterized as 'long-long' (LL; i.e. biotic diaspore vector, long-distance pollinator), LS, SL and SS. This two by two classification of seed and pollen dispersal vectors reveals significant non-independence between the two factors ($p=0.007$, Fisher's exact test), with higher than expected numbers in each of the two LS/SL categories, and lower than expected numbers in LL and SS. That is, the implication is that even in unfragmented conditions (i.e. those under which the mechanisms in question evolved), for persistence in most niches it has been advantageous to have at least one (and perhaps not more than one) dispersal mechanism more suited to long-distance dispersal, and only relatively rarely has it not been disadvantageous to have neither. It might be speculated that, in undisturbed forest, species with neither long distance pollinators or highly mobile diaspore vectors tend to form relatively large contiguous stands or may be habitat generalists, i.e. with less clumped natural distributions. For such species (e.g. the dioecious Anacardiaceae *Astronium graveolens*, which is wind-dispersed, apparently pollinated by small insects, and found in both riparian and dry conditions) fragmentation may imply a very much more aggregated population distribution, and the SS dispersal syndrome may no longer be capable of maintaining adequate levels of inter-population gene flow.

P. rubra and *A. excelsum* both fall in the intermediate, 'long-short' dispersal categories. However, even in this case, there are several strands of evidence that gene flow correlates with distance: firstly, the presence of correlation between genetic and geographic distance in both species; secondly, the correlation between outcrossing rate and tree density in *A. excelsum*; thirdly, correlation between fruit production and tree density in *P. rubra*; fourthly, the low genetic variation present in the Sandhill population of *P. rubra*. It seems reasonable to suggest that species without either highly mobile pollen or diaspore vectors will tend to be less able than either species to maintain gene flow in fragmented landscapes. On the

basis of gene flow considerations alone, the following species would all appear to be at greater risk of fragmentation effects than either of the two study species: *Alvaradoa amorphoides*, *Acosmium panamense*, *Astronium graveolens*, *Calcyophyllum candidissimum*, *Cedrela odorata*, *Cordia alliodora*, *Esenbeckia littoralis*, *Lonchocarpus minimiflorus*, *Maecbaerium bivulatum*, *Swietenia macrophylla*, *Thouinidium decandrum*. The presence of trees in intervening pastureland matrices would tend to reduce risk. Of the above species, three (*Acosmium panamense*, *Cordia alliodora*, *Cedrela odorata*) are common in pastureland in the Cañas area (Morales and Kleinn, 2001). However, at present, pastureland trees, which are generally remnants of forests or traditional grazing *sitios*, appear not to be being replaced. Although their final disappearance may take many decades, their eventual loss seems inevitable under current farming practice. It follows that their contribution to maintenance of gene flow may be transient.

The classification of dispersal categories in Table 8-3 also assumes that dispersal vectors are actually present. This may not always be the case. Janzen and Martin (1982) argued that seeds of many tree species of lowland Guanacaste (e.g. *Crescentia alata*, *Enterolobium cyclocarpum*, *Hymenaea courbaril*, *Pithecolobium saman*) were formerly dispersed by the extinct Central American Pleistocene megafauna. Although their argument has been strongly criticized by Howe (1985), the latter author nevertheless concedes its possible applicability to particular species (i.e. those with 'botanical anomalies' such as indehiscent, pulp-filled fruit). Particularly in the case of seed that is never or rarely eaten by birds, bats or monkeys, this implies truncation of present-day seed shadows, i.e. relative to pre-extinction shadows. Although Janzen and Martin (1982) point out the role of domesticated bovine and equine cattle in substituting for extinct species, cattle are not always present. Even in pastureland matrices, they may be fenced out of forest fragments, particularly riparian fragments. Furthermore, it is necessary to distinguish dispersal in general from long-distance dispersal. The Pleistocene megafauna, like the semi-feral cattle herds of colonial Costa Rica (see Chapter Three), could wander and graze over vast areas. By contrast, under modern stock management practices, cattle are regularly moved between relatively small enclosed pastures. The largest current pastures rarely exceed 20 ha (Francisco Mesén, personal communication). In a roughly square pasture of this size, seed dispersal would be limited to approximately 500m, although occasional longer distance dispersal could occur if seed ingested in one pasture is defecated after rotation of cattle to a different pasture.

Ultimately, the impact of fragmentation depends not only on biological characteristics but also on what might be done in mitigation. In this sense, the enhanced persistence of forest in riparian situations provides ground for hope and also a basis for action. Sánchez-Azofeifa *et al.* (2001) remark that 'the current conservation strategy in Costa Rica may only preserve a small fraction of nature's wealth. More attention must be given to ecosystem restoration and regeneration in highly degraded tropical environments'. Effective application of existing forest law would constitute a major contribution to meeting these objectives. The present partial effectiveness of forest law in protecting riparian 'buffer' strips might imply that the current legal framework is flawed. However, the multiple benefits of conservation and restoration of riparian forest give ground for hope that, given adequate publicity, review and improvement of current policy might be feasible.

Given the long-distance gene flow that has been observed in a number of species, the improved conservation and restoration of riparian fragments, coupled with more active husbandry of pastureland trees, represents a promising approach to both maintaining biodiversity outside protected areas and ensuring reproductive connectivity between them. At present, such measures are clearly not in place, and the impact of forest fragmentation will depend largely on the biological characteristics of species in the prevailing environmental conditions. Inaction is likely to lead to gradual erosion of species richness of remaining fragments, coupled with increased isolation and genetic erosion of remaining populations of both rare and more common species.

BIBLIOGRAPHY

- Acuña, V.H. and Molina, I. 1991. Historia económica y social de Costa Rica (1750-1950). Editorial Porvenir S.A..
- Ågren, J. 1996. Population size, pollinator limitation, and seed set in the self-incompatible herb *Lythrum salicaria*. *Ecology* 77(6): 1779-1790.
- Aizen, M.A.; Feinsinger, P. 1994. Habitat fragmentation, native insect pollinators. And feral bees in Argentine "chaco serrano". *Ecological Applications* 4(2): 378-392.
- Aizen, M.A.; Feinsinger, P. 1994b. Forest fragmentation, pollination, and plant reproduction in a chaco dry forest, Argentina. *Ecology* 75(2): 330-351.
- Aldrich, P.R.; Hamrick, J.L. 1998. Reproductive dominance of pasture trees in a fragmented tropical forest mosaic. *Science* 281: 103-105.
- Anderson, W.R. 1983. *Byrsonima crassifolia*. Pp. 202-204 in Janzen, D.H. (ed.). Costa Rican Natural History. Chicago, University of Chicago Press. 816p.
- Anonymous, 2002. Index to plant chromosome numbers. Online at <http://mobot.mobot.org/W3T/Search/ipcn.html>. Accessed August 2002.
- Apsit, V.J.; Hamrick, J.L.; Nason, J.D. 2001. Breeding population size of a fragmented population of a Costa Rican dry forest tree species. *Journal of Heredity* 92(5): 415-420.
- Area de Conservación Guanacaste. Undated. Species home pages. Online at http://www.acguanacaste.ac.cr/paginas_especie/pag_principal/principal.html. Accessed September 2002.
- Augsberger, C.K. 1986. Morphology and dispersal potential of wind-dispersed diaspores of tropical trees. *American Journal of Botany* 73(3): 353-363.
- Baker, H.G. 1955. Self-compatibility and establishment after 'long-distance' dispersal. *Evolution* 9: 347-348.
- Bawa, K.S. 1973. Chromosome numbers of tree species of a lowland tropical community. *Journal of the Arnold Arboretum* 54: 422-434.

- Bawa, K.S. 1974. Breeding systems of tree species of a lowland tropical community. *Evolution* 28: 85-92.
- Bawa, K.S. 1990. Plant-pollinator interactions in tropical rain forests. *Annual Review of Ecology and Systematics* 21: 399-422.
- Bawa, K.S.; Bullock, S.H.; Perry, D.R.; Coville, R.E.; Grayum, M.H. 1985. Reproductive biology of tropical lowland rain forest trees. II. Pollination systems. *American Journal of Botany* 72(3): 346-356.
- Bawa, K.S.; Bullock, S.H.; Perry, D.R.; Coville, R.E.; Grayum, M.H. 1985a. Reproductive biology of tropical lowland rain forest trees. II. Sexual systems and incompatibility mechanisms. *American Journal of Botany* 72(3): 346-356.
- Bawa, K.S.; Frankie, G.W. 1983. *Cochlospermum vitifolium*. Pp. 215-216 in Janzen, D.H. (ed.). Costa Rican Natural History. Chicago, University of Chicago Press. 816p.
- Bawa, K.S.; Opler, P.A. 1975. Dioecism in tropical forest trees. *Evolution* 29: 167-179.
- Bawa, K.S.; Webb, C.J. 1984. Flower, fruit and seed abortion in tropical forest trees: implications for the evolution of paternal and maternal reproductive patterns. *American Journal of Botany* 71(5): 736-751.
- Becker, H.F. 1943. Land utilization in Guanacaste province of Costa Rica. *Geographical Review* 33: 74-85.
- Begon, M.; Harper, J.L.; Townsend, C.R.; 1990. Ecology: individuals, populations and communities, 2nd edition. Boston, USA. Blackwell Scientific, 945p.
- Berg, E.E.; Hamrick, J.L. 1997. Quantification of genetic diversity of allozyme loci. *Canadian Journal of Forest Research* 27: 415-424.
- Berge, G.; Nordal, I.; Hestmark, G. 1998. The effect of breeding systems and pollination vectors on the genetic variation of small plant populations within an agricultural landscape. *Oikos* 81: 17-29.
- Billington, H.L. 1991. Effect of population size on genetic variation in a dioecious conifer. *Conservation Biology* 5: 115-119.

Brook, B.W.; Tonkyn, D.W.; O'Grady, J.J.; Frankham, R. 2002. Contribution of inbreeding to extinction risk in threatened species. *Conservation Ecology* 6(1): 16.[online] URL: <http://www.consecol.org/vol6/iss1/art16>.

Brown and Allard. 1970. Estimation of the mating system in open-pollinated maize populations using isozyme polymorphisms. *Genetics* 66: 133-145.

Brown, A.H.D.; Barrett, S.C.H.; Moran, G.F. 1985. Mating system estimation in forest trees: models, methods and meanings. Pp. 32-49 in Gregorius, H-L. (ed.), *Population genetics in forestry (Lecture notes in biomathematics 60)*, Berlin, Springer, 287pp.

Brzustowski, J. Undated. Rarefaction calculator. On-line at http://www.biology.ualberta.ca/old_site/jbrzusto//rarefact.php/Calculator. Accessed August 2002.

Bullock, S.H. 1985. Breeding systems in the flora of a tropical deciduous forest in Mexico. *Biotropica* 17(4): 287-301.

Bullock, S.H. 1995. Plant reproduction in neotropical dry forests. Pp277-303 in Bullock, S.H.; Mooney, H.A.; Medina, E. (eds.) *Seasonally dry tropical forests*. Cambridge, England, Cambridge University Press, 450pp.

Burger, W.; Huft, M. 1995. *Flora costaricensis*. Fieldiana Botany New Series No. 36.

Buza, L.; Young, A.; Thrall, P. 2000. Genetic erosion, inbreeding and reduced fitness in fragmented populations of the endangered tetraploid pea *Swainsona recta*. *Biological Conservation* 93: 177-186.

Byers, D.L.; Waller, D.M. 1999. Do plant populations purge their genetic load? Effects of population size and mating history on inbreeding depression. *Annual Review of Ecology and Systematics* 30: 479-513.

Caballero, A. 1994. Developments in the prediction of effective population size. *Heredity* 73: 657-679.

Camacho, M.; Orozco, L. 1998. Patrones fenológicos de doce especies arbóreas del bosque montano del la Cordillera de Talamanca, Costa Rica. Online at www.ots.duke.edu/tropibiojnl/claris/46-3/CAMACHO.

- Campos, J.J.; Camacho, M.; Villalobos, R.; Rodríguez, C.M.; Gómez, M. 2001. Tala ilegal en Costa Rica: un análisis para la discusión. Online at <http://www.catie.ac.cr/noticias>.
- CBD (Convention on Biological Diversity). 1992. Convention on Biological Diversity, 5 June 1992. Online at <http://www.biodiv.org/doc/legal/cbd-en.pdf> (accessed June 2002).
- Centre for Tropical Forest Science. Undated. Tree Atlas of the Panama Canal Watershed. Online at <http://www.ctfs.si.edu/webatlas/maintreeatlas.html>
- Charlesworth, D. and Charlesworth, B. 1987. Inbreeding depression and its evolutionary consequences. *Annual Review of Ecology and Systematics* 18:237-268.
- Chase, M.R.; Boshier, D.H.; Bawa, K.S. 1995. Population genetics of *Cordia alliodora* (Boraginaceae), a neotropical tree. 1. Genetic variation in natural populations. *American Journal of Botany* 82(4): 468-475.
- Cheliak, W.M. and Pitel, J.A. 1985. Inheritance and linkage of allozymes in *Larix laricina*. *Silvae Genetica* 34 (4-5): 142-147.
- CITES, 1997. Proposal to the Convention on international trade in endangered species to include *Swietenia macrophylla* on Appendix II. Government of the Netherlands. Unpublished document.
- Coates, D.J. 1988. Genetic diversity and population genetic structure in the rare Chattering grass wattle, *Acacia anomala* Court. *Australian Journal of Botany* 36: 273-286.
- Coen, E. 1983. Climate. Pp.35-46 in Janzen, D.H. (Ed.). Costa Rican Natural History. University of Chicago Press, 816pp.
- Colegial, C. 1989. Hacienda Tenorio. Pp. 205-209 in Stephens, C.S. (ed.), Bananeros in Central America, True stories of the Tropics. Alva, Florida, U.S.A.. Banana Books.
- Costa Rica Legislative Assembly, 1996. Ley No 7575, Ley Forestal. Alcance No 21 a la Gaceta No. 72, 16 April 1996, 8pp.
- Crnokrak, P.; Roff, D. 1999. Inbreeding depression in the wild. *Heredity* 83: 260-270.
- Croat, T.B. 1979. The sexuality of the Barro Colorado Island flora (Panama). *Phytologia* 42(4): 319-348.

- Crow, J.F. and Aoki, K. 1984. Group selection for a polygenic behavioural trait: Estimating the degree of population subdivision. *Proceedings of the National Academy of Sciences, USA* 81: 6073-6077.
- Cunningham, S.A. 2000. Effects of habitat fragmentation on the reproductive ecology of four plant species in Mallee Woodland. *Conservation Biology* 14: 758-768.
- Dayanandan, S.; Dole, J.; Bawa, K.; Kesseli, R. 1999. Population structure delineated with microsatellite markers in fragmented populations of a tropical tree, *Carapa guianensis* (Meliaceae). *Molecular Ecology* 8: 1585-1592.
- Demauro, M.M. 1993. Relationship of breeding system to rarity in the Lakeside Daisy (*Hymenoxys acaulis* var. *glabra*). *Conservation Biology* 7: 542-550.
- Dick, C.W. 2001. Genetic rescue of remnant tropical trees by an alien pollinator. *Proceedings of the Royal Society, London, Series B, Biological Sciences*, 268: 2391-2396.
- Didham, R.K.; Ghazoul, J.; Stork, N.; Davis, A. 1996. Insects in fragmented forests: a functional approach. *Trends in Ecology and Evolution* 11(6): 255-259
- Edelman, M. 1992. The logic of the latifundio. Stanford, California. Stanford University Press, 478pp.
- Elizondo, L.H. 1999. *Cebus capucinus* Linnaeus. Online at <http://www.inbio.ac.cr/bims/ubi/mamiferos/ubiespecjo/ubiid=&find.html>. Accessed September 2002.
- Enquist, B.J.; Sullivan, J.J. 2001. Vegetative key and descriptions of tree species of the tropical dry forests of upland sector Santa Rosa, Area de Conservación Guanacaste, Costa Rica. Online at http://www.acguanacaste.ac.cr/paginas_especie/plantae_online/EnquistSullivanTreeKey.pdf.
- Falconer, D.S. 1989. Introduction to quantitative genetics, third edition. Harlow, Essex, Longman Scientific and Technical, 438pp.
- FAO (Food and Agriculture Organization of the United Nations) 2001. State of the World's Forests. Rome, Italy: FAO, 181pp.
- Fernández Bonilla, L. 1884. Historia de Costa Rica durante la dominación española. Madrid.

- Finkeldey, R.; de Guzman, N.; Changtragoon, S. 1998. Inheritance of isozyme phenotypes in *Pterocarpus indicus* Willd. *Forest Genetics* 5(3): 165-170.
- Fischer, M. ; Matthies, D. 1998. Effects of population size on performance in the rare plant *Gentianella germanica*. *Journal of Ecology* 86: 195-204.
- FONTAGRO (Fondo Regional de Tecnología Agropecuaria). 1997. Plan de Mediano Plazo. Online at <http://www.fontagor.org/pdf/5.pdf>. Accessed 13 August 2002.
- Foré, S.A.; Hickey, R.J.; Vankat, J.L.; Guttman, S.I.; Scafer, R.L. 1992. Genetic structure after forest fragmentation: a landscape ecology perspective on *Acer saccharum*. *Canadian Journal of Botany* 70: 1659-1668.
- Foster, M.S. 1983. *Chirocipha linearis*. 563-564 in Janzen, D.H. (ed.). Costa Rican Natural History. Chicago, University of Chicago Press. 816p.
- Foster, M.S.; McDiarmid, R.W. 1983. *Stemmadenia donnell-smithii*. Pp. 328-329 in Janzen, D.H. (ed.). Costa Rican Natural History. Chicago, University of Chicago Press. 816p.
- Fournier, L.A. 1992. Desarrollo y perspectivas del movimiento conservacionista costarricense. San José, Costa Rica. Editorial de la Universidad de Costa Rica, 113pp.
- Fowler, D.P.; Morris, R.W. 1977. Genetic diversity in red pine: evidence for low genic heterozygosity. *Canadian Journal of Forest Research* 7: 343-347.
- Frankel, O.H.; Soule, M.E. 1981. Conservation and evolution. Cambridge, England. Cambridge University Press, 327pp.
- Frankham, R. 1995. Effective population size/adult population size in wildlife: a review. *Genetical Research* 66: 95-107.
- Frankham, R. 1998. Inbreeding and extinction: island species. *Conservation Biology* 12(3): 665-675.
- Frankham, R.; Ralls, K. 1995. Inbreeding leads to extinction. *Nature* 392: 441-442.
- Frankie, G.W. and Haber, W.A. 1983. Why bees move among mass-flowering neotropical trees. In C.E. Jones and R.J. Little (Eds.). Handbook of experimental pollination biology, pp. 360-372. Van Nostrand Reinhold, New York, New York, USA.

- Frankie, G.W.; Baker, H.G.; Opler, P. 1976. Foraging behavior of solitary bees: implications for outcrossing of a neotropical forest tree species. *Journal of Ecology* 64: 1049-1057.
- Frankie, G.W.; Baker, H.G.; Opler, P.A. 1974. Comparative phenological studies of trees in the tropical wet and dry forests in the lowlands of Costa Rica. *Journal of Ecology* 62: 881-919.
- Frankie, G.W.; Haber, W.A.; Opler, P.A.; Bawa, K.S. 1983. Characteristics and organization of the large bee pollination system in the Costa Rican dry forest. Pp. 411-447 in Jones, C.E. and Little, R.J. (eds.), Handbook of experimental pollination biology. New York, Scientific and Academic Editions.
- Freitas, B.M.; Paxton, R.J. 1996. The role of wind and insects in cashew (*Anacardium occidentale*) pollination in NE Brazil. *Journal of Agricultural Science* 126: 319-326.
- Futuyma, D.J. 1986. Evolutionary Biology. Sunderland, Massachusetts, Sinauer Associates, 600pp.
- Fyfe, J.L.; Bailey, N.T.J. 1951. Plant breeding studies in leguminous forage crops. 1. Natural cross breeding in winter beans. *Journal of Agricultural Science* 41: 371-378.
- Gascon, C.; Lovejoy, T.E. 1998. Ecological impacts of forest fragmentation in central Amazonia. *Zoology* 101: 273-280.
- Gentry, A.H. 1983. *Tabebuia ochraceae* ssp. *neochrysantha*. Pp. 335-337 in Janzen, D.H. (ed.). Costa Rican Natural History. Chicago, University of Chicago Press. 816p.
- Ghazoul, J.; Liston, K.A.; Boyle, T.J.B. 1998. Disturbance-induced density dependent seed set in *Shorea siamensis* (Dipterocarpaceae), a tropical tree. *Journal of Ecology* 86: 462-474.
- Ghazoul, J.; McLeish, M. 2001. Reproductive ecology of tropical forest trees in logged and fragmented habitats in Thailand and Costa Rica. *Plant Ecology* 153: 335-345.
- Gigord, L.; Lavigne, C.; Shykoff, J.A. 1998. Partial self-incompatibility and inbreeding depression in a native tree species of la Réunion (Indian Ocean). *Oecologia* 117: 342-352.
- Gigord, L.; Picot, F.; Shykoff, J.A. 1999. Effects of habitat fragmentation on *Dombeya acutangula* (Sterculiaceae), a native tree on La Réunion (Indian Ocean). *Biological Conservation* 88: 43-51.

- Gilbert, F.; Willmer, P.; Semida, F.; Ghazoul, J.; Zalut, S. 1996. Spatial variation in selection in a plant-pollinator system in the wadis of Sinai, Egypt. *Oecologia* 108: 479-487.
- Gillet, E.; Hattemer, H.H. 1989. Genetic analysis of isozyme phenotypes using single-tree progenies. *Heredity* 63: 135-141.
- Gilpin, M.E.; Soulé, M.E. 1986. Minimum viable populations: processes of population extinction. Pp. 19-34 in Soulé, M.E. (Ed.), *Conservation Biology: the science of scarcity and diversity*. Sunderland, Massachusetts, Sinauer Associates.
- Glander, K.E. 1979. Feeding associations between howling monkeys and basilisk lizards. *Biotropica* 11(3): 235-236.
- Glander, K.E.; Nisbett, R.A. 1996. Community structure and species density in tropical dry forest associations al (sic) Hacienda La Pacífica in Guanacaste province, Costa Rica. *Brenesia* 45-46: 113-142.
- Gottsberger, G. 1986. Some pollination strategies in neotropical savannas forests. *Plant systematics and evolution* 152: 29-45.
- Granger, A.R. 1996. Inheritance and linkage of isozymes in sweet cherry. *Theoretical and Applied Genetics* (1996) 93: 426-430.
- Groom, M. 1994. Quantifying the loss of species due to tropical deforestation. Box 5B (p. 121) in Meffe, G.K. and Carroll, C.R. (eds.). *Principles of Conservation Biology*. Sunderland, Massachusetts, USA: Sinauer Associates, Inc., 600pp.
- Groom, M.J. 1998. Allee effects limit population viability of an annual plant. *American Naturalist* 151: 487-496.
- Groombridge, B.; Jenkins, M.D. 2002. *World Atlas of Biodiversity*. University of California Press, 340pp.
- Gross, C.L.; Mackay, D. 1998. Honeybees reduce fitness in the pioneer shrub *Melastoma affine* (Melastomataceae). *Biological Conservation* 86: 169-178.
- Gudmundson, L. 1983. Hacendados, políticos y precaristas: la ganadería y el latifundismo guanacasteco, 1800-1950. San José, Costa Rica. Editorial Costa Rica, 255pp.

- Haber, W.A. 1984. Pollination by deceit in a mass-flowering tropical tree, *Plumeria rubra* L. (Apocynaceae). *Biotropica* 16: 269-275.
- Haber, W.A. and Frankie, G.W. 1989. A tropical hawkmoth community: Costa Rican dry forest Sphingidae. *Biotropica* 21(2): 155-172.
- Hall, P.; Chase, M.R.; Bawa, K.S. 1994b. Low genetic differentiation but high population differentiation in a common tropical forest tree species. *Conservation Biology* 8(2): 471-482.
- Hall, P.; Orrell, L.C.; Bawa, K.S. 1994a. Genetic diversity and mating system in a tropical tree, *Carapa guianensis* (Meliaceae). *American Journal of Botany* 81(9): 1104-1111.
- Hall, P.; Walker, S.; Bawa, K. 1996. Effect of forest fragmentation on genetic diversity and mating system in a tropical tree, *Pithecellobium elegans*. *Conservation Biology* 10(3): 757-768.
- Hall, P.; Walker, S.; Bawa, K.S. 1994. Effect of forest fragmentation on genetic diversity and mating system in a tropical tree, *Pithecellobium elegans*. *Conservation Biology* 10(3): 757-768
- Hamilton, M.B. 1999. Tropical tree gene flow and seed dispersal. *Nature* 401: 120.
- Hamrick, J.L. 1987. Gene flow and the distribution of genetic variation in plant populations. Pp. 53-67 in Urbanska, K.M: (ed.), *Differentiation patterns in higher plants*. Academic Press, London.
- Hamrick, J.L. and Nason, J.D. 2000. Gene flow in forest trees. In Young, A.; Boshier, D; Boyle, T. (eds.). *Forest conservation genetics*, pp. 81-90. CABI Publishing, Wallingford, England / CSIRO Publishing, Collingwood, Victoria.
- Hamrick, J.L.; Godt, M.J.W.; Sherman-Broyles, S.L. 1992. Factors influencing levels of genetic diversity in woody plant species. *New Forests* 6: 95-124.
- Hamrick, J.L.; Murawski, D.A.; Nason, J.D. 1993. The influence of seed dispersal mechanisms on the genetic structure of tropical tree populations. *Vegetatio* 107/108: 281-297.
- Hartl, D.L. 2000. *A primer of population genetics*. Sunderland, Massachusetts. Sinauer Associates, 221pp.
- Hartl, D.L. and Clark, A.G. 1989. *Principles of population genetics*. Second edition. Sunderland, MA, USA, Sinauer Associates, 682pp.

- Hartshorn, 1983. *Trema micrantha*. Pp. 337-338 in Janzen, D.H. (ed.). Costa Rican Natural History. Chicago, University of Chicago Press. 816p
- Hartshorn, G.S. and Gentry, A.H. 1983. *Anacardium excelsum* (Espavél (*sic*), espavé, Acajou). Pp 189-190 in D.H. Janzen (ed.), Costa Rican natural history. The University of Chicago Press.
- Hedrick, P.W. and Kalinowski, S.T. 2000. Inbreeding depression in conservation biology *Annual Review of Ecology and Systematics* 31: 139-62.
- Heithaus, E.R.; Fleming, T.H.; Opler, P.A. 1975. Foraging patterns and resource utilization in seven species of bats in a seasonal tropical forest. *Ecology* 56: 841-854.
- Hodgkiss, P. 2001. Allozyme, isozyme electrophoresis: assaying genetic variation. Online at [Http://dendrome.ucdavis.edu/~phodgski/iso.index.html](http://dendrome.ucdavis.edu/~phodgski/iso.index.html).
- Howe, H.F. 1985. Gomphothere fruits: a critique. *The American Naturalist* 156(6): 853-865.
- Huang, H.; Dane, F.; Norton, J.D.; Weaver, D.B.; Hebard, F.V. 1996. Linkage relationships of isozymes and morphological traits in interspecific chestnut crosses. *Hortscience* 31(3): 419-420.
- Hubbell, S.P. 1979. Tree dispersion, abundance and diversity in a tropical dry forest. *Science* 203: 1299-1309.
- Hurlbert, S.H. 1971. The nonconcept of species diversity: a critique and alternative parameters. *Ecology* 52(4): 577-586.
- Husband, B.C. and Schemske, D.W. 1996. Evolution of the magnitude and timing of inbreeding depression in plants. *Evolution* 50(1): 54-70.
- Ibarra, E. 1990. Las sociedades cacicales de Costa Rica (siglo XVI). Colección Historia de Costa Rica. San José, Costa Rica, Editorial de la Universidad de Costa Rica, 246pp.
- Ibarra-Manríquez, G.; Martínez Ramos, M.M.; Oyama, K. 2001. Seedling functional types in a lowland rain forest in Mexico. *American Journal of Botany* 88: 1801-1812. Appendix. Specimen vouchers and germination data. American Journal of Botany supplementary data online at <http://ajbsupp.botany.org/v88>.

IGCR (Instituto Geográfico de Costa Rica). 1955. Lajas. 1:25000 topographic map series, sheet 3146 I NE. San José, Costa Rica, Instituto Geográfico de Costa Rica.

IGCR (Instituto Geográfico de Costa Rica). 1956a. Bebedero. 1:25000 topographic map series, sheet 3147 II SW. San José, Costa Rica, Instituto Geográfico de Costa Rica.

IGCR (Instituto Geográfico de Costa Rica). 1956b. Pelado. 1:25000 topographic map series, sheet 3147 II SE. San José, Costa Rica, Instituto Geográfico de Costa Rica.

IGCR (Instituto Geográfico de Costa Rica). 1956c. Taboga. 1:25000 topographic map series, sheet 3146 I NW. San José, Costa Rica, Instituto Geográfico de Costa Rica.

IGCR (Instituto Geográfico de Costa Rica). 1956d. Corobicí. 1:25000 topographic map series, sheet 3147 II NW. San José, Costa Rica, Instituto Geográfico de Costa Rica.

IGN (Instituto Geográfico Nacional). 1972. Mapa histórico-geográfico de Costa Rica, 3rd edition. San José, Costa Rica, Instituto Geográfico Nacional.

IGN (Instituto Geográfico Nacional). 1973. Abangares. 1:50000 topographic map series, sheet 3146 I. San José, Costa Rica, Instituto Geográfico Nacional.

IGN (Instituto Geográfico Nacional). 1988a. Liberia. 1:200000 topographic map series, sheet CR-2CM-1. San José, Costa Rica, Instituto Geográfico Nacional.

IGN (Instituto Geográfico Nacional). 1988b. Nicoya. 1:200000 topographic map series, sheet CR-2CM-4. San José, Costa Rica, Instituto Geográfico Nacional.

INEC (Instituto Nacional de Estadística y Censo). 2002. Resultados Generales IX Censo Nacional de Población y V de Vivienda. Online at <http://www.inec.go.cr/INEC2/resultadoscenso2000.htm>.

James, T.; Vege, S.; Aldrich, P. 1998. Mating system of three tropical dry forest tree species. *Biotropica* 30(4): 587-594.

Janzen, D.H. 1967. Synchronization of sexual reproduction of trees within the dry season in Central America. *Evolution* 21: 620-637.

Janzen, D.H. 1971. Euglossine bees as long distance pollinators of tropical plants. *Science* 171: 203-205.

Janzen, D.H. 1974. The deflowering of Central America. *Natural History* 83: 48-53.

- Janzen, D.H. 1976. The microclimate difference between a deciduous forest and adjacent riparian forest in Guanacaste province, Costa Rica. *Brenesia* 8: 29-33.
- Janzen, D.H. 1983. *Gnazuma ulmifolia*. Pp. 246-248 in Janzen, D.H. (ed.). Costa Rican Natural History. Chicago, University of Chicago Press. 816p.
- Janzen, D.H. 1983a. *Sapranthus palanga*. Pp. 320-322 in Janzen, D.H. (ed.). Costa Rican Natural History. Chicago, University of Chicago Press. 816p.
- Janzen, D.H. 1983b. *Crescentia alata*. Pp. 222-224 in Janzen, D.H. (ed.). Costa Rican Natural History. Chicago, University of Chicago Press. 816p.
- Janzen, D.H. 1983c. *Hymenaea courbaril*. Pp. 253-257 in Janzen, D.H. (ed.). Costa Rican Natural History. Chicago, University of Chicago Press. 816p.
- Janzen, D.H. 1983d. *Casearia corymbosa*. Pp.208-210 in Janzen, D.H. (ed.). Costa Rican Natural History. Chicago, University of Chicago Press. 816p.
- Janzen, D.H. 1986. Guanacaste National Park: tropical ecological and cultural restoration. San José, Costa Rica, Editorial Universidad Estatal a Distancia (UNED), 103pp.
- Janzen, D.H. 1986. Tropical dry forests, the most endangered major tropical ecosystem. Pp. 130-144 in Wilson, E.O. (Ed.). Biodiversity. Washington, National Academy of Sciences, 521p.
- Janzen, D.H. 1988. Management of habitat fragments in a tropical dry forest: growth. *Annals of the Missouri Botanical Garden* 75: 105-116.
- Janzen, D.H.; Liesner, R. 1980. Annotated check-list of plants of lowland Guanacaste province, Costa Rica, exclusive of grasses and non-vascular cryptogams. *Brenesia* 18: 15-90.
- Janzen, D.H.; Martin, P.S. 1982. Neotropical anachronisms: the fruits the gomphotheres ate. *Science* 215: 19-27.
- Janzen, D.H.; Miller, G.A.; Hackforth-Jones. J.; Pond, C.M.; Hooper, K.; Janos, D.P. 1976. Two Costa Rican bat-generated seed shadows of *Andira inermis* (Leguminosae). *Ecology* 57: 1068-1075.
- Janzen, D.H.; Wilson, D.E. 1983. Mammals. Pp. 426-442 in Janzen, D.H. (ed.). Costa Rican Natural History.

- Jennersten, O. 1988. Pollination in *Dianthus deltooides* (Caryophyllaceae): effects of habitat fragmentation on visitation and seed set. *Conservation Biology* 2: 359-366.
- Jiménez, Q.; Elizondo, L.H.; Corrales, L.F.; Alfaro, R.M. 1987. Estudio Ecológico de Hacienda La Pacífica, Cañas, Guanacaste. Fundación Neotrópica, Fundación de Parques Nacionales, Programa de Patrimonio Natural, Informe de Consultoría Preparado para Asociación Centro Ecológico La Pacífica.
- Jules, E.S. 1998. Habitat fragmentation and demographic change for a common plant: *Trillium* in old-growth forest. *Ecology* 79: 1645-1656.
- Kahmen, S.; Poschlod, P. 2000. Population size, plant performance, and genetic variation in the rare plant *Arnica montana* L. in the Rhön, Germany. *Basic and Applied Ecology* 1: 43-51.
- Karron, J.D.; Thumser, N.N.; Tucker, R.; Hessenauer, A.J. 1995. The influence of population density on outcrossing rates in *Mimulus ringens*. *Heredity* 75: 175-180.
- Kellman, M. Tackaberry, R.; Meave, J. 1996. The consequences of long-term fragmentation.. Pp 37-58 in Schelhas, J.; Greenberg, R., Forest patches in tropical landscapes. Washington D.C., Island press. 426pp.
- Kellman, M., Tackaberry, R., Brokaw, N.; Meave, J., 1994, Tropical Gallery Forests. National Geographic Research & Exploration, 10(1), 92-103.
- Kelly, D.; Ladley, J.J.; Robertson, A.W.; Norton, D.A. 2000. Limited forest fragmentation improves reproduction in the declining New Zealand mistletoe *Peraxilla tetrapetala* (Loranthaceae). Pp. 241-252 in Young, A.G. and Clarke, G.M. (Eds.), Genetics, demography and viability of fragmented populations. Cambridge, England, Cambridge University Press, 438pp.
- Kimura, M.; Weiss, G.H. 1964. The stepping stone model of population structure and the decrease of genetic correlation with distance. *Genetics* 49: 561-576.
- Koptur, S. 1983. *Inga*. Pp. 259-261 in Janzen, D.H. (ed.). Costa Rican Natural History. Chicago, University of Chicago Press. 816p
- Kremer, A.; and Zanetto, A. 1997. Geographical structure of gene diversity in *Quercus petraea* (Matt.) Liebl. II: Multilocus patterns of variation. *Heredity* 78: (1997): 476-489.

- Laan, F.M. van der; Arends, J.C. 1985. Cytotaxonomy of the Apocynaceae. *Genetica* 68: 3-35.
- Lacy, R.C. 1993. VORTEX: a computer simulation model for population viability analysis. *Wildlife Research* 20: 45-65.
- Lande, R. and Barrowclough, G.F. 1987. Effective population size, genetic variation, and their use in population management. Pp. 87-125 in Soulé, M.E. (Ed.), *Viable populations for conservation*. Cambridge University Press, 1987
- Laurance, W.F. 1999. Reflections on the tropical deforestation crisis. *Biological Conservation* 91: 109-117.
- Laurance, W.F.; Laurance, S.G.; Ferreira, L.V.; Rankin-de Meona, J.M.; Gascon, C.; Lovejoy, T.E. 1997. Biomass collapse in Amazonian forest fragments. *Science* 278: 117-118.
- León, J. 1952. Nueva geografía de Costa Rica. San José, Costa Rica. Librería la Española.
- Liedloff, A. 1999. Mantel nonparametric test calculator, Version 2.00. <http://www.sci.qut.edu.au/nrs/mantel.htm>.
- Liengsiri, C.; Boyle, T.J.B.; Yeh, F.C. 1998. Mating system in *Pterocarpus macrocarpus* Kurz in Thailand. *Journal of Heredity* 89: 216-221.
- Liengsiri, C; Piewluang, C.; Boyle, T.J.B. 1990. Starch gel electrophoresis of tropical trees. A manual. Saraburi, Thailand, ASEAN-Canada Forest Tree Seed Centre, 52pp.
- Linhart, Y.B.; Grant, M.C. 1996. Evolutionary significance of local genetic differentiation in plants. *Annual Review of Ecology Systematics* 27: 237-277.
- Little, E.L.; Wadsworth, F.H. 1964. Common trees of Puerto Rico and the Virgin Islands. USDA Agricultural Handbook 249.
- Little, E.L.; Woodbury, R.O.; Wadsworth, F.H. 1974. Trees of Puerto Rico and the Virgin Islands. Vol 2. USDA Agricultural Handbook 449, 1024pp.
- Loiselle, B.A.; Sork, V.L.; Nason, J.; Graham, C. 1995. Spatial genetic structure of a tropical understory shrub, *Psychotria officinalis* (Rubiaceae). *American Journal of Botany* 82: 1420-1425.
- Loveless, M.D. 1992. Isozyme variation in tropical trees: patterns of genetic organization. *New Forests* 6: 67-94

- Manly, B.F.J. 1985. The statistics of natural selection on animal populations. Chapman and Hall, London.
- Marshall, D.R.; Brown, A.H.D. 1975. Optimum sampling strategies in genetic conservation. Pp53-80 in Frankel, O.H.; Hawkes. J.G. (Eds.), Crop genetic resources for today and tomorrow. Cambridge, U.K., Cambridge University Press, 492pp.
- Másís, A.; Espinoza, R.; Guadamuz, A.; Pérez, D.; Chavarría, F. 1998. Species page of *Ocotea veraguensis*. Online at www.acguanacaste.ac.cr/paginas_especie/plantae_online/magnoliophyta/lauraceae/ocotea_veraguensis/prin_o.veraguensis.html. Accessed September 2002.
- May, B. 1998. Starch gel electrophoresis of allozymes. Pp.1-28 in Hoelzel, A.R. (Ed.). Molecular genetic analysis of populations. A practical approach. Oxford, England, IRL Press at Oxford University Press, 445 pp.
- Meffe, G.K.; Carroll, C.R. 1994. Genetics: conservation of diversity within species. Pp.143-178 in Meffe, G.K. and Carroll, C.R. (eds.). Principles of Conservation Biology. Sunderland, Massachusetts, USA: Sinauer Associates, Inc., 600pp.
- Meléndez, C. 1955. Bagaces, un pueblo en olvido. *Boletín del Museo Nacional* 2 (2): 11-12.
- Meléndez, C. 1974. Viajeros por Guanacaste. San José, Costa Rica. Ministerio de Juventud, Cultura y Deportes, Costa Rica, 557pp.
- Mesquita, R.C.G.; Delamônica, P.; Laurance, W.F. 1999. Effect of surrounding vegetation on edge-related tree mortality in Amazonian forest fragments. *Biological Conservation* 91: 129-134.
- Mills, L.S.; Allendorf, F.W. 1996. The one-migrant-per-generation rule in conservation and management. *Conservation Biology* 10(6): 1509-1518.
- MIRENEM (Ministerio de Recursos Naturales, Energía y Minas, Costa Rica). 1988. Catastro de las series de precipitaciones medidas en Costa Rica. San José, Costa Rica, 361 pp.
- Mitchell, J.D. and Mori, S.A. 1987. The cashew and its relatives (*Anacardium*. Anacardiaceae). *Memoirs of the New York Botanical Garden* 42: 1-76.
- Molina, M. 1996. Tratamientos pregermanitivos para especies tropicales del bosque seco tropical. *Rothschildia* 3(1). Online at

- http://www.acguanacaste.ac.cr/paginas_especie/pag_principal/principal.html. Accessed September 2002.
- Mooney, H.A.; Bullock, S.H.; Medina, E. 1995. Introduction. Pp.1-8 in Bullock, S.H.; Mooney, H.A.; Medina, E. (Eds.) Seasonally dry tropical forests. Cambridge, England, Cambridge University Press, 450pp.
- Morales, D.; Kleinn, C. 2001. Arboles fuera del bosque, conceptos, importancia e evaluación en Costa Rica. Online at <http://www.rlc.fao.org/proyecto/rla133ec/AFB-pdf/AFBpercent20CR.pdf>. Accessed September 2002.
- Murawski, D.A. and Hamrick, J.L. 1991. The effect of density of flowering individuals on the mating systems of nine tropical tree species. *Heredity* 67: 167-174.
- Murawski, D.A. and Hamrick, J.L. 1992. The mating system of *Cavanillesia platanifolia* under extremes of flowering-tree density: a test of predictions. *Biotropica* 24: 99-101.
- Murawski, D.A.; Gunatilleke, I.A.U.N.; Bawa, K.S. 1994. The effects of selective logging on inbreeding in *Shorea megistophylla* (Dipterocarpaceae) from Sri Lanka. *Conservation Biology* 8(4): 997-1002.
- Murawski, D.A.; Hamrick, J.L. 1991. The effect of the density of flowering individuals on the mating systems of nine tropical tree species. *Heredity* 67: 167-174.
- Murawski, D.A.; Hamrick, J.L.; Hubbell, S.P.; Foster, R.B. 1990. Mating systems of two bombacaceous trees of a neotropical moist forest. *Oecologia* 82: 501-506.
- Murcia, C. 1996. Forest fragmentation and the pollination of neotropical plants. Pp. 19-36 in Schelhas, J.; Greenberg, R., Forest patches in tropical landscapes. Washington D.C., Island press. 426pp.
- Murphy, P.G.; Lugo, A.E. 1995. Dry forests of Central America and the Caribbean. Pp. 9-34 in Bullock, S.H.; Mooney, H.A.; Medina, E. (Eds.) Seasonally dry tropical forests. Cambridge, England, Cambridge University Press, 450pp.
- Mustajärvi, K.; Siikamäri, P.; Rytönen, S.; Lammi, A. 2001. Consequences of plant population size and density for plant-pollinator interactions and plant performance. *Journal of Ecology* 89: 80-87.

- Namkoong, G.; Boyle, T.J.B.; Gregorius, H-R.; Joly, H.; Savolainen, O.; Wickneswari, R.; Young, A. 1996. CIFOR working paper No. 10. Bogor, Indonesia. Center for International Forestry Research. 12pp.
- Nason, J.D. and Hamrick, J.L. 1997. Reproductive and genetic consequences of forest fragmentation: two case studies of neotropical canopy trees. *Journal of Heredity* 88: 264-276.
- Nason, J.D.; Aldrich, P.R.; Hamrick, J.L. 1997. Dispersal and the dynamics of genetic structure in fragmented tropical tree populations. Pp. 304-320 in Laurance, W.F.; Bierregaard, R.O. (eds.), *Tropical forest remnants: ecology, management and conservation*. Chicago, Illinois. University of Chicago Press.
- Nason, J.D.; Hamrick, J.L. 1997. Reproductive and genetic consequences of forest fragmentation: two case studies of neotropical canopy trees. *Journal of Heredity* 88:246-276.
- Nunney, L. 1993. The influence of mating system and overlapping generations on effective population size. *Evolution* 47: 1329-1341.
- Oficina de Planificación Sectorial Agropecuaria. 1978. Asociación de subgrupos de suelos de Costa Rica. Edición 1, Liberia and Nicoya Sheets. San José, Costa Rica.
- Ohta, T. 1982. Linkage disequilibrium with the island model. *Genetics* 101: 1139-155.
- Oostermeijer, J.G.B.; van Eijik, M.E.; den Nijs, J.C.M. 1994. Offspring fitness in relation to population size and genetic variation in the rare perennial plant species *Gentiana pneumonanthe* (Gentianaceae). *Oecologia* 97: 289-296.
- Pitel, J.A. and Cheliak, W.M. 1984. Effect of extraction buffers on characterization of isoenzymes from vegetative tissues of five conifer species: a users manual. Report PI-X-34, Petawawa National Forestry Inst., Canadian Forestry Service.
- Pittier, H. 1978. Plantas usuales de Costa Rica. Editorial Costa Rica, San José, Costa Rica.
- Porras, A; Villareal, B. 1993. Deforestación en Costa Rica (implicaciones sociales, económicas y legales). San José, Costa Rica. Editorial Costa Rica. 118pp.
- Prober, S.M.; Brown, A.H.D. 1994. Conservation of the grassy white box woodlands: population genetics and fragmentation of *Eucalyptus albens*. *Conservation Biology* 8: 1003-1013.

- Prober, S.M.; Spindler, L.H.; Brown, A.H.D. 1998. Conservation of the grassy white box woodlands: effects of remnant population size on genetic diversity in the allotetraploid herb *Microseris lanceolata*. *Conservation Biology* 12(6): 1279-1290.
- Quirós, C. 1990. La era de la encomienda. San José, Costa Rica. Editorial de la Universidad de Costa Rica, 376pp.
- Raijmann, L.E.; van Leeuwen, N.C.; Kersten, R.; Oostermeijer, J.G.B.; den Nijs, H.C.M.; Menken, S.B.J. 1994. Genetic variation and outcrossing rate in relation to population size in *Gentiana pneumonanthe* L. *Conservation Biology* 8: 1014-1026.
- Ramsey, F.L. and Schafer, D.W. 1997. The statistical sleuth. Belmont, California. Duxbury Press. 742pp.
- Rathke, B.J.; Jules, E.S. 1993. Habitat fragmentation and plant-pollinator interactions. *Current Science* 65(3): 273-277.
- Ridgeway, G.; Sherburne, S.W.; Lewis, R.D. 1970. Polymorphism in the esterases of Atlantic herring. *Transactions of the American Fisheries Society* 99: 147-151.
- Riitters, K; Wickham, J.; O'Neill, R; Jones, B.; Smith, E. 2000. Global-scale patterns of forest fragmentation. *Conservation Ecology* 4(2): 3. [online] URL: <http://www.consecol.org/vol4/iss2/art3>.
- Ritland, K. 1996. Multilocus mating system program MLTR. Program and accompanying documentation. Online at <http://genetics.forestry.ubc.ca/ritland/programs.html>.
- Ritland, K. and Jain, S. 1981. A model for the estimation of outcrossing rate and gene frequencies using n independent loci. *Heredity* 47(1): 35-52.
- Roberds, J.H. and Brotschol, J.V. 1985. Linkage disequilibria among allozyme loci in natural populations of *Liriodendron tulipifera* L. *Silvae Genetica*. 34(4-5): 137-141.
- Rocha, O.J. and Lobo, J.A. Genetic variation and differentiation among five populations of the Guanacaste tree (*Enterolobium cyclocarpum* Jacq.) in Costa Rica. *International Journal of Plant Sciences* 157(2): 234-239.
- Roubik, D.W. (ed.). 1995. Pollination of cultivated plants in the tropics. FAO Agricultural Services Bulletin 118. Rome, Italy. 198p

- Sader, S.A.; Joyce, A.T. 1988. Deforestation rates and trends in Costa Rica, 1940-1983. *Biotropica* 20(1): 11-19.
- Sampson, J.F.; Hopper, S.D.; James, S.H. 1988. Genetic diversity and the conservation of *Eucalyptus crucis* Maiden. *Australian Journal of Botany* 36: 447-460.
- Sánchez-Azofeifa, G.A.; Harriss, R.C.; Skole, D.L. 2001. Deforestation in Costa Rica: a quantitative analysis using remote sensing imagery. *Biotropica* 33(3): 378-384.
- SAS (Statistical Analysis System). 1999. Online documentation Version 8.0.
- Savolainen, O.; Kuittinen, H. 2000. Small population processes. Pp. 91-100 in Young, A.; Boshier, D.; Boyle, T. (eds.). Forest conservation genetics. CABI Publishing, Wallingford, England / CSIRO Publishing, Collingwood, Victoria.
- Schelhas, J. 1996 Land-use choice and forest patches in Costa Rica. Pp. 258-284 in Schelhas, J.; Greenberg, R. (eds.) Forest patches in tropical landscapes. Washington D.C., Island press. 426pp.
- Sequiera, W. 1985. La hacienda ganadera en Guanacaste: aspectos económicos y sociales 1850-1900. UNED, Costa Rica.
- Sizer, N.; Tanner, E.V.J. 1999. Responses of woody plant seedlings to edge formation in a lowland tropical rainforest, Amazonia. *Biological Conservation* 91: 135-142.
- Smith, N.J.H.; Williams, J.T.; Plucknett, D.L.; Talbot, J.P. 1992. Tropical forests and their crops. Ithaca, U.S.A. and London, England. Comstock Publishing Associates (Cornell University Press), 568pp.
- Sokal, R.R. and Rohlf, F.J. 1995. Biometry. New York, Freeman and Company. 887pp.
- Soulé, M.E. (ed.). 1987. Viable populations for conservation. Cambridge University Press, New York.
- SPSS Inc. 1996. SPSS Base 7.0 Applications Guide. Chicago, U.S.A. SPSS Inc., 214pp.
- Standley, P.C. and Williams, L.O. 1969 Flora of Guatemala. Apocynaceae. *Fieldiana* 24(4): 234-407.
- Standley, P.C.; Steyermark, J.A. 1946b. Capparidaceae. Flora of Guatemala. *Fieldiana* 24, Part V.

- Standley, P.C.; Steyermark, J.A. 1946a. Flora of Guatemala. *Fieldiana* 24, Part IV.
- Stevens, G. 1983. *Bursera simaruba*. Pp. 201-202 in Janzen, D.H. (ed.). Costa Rican Natural History. Chicago, University of Chicago Press. 816p.
- Strauss, S.H. and Conkle, M.T. 1986. Segregation, linkage, and diversity of allozymes in knobcone pine. *Theoretical and Applied Genetics* (1986) 72: 483-493.
- Sun, M.; Ritland, K. 1998. Mating system of yellow starthistle (*Centaurea solstitialis*), a successful colonizer in North America. *Heredity* 80: 225-232.
- Van Dyck, H; Matthysen, E. 1999. Habitat fragmentation and insect flight: a changing 'design' in a changing landscape. *Trends in Ecology and Evolution* 14(5): 172-174.
- Van Treuren, R.; Bijlsma, R.; Van Delden; Ouborg, N.J. 1991. The significance of genetic erosion in the process of extinction. I. Genetic differentiation in *Salvia pratensis* and *Scabiosa columbaria* in relation to population size. *Heredity* 66: 181-189.
- Vargas Ulate, G. 2001. Fitogeografía de ecosistemas secos en la meseta de ignimbritas de Guanacaste, Costa Rica. *Revista de Biología Tropical* 49(1): 227-238.
- Vaughton, G.; Ramsey, R. 1995. Pollinators and seed production. Pp. 475-490 in Kigel, J., Galili, J., (eds.), Seed development and germination. New York, USA, Dekker.
- Weeden, N.F. and Wendel, J.F. 1990. Genetics of plant isozymes. Pp 46-72 in Soltis, D.E. and Soltis, P.S. Isozymes in plant biology. London, Chapman and Hall.
- Weir, B.S. 1979. Inferences about linkage disequilibrium. *Biometrics* 35: 235-254.
- Weir, B.S. 1979. Inferences about linkage disequilibrium. *Biometrics* 35: 235-254.
- Weir, B.S. 1996. Genetic data analysis 2. Sunderland, Massachusetts, Sinauer Associates. 445pp.
- Wendel, J.F.; Weeden, N.F. 1990. Visualization and interpretation of plant isozymes. Pp 5-45 in Soltis, D.E. and Soltis, P.S. Isozymes in plant biology. London, Chapman and Hall.
- White, G.M.; Boshier, D.H. 2000. Fragmentation in Central American dry forests: genetic impacts on *Swietenia humilis* (Meliaceae). Pp. 293-311 in Young, A.G. and Clarke, G.M. (eds.), Genetics, demography and viability of fragmented populations. Cambridge, England, Cambridge University Press, 438pp.

- White, G.M.; Boshier, D.H.; Powell, W. 1999. Genetic variation in a fragmented population of *Swietenia humilis*. *Molecular Ecology* 8: 1899-1909.
- Whitlock, M.C.; McCauley, D.E. 1999. Indirect measures of gene flow and migration: $F_{ST} \approx 1/(4Nm+1)$. *Heredity* 82: 117-125.
- Witsberger, D.; Current, D.; Archer, E. 1982. Arboles del Parque Deininger. San Salvador, El Salvador. Ministerio de Educación, 336p.
- Woodson, 1938. Studies in the Apocynaceae. VII. *Annals of the Missouri Botanic Gardens* 25: 189-223.
- Wright, S. 1938. Size of population and breeding structure in relation to evolution. *Science* 87: 430-431.
- Wright, S. 1969. Evolution and the genetics of populations. Volume 2. The theory of gene frequencies. Chicago and London, The University of Chicago Press, 511pp.
- Xie, C.Y.; Dancik, B.P.; Yeh, F.C. 1991. Inheritance and linkage of isozymes in *Thuja orientalis*. *Journal of Heredity* 82: 329-334.
- Yang, R.-C.; Yeh, F.C. 1993. Multilocus structure in *Pinus contorta* Dougl. *Theoretical and Applied Genetics* (1993) 87: 568-576.
- Yeh, F.C. 2000. Population genetics. In Young, A.; Boshier, D.; Boyle, T. (eds.). Forest conservation genetics, pp. 21-37. CABI Publishing, Wallingford, England / CSIRO Publishing, Collingwood, Victoria..
- Yeh, F.C. and Boyle, T.J.B. 1997. Population genetic analysis of codominant and dominant markers and quantitative traits. *Belgian Journal of Botany* 129: 157.
- Yeh, F.C.; Layton, C. 1979. The organization of genetic variability in central and marginal populations of lodgepole pine *Pinus contorta* ssp. *latifolia*. *Canadian Journal of Genetics and Cytology* 21: 487-503.
- Yeh, F.C.; Morgan, K. 1987. Mating system and multilocus associations in a natural population of *Pseudotsuga menziesii* (Mirb.) Franco. *Theoretical and Applied Genetics* (1987) 73: 799-808.

- Yeh, F.C.; Shi, J.; Yang, R.; Hing, J.; Ye, Z. 1994. Genetic diversity and multilocus associations in *Cunninghamia lanceolata* (Lamb.) Hook from The People's Republic of China. *Theoretical and Applied Genetics* (1996) 88: 465-471.
- Young, A.; Boyle, T.; Brown, T. 1996. The population genetic consequences of habitat fragmentation for plants. *Trends in Ecology and Evolution* 11: 413-418.
- Young, A.G.; Brown, A.H.D.; Murray, B.G.; Thrall, P.H.; Miller, C.H. 2000. Genetic erosion, restricted mating and reduced variability in fragmented populations of the endangered grassland herb *Rutidosia leptorrhynchoides*. Pp. 335-359 in Young, A.G. and Clarke, G.M. (Eds.), *Genetics, demography and viability of fragmented populations*. Cambridge, England, Cambridge University Press, 438pp.
- Young, A.G.; Brown, A.H.D.; Zich, F.A. 1999. Genetic structure of fragmented populations of the endangered daisy *Rutidosia leptorrhynchoides*. *Conservation Biology* 13: 256-265.
- Young, A.G.; Merriam, H.G.; Warwick, S.I. 1993. The effects of forest fragmentation on genetic variation in *Acer saccharum* Marsh. (sugar maple) populations. *Heredity* 71: 277-289.
- Zamora, N: 2000. *Cordia bicolor* DC. Online at <http://guestguest@darnis.inbio.ac.cr/FFMPro?-DB=UBI&-lay=WebAll&-Format=/ubi/detail.html&-Op=bw&id=1965&-Find>.
- Zanetto, A.; Kremer, A.; Müller-Starck, G.; Hattemer, H.H. 1996. Inheritance of isozymes in pedunculate oak (*Quercus robur* L.). *Journal of Heredity* 87: 364-370.

APPENDICES

APPENDIX I: TABLES

Table 3-1. Two-way contingency tables of location with respect to nearest watercourse ($\leq 100\text{m}$, $>100\text{m}$) and presence of forest in 1945 (southern sector) and 1945-61 (northern sector)

	Forest cover		Totals
	Unforested	Forested	
Distance to nearest watercourse			
Southern zone			
$\leq 100\text{m}$	11	12	23
$>100\text{m}$	56	21	77
Totals	67	33	100
One-tailed probability (Fisher's exact test): 0.026			
Odds ratio (95 per cent Confidence limits) for distance $\leq 100\text{m}$ v. distance $>100\text{m}$ = 2.909 (1.114, 7.595)			
	Unforested	Forested	Totals
Northern zone			
$\leq 100\text{m}$	6	1	7
$>100\text{m}$	27	5	32
Totals	33	6	39
One-tailed probability (Fisher's exact test): 1.0			
Odds ratio (95 per cent Confidence limits) for distance $\leq 100\text{m}$ v. distance $>100\text{m}$ = 0.9 (0.088, 9.178)			

Table 3-2. Two-way contingency tables of location with respect to nearest watercourse ($\leq 100\text{m}$, $>100\text{m}$) and presence of forest, 1998

	Forest cover		Totals
	Unforested	Forested	
Distance to nearest watercourse			
Southern zone			
$\leq 100\text{m}$	44	23	67
$>100\text{m}$	181	18	199
Totals	225	41	266
One-tailed probability (Fisher's exact test): 0.000			
Odds ratio (95 per cent Confidence limits) (distance $\leq 100\text{m}$ v. distance $>100\text{m}$) = 5.256 (2.612, 10.577)			
	Unforested	Forested	Totals
Northern zone			
$\leq 100\text{m}$	12	9	21
$>100\text{m}$	57	15	72
Totals	69	24	93
One-tailed probability (Fisher's exact test): 0.044			
Odds ratio (95 per cent Confidence limits) (distance $\leq 100\text{m}$ v. distance $>100\text{m}$) = 2.850 (1.013, 8.020)			

Table 3-3. Two-way contingency tables of presence of trees within radius of 27.5m in agricultural and pastureland, 1998

	Land use		Totals
	Agriculture	Pastureland	
Presence of trees in radius of			
Southern zone			
Present	27	62	89
Absent	97	27	124
Totals	124	89	213
One-tailed probability (Fisher's exact test): 0.000			
Odds ratio (95 per cent Confidence limits) (for pasture v. agricultural land) = 8.250 (4.432, 15.357)			
	Agriculture	Pastureland	Totals
Northern zone			
Present	1	36	37
Absent	1	16	17
Totals	2	52	54
One-tailed probability (Fisher's exact test): .535			
Odds ratio (95 per cent Confidence limits) (for pasture v. agricultural land) = 2.25 (.132, 38.27)			

Table 3-4. Two-way contingency tables of presence of forest within 100m of sample points in agricultural and pastureland, 1998

	Land use		Totals
	Agriculture	Pastureland	
Distance to nearest forest			
Southern zone			
≤ 100m	25	41	66
>100m	99	48	147
Totals	124	89	213
One-tailed probability (Fisher's exact test): 0.000			
Odds ratio (95 per cent Confidence limits) (for pasture v. agricultural land) = 3.383 (1.847, 6.195)			
	Agriculture	Pastureland	Totals
Northern zone			
≤ 100m	2	19	21
>100m	0	33	33
Totals	2	52	54
One-tailed probability (Fisher's exact test): 0.147			
Odds ratio (95 per cent Confidence limits) (for pasture v. agricultural land) = 1.105 (.962, 1.270)			

Table 4-1. Segregation ratios in pooled progeny arrays of five enzyme systems of *Anacardium excelsum*

Enzyme	Maternal genotype	Classes for G-test	Observed numbers per class	n_{total}	G_{adj}^1
AK2	AB	AA+BB AB	259 176	435	15.9
LAP	AB	AA+BB AB	247 217	464	1.94
LAP	AC	AA+CC AC	22 25	47	0.19
LAP	AB	AC BC	6 6	12	0.00
PGD	AB	AA+BB AB	321 225	546	16.90
PGM	AB	AA+BB AB	335 334	671	0.001
UGPU	AB	AA+BB AB	181 153	334	2.35

¹Critical value for p of 0.05 = 3.841, with Williams's adjustment (Sokal and Rohlf, 1995)

Table 4-2. Segregation ratios and associated probabilities in progeny of mother trees of *Anacardium excelsum* putatively heterozygous for the AK2 locus.

Population	Family	n_{AA}	n_{BB}	$n_{AA}+n_{BB}$	n_{AB}	n_{total}	n_{exp}^1	p^2
Duquesa	706	5	4	9	5	14	7	0.42
El Cepo	1512	8	3	11	9	20	10	0.82
El Cepo	1521	3	2	5	7	12	6	0.77
El Cepo	1532	5	4	9	11	20	10	0.82
El Cepo	1550	5	4	9	11	20	10	0.82
El Cepo	1551	2	3	5	11	16	8	0.21
El Cepo	1554	15	4	19	0	19	9.5	0.00
El Rodeo	6	5	3	8	10	18	9.0	0.81
La Isla	15	10	3	13	7	20	10	0.26
Las Congojas	4	6	6	12	8	20	10	0.50
Las Congojas	5	10	3	13	6	19	9.5	0.17
Las Congojas	6	1	9	10	2	12	6.0	0.04
Las Congojas	8	4	3	7	5	12	6	0.77
Las Congojas	16	6	7	13	7	20	10	0.26
Las Congojas	17	1	8	9	1	10	5	0.02
Las Congojas	20	8	2	10	10	20	10	1.00
E.J.N.	110	7	1	8	7	15	7.5	1.00
Marcela	416	10	2	12	4	16	8	0.08
Palmira	1	3	5	8	5	13	6.5	0.58
Palmira	4	6	10	16	5	21	10.5	0.03
Paso Hondo	2	4	3	7	6	13	6.5	1.00
Paso Hondo	3	4	1	5	2	7	3.5	0.45
R.S.R.	3	7	2	9	10	19	9.5	1.00
R.S.R.	7a	6	4	10	10	20	10	1.00
R.S.R.	13	8	4	12	8	20	10	0.50
Toroaja	115	8	2	10	9	19	9.5	1.00

¹i.e. expected number in either of either n_{AB} or $n_{AA}+n_{BB}$ under null hypothesis of 1:1 ratio; ²two-tailed exact binomial cumulative probability of n_{AB} if $n_{AB} < n_{AA}+n_{BB}$, or of $n_{AA}+n_{BB}$ if $n_{AB} > n_{AA}+n_{BB}$.

Table 4-3. Segregation ratios and associated probabilities in progeny of mother trees of *Anacardium excelsum* putatively heterozygous for the LAP locus.

Maternal genotype, population	Family	Number of progeny per genotypic class				n_{total}	n_{exp} per class ¹	p^2
		n_{AA}	n_{BB}	$n_{AA}+n_{BB}$	n_{AB}			
AB mothers								
Duquesa	706	2	3	5	13	18	9	0.10
El Cepo	1513	5	2	7	8	15	7.5	1.00
El Cepo	1535	6	1	7	3	10	5	0.34
El Cepo	1566	16	1	17	3	20	10	0.003
El Ojoche	10	2	2	4	6	10	5	0.75
El Ojoche	21	7	7	14	18	32	16	0.60
La Isla	3	2	7	9	11	20	10	0.82
La Isla	4	4	2	6	12	18	9	0.24
La Isla	15	4	3	7	9	16	8	0.80
Las Congojas	5	3	1	4	11	15	7.5	0.12
Las Congojas	9	3	5	8	10	18	9	0.81
Las Congojas	16	10	2	12	6	18	9	0.24
Las Congojas	25	2	2	4	6	10	5	0.75
Marcela	304	2	1	3	4	7	3.5	1.00
Marcela	357	9	1	10	7	17	8.5	0.63
Marcela	371	2	1	3	0	3	1.5	0.25
Marcela	409	3	5	8	9	17	8.5	1.00
Marcela	415	2	6	8	10	18	9	0.81
Marcela	416	6	4	10	8	18	9	0.81
Marcela	418	5	2	7	3	10	5	0.34
Marcela	419	13	1	14	1	15	7.5	0.001
Marcela	503	5	3	8	4	12	6	0.39
Palmira	1	2	1	3	13	16	8.0	0.02
Palmira	2	6	5	10	8	18	9	0.81
Palmira	3	5	7	12	3	15	7.5	0.04
Palmira	4	4	8	12	7	19	9.5	0.36
R.S.R.	0	6	5	11	6	17	8.5	0.33
R.S.R.	5	1	8	9	3	12	6	0.15
R.S.R.	13	4	5	9	11	20	10	0.82
Toronja'	116	5	1	6	4	10	5	0.75
		n_{AC}	n_{BC}					
El Cepo	1554	1	1			2	1	1.00
Las Congojas	5	0	1			1	0.5	1.00
Palmira	3	1	1			2	1.0	1.00
La Isla	4	0	1			1	0.5	1.00
Marcela	357	3	0			3	1.5	0.25
El Ojoche	10	0	2			2	1	0.50
R.S.R.	5	1	0			1	0.5	1.00

¹i.e. expected number in either of the genotypic classes being compared under null hypothesis of 1:1 ratio; ²two-tailed exact binomial cumulative probability of less frequent class.

Table 4-3. Segregation ratios and associated probabilities in progeny of mother trees of *Anacardium excelsum* putatively heterozygous for the LAP locus (continued)

Population, maternal genotypes	Family	Number of progeny per genotypic class				n_{total}	n_{exp} per class	p^1
		n_{AA}	n_{CC}	$n_{AA+n_{CC}}$	N_{AC}			
AC mothers								
El Cepo	1550	2	2	4	6	10	5	0.75
El Cepo	1551	3	6	9	8	17	8.5	1.00
El Ojoche	12	3	2	5	7	12	6	0.77
El Ojoche	16	3	1	4	4	8	4	1.00
BC mothers								
		N_{AC}	N_{BC}					
La Isla	1	2	6	8	3	11	5.5	0.23

¹ p : expected number in either of the genotypic classes being compared under null hypothesis of 1:1 ratio; ²two-tailed exact binomial cumulative probability of less frequent class.

Table 4-4. Segregation ratios and associated probabilities in progeny of mother trees of *Anacardium excelsum* putatively heterozygous for the PGD locus.

Population	Family	n_{AA}	n_{BB}	$n_{AA} + n_{BB}$	n_{AB}	n_{total}	n_{exp}^1	\hat{p}^2
Duquesa	700	8	1	9	11	20	10	0.82
Duquesa	706	2	4	6	8	14	7	0.79
Duquesa	700a	4	3	7	13	20	10	0.26
El Cepo	1511	18	1	19	1	20	10	0.00
El Cepo	1512	8	4	12	8	20	10	0.50
El Cepo	1521	8	2	10	8	18	9	0.81
El Cepo	1535	4	5	9	11	20	10	0.82
El Cepo	1542	4	5	9	10	19	9.5	1.00
El Cepo	1550	5	5	10	10	20	10	1.00
El Cepo	1551	7	1	8	11	19	9.5	0.65
El Cepo	1554	14	1	15	5	20	10	0.04
El Cepo	1556	7	3	10	7	17	8.5	0.63
El Cepo	1566	17	1	18	2	20	10	0.00
Ojoche	2923	9	1	10	4	14	7	0.18
La Isla	1	11	2	13	5	18	9	0.10
La Isla	3	5	4	9	12	21	10.5	0.66
Las Congojas	6	9	5	14	5	19	9.5	0.06
Marcela	371	1	1	2	1	3	1.5	1.00
Marcela	418	7	2	9	1	10	5	0.02
Palmira	1	4	7	11	5	16	8.0	0.21
Palmira	4	10	6	16	4	20	10	0.01
Paso Hondo	2	2	8	10	3	13	6.5	0.09
Paso Hondo	3	1	6	7	0	7	3.5	0.02
R.S.R.	0	8	1	9	7	16	8	0.80
R.S.R.	3	5	3	8	12	20	10	0.50
R.S.R.	5	4	3	7	7	14	7	1.00
R.S.R.	7a	9	4	13	7	20	10	0.26
Toronja	104	7	4	11	9	20	10	0.82
Toronja	107	3	6	9	6	15	7.5	0.61
Toronja	108	2	2	4	1	5	2.5	0.38
Toronja	115	5	1	6	13	19	9.5	0.17
Toronja	116	4	1	5	5	10	5	1.00
Toronja	115	5	1	6	13	19	9.5	0.17

¹i.e. expected number in either of either n_{AB} or $n_{AA} + n_{BB}$ under null hypothesis of 1:1 ratio; ²two-tailed exact binomial cumulative probability of n_{AB} if $n_{AB} < n_{AA} + n_{BB}$, or of $n_{AA} + n_{BB}$ if $n_{AB} > n_{AA} + n_{BB}$.

Table 4-5. Segregation ratios and associated probabilities in progeny of mother trees of *Anacardium excelsum* putatively heterozygous for the PGM1 locus.

Population	Family	n_{AA}	n_{BB}	$n_{AA} + n_{BB}$	n_{AB}	n_{total}	n_{exp}^1	p^2
Duquesa	700	5	4	9	11	20	10	0.82
Duquesa	700A	7	4	11	9	20	10	0.82
El Cepo	1511	4	6	10	10	20	10	1.00
El Cepo	1512	4	5	9	11	20	10	0.82
El Cepo	1521	4	5	9	11	20	10	0.82
El Cepo	1539	1	3	4	6	10	5	0.75
El Cepo	1549	8	8	16	4	20	10	0.01
El Cepo	1550	1	2	3	7	10	5	0.34
El Cepo	1553	5	7	12	8	20	10	0.50
El Cepo	1554	3	1	4	5	9	4.5	1.00
El Cepo	1556	6	3	9	9	18	9	1.00
El Cepo	1566	2	11	13	5	18	9	0.10
El Ojoche	16	2	4	6	9	15	7.5	0.61
El Ojoche	21	7	5	12	20	32	16	0.22
El Ojoche	23	3	5	8	6	14	7	0.79
La Isla	1	7	4	11	7	18	9	0.48
La Isla	3	4	3	7	14	21	10.5	0.19
La Isla	15	4	7	11	9	20	10	0.82
Las Congojas	4	2	1	3	7	10	5	0.34
Las Congojas	6	7	4	11	9	20	10	0.82
Las Congojas	9	6	4	10	10	20	10	1.00
Las Congojas	12	8	3	11	9	20	10	0.82
Las Congojas	17	4	4	8	2	10	5	0.11
Las Congojas	22	10	2	12	8	20	10	0.50
Las Congojas	24	6	1	7	3	10	5	0.34
MAG	110	2	3	5	11	16	8	0.21
Marcela	357	6	6	12	7	19	9.5	0.36
Marcela	415	6	2	8	11	19	9.5	0.65
Marcela	416	3	4	7	11	18	9	0.48
Marcela	418	6	1	7	3	10	5	0.34
Marcela	419	3	4	7	8	15	7.5	1.00
Marcela	503	1	4	5	7	12	6	0.77
Palmira	3	3	4	7	12	19	9.5	0.36
Paso Hondo	1	6	1	7	3	10	5.0	0.34
Paso Hondo	2	7	2	9	4	13	6.5	0.27
R.S.R.	0	2	6	8	9	17	8.5	1.00
R.S.R.	5	1	7	8	5	13	6.5	0.58
R.S.R.	7	3	2	5	15	18	9	0.04
Toronja	115	8	2	10	10	20	10	1.00
Toronja	116	3	1	4	9	13	6.5	0.27

¹i.e. expected number in either of either n_{AB} or $n_{AA} + n_{BB}$ under null hypothesis of 1:1 ratio; ²two-tailed exact binomial cumulative probability of n_{AB} if $n_{AB} < n_{AA} + n_{BB}$, or of $n_{AA} + n_{BB}$ if $n_{AB} > n_{AA} + n_{BB}$.

Table 4-6. Segregation ratios and associated probabilities of putatively heterozygous mother trees of *Anacardium excelsum*, UTP-glucose-1-phosphate uridylyltransferase

Population	Family	n_{AA}	n_{BB}	$n_{AA}+n_{BB}$	n_{AB}	n_{total}	n_{exp}^1	p^2
Duquesa	706	5	1	6	7	13	6.5	1.00
El Cepo	1512	6	2	8	12	20	10	0.50
El Cepo	1535	8	2	10	10	20	10	1.00
El Cepo	1541	9	2	11	9	20	10	0.82
El Cepo	1550	4	4	8	12	20	10	0.50
El Cepo	1557	6	2	8	6	14	7	0.79
El Ojoche	10	2	3	5	8	13	6.5	0.58
El Ojoche	12	2	3	5	5	10	5	1.00
Las Congojas	1	4	2	6	10	16	8	0.45
Las Congojas	4	4	3	7	13	20	10	0.26
Las Congojas	8	2	1	3	9	12	6	0.15
Las Congojas	20	10	3	13	7	20	10	0.26
Marcela	415	11	1	12	5	17	8.5	0.14
Marcela	416	7	2	9	9	18	9	1.00
Marcela	418	5	1	6	4	10	5	0.75
Palmira	1	5	4	9	6	15	7.5	0.61
Palmira	2	7	8	15	5	20	10	0.04
Palmira	4	8	6	14	7	21	10.5	0.19
R.S.R.	4	6	6	12	6	18	9	0.24
R.S.R.	5	11	1	12	2	14	7	0.01
R.S.R.	9	1	1	2	1	3	1.5	1.00

¹i.e. expected number in either of either n_{AB} or $n_{AA}+n_{BB}$ under null hypothesis of 1:1 ratio; ²two-tailed exact binomial cumulative probability of n_{AB} if $n_{AB} < n_{AA}+n_{BB}$, or of $n_{AA}+n_{BB}$ if $n_{AB} > n_{AA}+n_{BB}$. i.e. of either n_{AB} or $n_{AA}+n_{BB}$.

Table 4-7. Significant estimates of Burrows's composite linkage disequilibrium coefficient, correlation coefficient and significance of associated chi-squared test in populations of *Anacardium excelsum* in Guanacaste province, Costa Rica.

Loci and alleles ¹	Number +/- ²	Population	Δ_y	r	p	n
AK2-A v. LAP-A	6/6	Toronja:	-.0564	-.2514	.0165	91
AK2-A v. UGPU2-A	10/0 ³	Duquesa	.0252	.2627	.0258	73
		Cepo	.0350	.1526	.0061	319
		Santa Rosa	.0439	.1952	.0168	148
AK2-A v. PGD-A	9/3	La Isla	0.0736	0.2746	0.0280	64
		Paso Hondo	-0.1104	0.3462	0.0102	55
LAP-A v. PGM1-A	8/4	Duquesa	-0.0532	-0.27	0.0220	72

¹All alleles except the most frequent were pooled to one synthetic allele; ²i.e. number of positive and negative correlations, irrespective of significance, over all 12 populations; ³total is < total number of populations (12) because $p=1$ (>0.9999) in remaining populations, i.e. disequilibrium=0

Table 4-8. Two by three classification of *Anacardium excelsum* progeny by PGD putative genotype and presence of putative AK1-A allele, with result of chi-square test of association.

	Number of individuals per progeny class		
	AK1-A present	AK1-A absent	totals
PGD putative genotypes			
AA	917	41	958
AB	108	171	279
BB	9	117	126
totals	1034	329	1363
Chi ² , df, p:	741.55, d.f.=2, $p < 0.0001$		

Table 4-9. Estimates of Ohta's multiple population linkage disequilibrium coefficients for 12 populations of *Anacardium excelsum* in northwestern Costa Rica.

Loci	D_{IT}^2	D_{IS}^2	D'_{IS}^2	D_{ST}^2	D'_{ST}^2
AK2/LAP	0.16675	0.00156	0.15416	0.16216	0.01259
AK2/PGD	0.13579	0.00844	0.12970	0.12196	0.00609
AK2/PGM	0.12596	0.00386	0.11996	0.12666	0.00599
AK2/UGPU	0.12176	0.00177	0.11533	0.11955	0.00644
LAP/PGD	0.14757	0.00106	0.14351	0.14080	0.00406
LAP/PGM	0.12940	0.00114	0.12655	0.12781	0.00285
LAP/UGPU	0.13974	0.00034	0.13529	0.13852	0.00445
PGD/PGM	0.09803	0.00108	0.09750	0.09385	0.00053
PGD/UGPU	0.06787	0.00034	0.06611	0.06795	0.00176
PGM/UGPU	0.08043	0.00149	0.07894	0.08241	0.00149
Overall	0.11035	0.00146	0.10619	0.10802	0.00416

Table 4-10. Results of Ewens-Watterson test of selective neutrality for 12 populations of *Anacardium excelsum* in northwestern Costa Rica.

Locus	Parameters	Population ¹											
		D	C	O	R	I	Co	E	M	P	PH	S	T
AK2	n ²	146	632	246	42	128	520	32	302	148	114	310	228
	F _{obs} ³	0.82	.64	.98	.50	.51	.59	.60	.88	.65	.62	.69	.76
	95% limits ⁴	.50-.99	.51-1.0	.50-.99	.50-.95	.50-.98	.50-1.0	.50-.94	.50-.99	.50-.99	.50-.99	.50-.98	.50-.99
LAP	n ²	146	718	218	42	148	508	34	302	142	108	270	188
	F _{obs} ³	.65	.80	.50	.95	.51	.76	.84	.53	.50	.76	.61	.53
	95% limits ⁴	.50-.99	.51-1.0	.50-.99	.50-.95	.50-.99	.51-1.0	.50-.94	.50-.99	.50-.99	.50-.99	.50-.98	.50-.99
PGD	n ²	148	788	254	42	162	532	34	306	150	114	312	226
	F _{obs} ³	0.56	.68	.87	.91	.67	.84	.94	.83	.57	.60	.69	.54
	95% limits ⁴	.50-.99	.50-1.0	.50-.99	.50-.95	.50-.99	.51-1.0	.50-.94	.50-.99	.50-.99	.50-.99	.50-.98	.50-.99
PGM	n ²	146	582	248	42	162	408	34	306	144	104	286	232
	F _{obs} ³	0.60	.58	.52	.66	.51	.56	.50	.52	.71	.60	.70	.54
	95% limits ⁴	.50-.99	.50-.99	.50-.99	.50-.95	.50-.99	.50-1.0	.50-.94	.50-.99	.50-.99	.50-.99	.50-.98	.50-.99
UGPU2	n ²	146	822	242	M ⁵	160	532	M ⁵	292	152	116	306	234
	F _{obs} ³	0.88	.74	.77		.98	.75		.81	.54	.98	.62	.93
	95% limits ⁴	.50-.99	.50-1.0	.51-.99		.50-.99	.50-1.0		.50-.99	.50-.99	.50-.98	.50-.99	.50-.99

¹Population abbreviations: D=Bosque Durquesa, C=El Capon, O=El Ochoque, R=El Rodeo, I=La Isla, Co=Las Conchas, E=EJN, M=Marcel, P=Palmar, PH=Paseo Honda, S=R.S.R.; T=Torrejón; number of alleles in sample; observed F_i 95 per cent confidence limits of sampling distribution ³M=monomorphic

Table 4-11. Estimates of proportions of extraneous seed in collections of putatively heterozygous mother trees of *Anacardium excelsum*.

Population/ Locus	N_{XY} ¹	$N_{XX}:N_{XY}$ (arrays) ²	$XX:XY$ (arrays) ³	$XX:XY$ (pop) ⁴	p_e ⁵
El Cepo					
AK2	5 ⁶	39:45	0.46:0.54	0.78:0.22	-ve
PGD	8 ⁷	83:70	0.54:0.46	0.76:0.24	0.15
PGM	10	89:76	0.54:0.46	0.67-0.33	0.23
UGPU	5	45:49	0.48:0.52	0.80-0.20	-ve
El Ojoche					
PGM	3	26:35	0.43:0.57	0.47-0.53	>1.0
Las Congojas					
AK2	5 ⁸	55:36	0.60:0.40	0.77-0.23	0.37
LAP	4	28:33	0.46:0.54	0.82:0.18	-ve
PGM	7	62:48	0.56:0.44	0.58-0.42	0.75
UGPU	4	29:39	0.43:0.57	0.77:0.23	-ve
Marcela					
LAP ²	8 ⁹	57:45	0.56:0.44	0.83:0.17	0.18
PGM	6	46:47	0.49:0.51	0.56:0.44	-ve
UGPU	3	27:18	0.60:0.40	0.84:0.16	0.29
R.S.R.					
AK	3	31:28	0.52:0.48	0.69:0.31	0.11
PGD	4	37:33	0.53:0.47	0.76:0.24	0.12
PGM	3	21:29	0.42:0.58	0.70:0.30	-ve
Toronja					
PGD	6	41:47	0.46:0.54	0.56:0.44	-ve

¹Number of putative heterozygous mother-trees; ²Numbers of homozygotes and heterozygotes in collections from putative heterozygotes; ³Ratio of homozygotes to heterozygotes in collections from putative heterozygotes heterozygotes; ⁴Ratio of homozygotes to heterozygotes in population as a whole; ⁵estimated proportion of extraneous seed; ⁶Tree 1554 excluded because extreme segregation ratio suggests spurious heterozygous designation; ⁷Tree 1554 excluded because extreme segregation ratio suggests spurious heterozygous designation; ⁸Trees 1511, 1566 excluded; ⁹Trees 6, 17excluded; ¹⁰Tree 419 excluded

Table 4-12. Results of simulation of effect on estimation of mating system parameters of presence of extraneous seed in progeny arrays for combinations of two initial outcrossing rate scenarios ($t_m=1.0$, $t_m=0.7$), three allele frequency scenarios and four seed admixture scenarios

Case ¹	Allele Frequencies	Settings (N, θ) ²	Parameter estimates (s.d.) before admixture	Parameter estimates (s.d.) after admixture		Number of aborted runs ³
				F	t_m	
1.1	.85, .10, .05 .48, .52 .44, .56	1: 20, 10 2: 20, 2 3: 2, 10	F=-.038 (.149), $t_m=1.112$ (.036)	1: -0.97 (0.039)	1: 1.31 (0.072)	0
				2: 2.036 (0.12)	2: 1.14 (0.020)	0
				3: -0.14 (0.054)	3: 1.11 (0.017)	0
1.2	.77, .15, .08 .49, .51 .83, .17	4: 2, 2 1: 20, 10 2: 20, 2 3: 2, 10 4: 2, 2	F=-.169 (.136), $t_m=1.027$ (.043)	4: -0.06 (0.024)	4: 1.11 (0.005)	0
				1: -0.96 (0.071)	1: 1.21 (0.041)	1
				2: -0.52 (0.106)	2: 1.02 (0.015)	2
1.3	.81, .12, .07 .82, .18 .87, .13	4: 2, 2 1: 20, 10 2: 20, 2 3: 2, 10 4: 2, 2	F=.11 (.366), $t_m=1.003$ (.031)	4: -0.195 (0.05)	4: .99 (0.009)	0
				1: -0.98 (0.025)	1: 1.14 (0.037)	0
				2: -0.99 (0.000)	2: 1.01 (0.033)	1
2.1	.79, .14, .07 .58, .42 .48, .52	4: 2, 2 1: 20, 10 2: 20, 2 3: 2, 10 4: 2, 2	F=-.028 (.145), $t_m=.759$ (.039)	4: 0.11 (0.006)	4: 1.005 (0.002)	1
				1: -0.98 (0.02)	1: 1.11 (0.081)	3
				2: -0.42 (0.131)	2: .81 (0.049)	0
2.2	.84, .11, .05 .44, .56 .87, .13	4: 2, 2 1: 20, 10 2: 20, 2 3: 2, 10 4: 2, 2	F=-.021 (.17), $t_m=.716$ (.036)	3: -0.11 (0.045)	3: .78 (0.018)	2
				4: 0.058 (0.018)	4: .76 (0.007)	0
				1: -0.99 (0.00)	1: 1.08 (0.039)	4
2.3	.83, .14, .03 .86, .14 .82, .18	4: 2, 2 1: 20, 10 2: 20, 2 3: 2, 10 4: 2, 2	F=.176 (.248), $t_m=.75$ (.054)	2: -0.51 (0.083)	2: .826 (0.019)	2
				3: -0.11 (0.05)	3: .73 (0.02)	1
				4: -0.07 (0.046)	4: .72 (0.01)	0
				1: -0.98 (0.076)	1: 1.0 (0.023)	0
				2: -0.96 (0.058)	2: 0.78 (0.060)	0
				3: 0.105 (0.027)	3: 0.79 (0.017)	0
				4: 0.119 (0.053)	4: .74 (0.014)	0

¹Case 1.n refers to outcrossing rate settings of $t_m=1.0$, n defines allele frequency settings (see column 3), case 2.n refers to $t_m=0.7$; 2N=number of trees affected by admixture, e=number of admixed seed per affected tree, accordingly, designations 1,2,3 and 4 refer to generalized and intense seed admixture, generalized but light, sporadic but intense, sporadic and light, respectively; %e, due to presence of 3 homozygotic types in ≥ 1 array

Table 5-1. Description of populations sampled in a study of genetic effects of forest fragmentation on *Anacardium excelsum* in northwestern Costa Rica.

Population	coordinates ¹	N ²	isolation ³	disturbance ⁴	matrix ⁵	ndf ⁶ , linearity ⁷
Corobici Group						
El Cepo	85°07.5', 10°28.5'	500+	L	L	P	24.2, 2.9
El Ojoché	85°06.3', 10°29.6'	23	H (5)	M	P	17.0, 1.87
Las Congojas	85°06.6', 10°28.5'	50	M (10)	M	P	34.5, 1.92
Oldemar	85°06.7', 10°26.4'	15	M	H	P	n.d., riparian strip
Quebrada Lajero	85° 06.6', 10°29'	54	M (30)	H	P	n.d., 3.1
River Santa Rosa	85°06.3', 10°27.6'	19	H (5)	H	A/P	7.57, 15.0
Taboga Group						
Bosque Duquesa	85°07', 10°19.5'	9	M (6)	L	P, S	5.0, 2.28
COSTASEM	85°09', 10°18.8'	10	M (4)	H	S, A	5.5, 5.25
E.J.N.	85°08.5', 10°20'	12	H (0)	L	P, A	22.4, 2.96
El Rodeo	85°04.3', 10°19'	6	H (n.d.)	H	P, S	7.2, 10.7
Hacienda Los Rios	85°05', 10°20'	15	H (0)	H	P	10.5, 9.44
La Gotera	85°04.2', 10°19.9'	109	L	L	P	2.4, 3.7
Las Avispas	85°08.2', 10°17.5'	27	H (0)	H	S	25.7, 5.22
Marcela	85°06', 10°18.2'	285	80 (M)	L	S	33.7, 2.96
Quebrada Duquesa	85°06.5', 10°20.1'	10	H (1)	M	P, S	4.14, 4.7
R. Reventado (N)	85°05', 10°19.2'	33	M (17)	H	P	n.d., 5.0
R. Reventado (S)	85°06.0', 10°18.6'	57	L (100)	M	S	n.d., 2.33
Sitio Caucante	85°08.6', 10°19'	15	H (10)	H	S	5.9, 3.25
Toroja	85°09.3', 10°18.3'	34	H (0)	H	S	44.6, 1.06

¹At population centre; ²Population size; ³L=low, M=medium, H=high; number in parentheses is estimated number of extrafragment trees within 500m of population centre; ⁴L=low, M=medium, H=high; ⁵P=pastureland, S=sugar-cane, A=other agriculture; ⁶neighbourhood density index (see text); ⁷l/a, population linearity; l/w, where l = length of a line drawn between the two most distant points of the population and w = maximum width of the population measured perpendicular to line l).

Table 5-1. Description of populations sampled in a study of genetic effects of forest fragmentation on *Anacardium excelsum* in northwestern Costa Rica (continued)

	population	coordinates ¹	N ²	isolation ³	disturbance ⁴	matrix ⁵	ndi ⁶ , linearity ⁷
Other fragments							
	Canatca	85°09.6', 10°31.7'	500+	L	L	P	n.d., riparian strip
	Hacienda Tenorio	85°06', 10°33.2'	500+	L	L	P	n.d., riparian strip
	La Isla	85°06.9', 10°23.7'	18	H (O)	M-H	S	10.6, 4.3
	Palmita	85°05.6', 10°33'	4	L (500)	H	P	4.25, 2.5
	Paso Hondo	85°10.5', 10°23.3'	4	H (O)	H	S	1.0, 14.75
	Q. Salitral (O)	85°06.2', 10°23.7'	17	M	H	S	n.d., riparian strip
	Q. Salitral (Libertad)	85°06.5', 10°22.9'	22	M	H	S	n.d., riparian strip

¹At population centre; ²Population size; ³L=low, M=medium, H=high, number in parentheses is estimated number of extant fragment trees within 500m of population centre; ⁴L=low, M=medium, H=high; ⁵P=pastureland, S=sugar-cane, A=other agriculture; ⁶neighbourhood density index (see text); ⁷l₁, population linearity; l₂/w, where l = length of a line drawn between the two most distant points of the population and w = maximum width of the population measured perpendicular to line l).

Table 5-2. Sampling of populations of in a study of effects of forest fragmentation on genetic and reproduction of *Anacardium excelsum* in northwestern Costa Rica.

population	genetic diversity ¹		mating system parameters		flowering study
	NF ²	NI ³	NF	NI	NI
Corobicí group					
El Cepo	19	420	19	420	10
El Ojoche	8	128	5	88	8
Las Congojas	20	270	17	259	7
Oldemar	0	0	0	0	11
Quebrada Lajero	0	0	0	0	7
River Santa Rosa	10	158	10	158	10
Taboga group					
Bosque La Duquesa	4	74	4	74	5
COSTASEM	0	0	0	0	7
E.J.N. Experiment Stn.	2	17	0	0	13
El Rodeo	3	21	0	0	4
Hacienda Los Ríos	0	0	0	0	12
La Gotera	0	0	0	0	10
Las Avispas	0	0	0	0	16
Marcela fragment	14	154	12	150	10
Q. La Duquesa	0	0	0	0	6
Quebrada Reventado (N)	0	0	0	0	10
Quebrada Reventado (S)	0	0	0	0	9
Sitio Cascante	0	0	0	0	10
Toronja fragment	12	117	6	71	10
Other fragments					
Canateca	0	0	0	0	10
Hacienda Tenorio	0	0	0	0	10
La Isla	5	81	5	81	9
Palmira	4	72	4	72	0
Paso Hondo	4	50	4	50	4
Salitral (Libertad)	0	0	0	0	11

¹Sample sizes were higher for genetic diversity as progeny of indeterminate maternity were included, i.e. those collected beneath crowns of >1 mother tree. ²Number of families, i.e. mother-trees; ³Number of individual progeny

Table 5-3. Representation by block of seed sources (fragments) and open-pollinated families within fragments in a common garden experiment of *Anacardium excelsum*

Fragment/ Family	Block									
	1	2	3	4	5	6	7	8	9	10
El Cepo										
2	1	0	0	0	0	0	0	0	0	0
5	1	1	1	0	0	1	1	0	1	0
9	1	1	0	0	1	1	1	1	0	0
11	1	1	1	1	0	1	0	1	1	1
12	0	1	1	1	1	1	1	1	1	1
19	0	1	1	1	0	1	1	1	1	1
21	1	1	0	0	1	1	1	1	1	1
32	0	1	1	1	1	1	1	1	1	1
35	1	1	1	1	1	1	1	1	1	1
39	1	0	1	1	1	1	1	1	1	1
41	0	1	0	0	1	1	0	1	1	0
42	0	1	1	1	1	1	1	1	1	1
50	1	1	0	1	1	1	1	1	1	0
51	1	1	0	0	0	1	1	1	1	1
53	1	1	0	0	1	1	1	1	1	1
54	1	1	1	0	0	1	0	1	0	0
56	0	1	1	1	1	1	1	0	0	1
57	1	1	1	0	1	1	0	1	0	1
66	1	1	1	1	1	1	1	1	1	1
Bosque Duquesa										
700	1	0	1	1	1	1	1	1	0	0
701	1	1	1	1	0	1	1	1	1	1
706	0	0	0	0	1	0	1	0	0	1
700a	1	1	1	1	1	1	1	1	1	1
Toronja										
100	0	1	1	1	0	1	1	1	0	0
104	1	1	1	0	0	0	1	1	1	0
115	1	1	1	1	1	1	1	1	1	1
116	1	1	1	1	1	0	1	0	0	1
Marcela										
355	0	1	1	1	0	1	0	1	1	1
357	1	1	1	0	0	1	1	0	1	0
409	0	1	0	1	0	0	0	1	1	0
415	1	1	1	0	0	0	1	1	1	0
416	1	1	0	0	1	0	0	0	0	1
419	1	1	1	0	0	0	0	0	1	0
Congojas										
1	0	0	0	0	1	1	0	0	0	0
4	1	1	1	1	1	1	0	1	1	1
5	1	1	1	1	1	1	1	1	1	1
6	0	1	1	1	1	0	0	1	1	1
8	1	1	1	0	1	1	1	1	1	0
9	1	1	1	1	1	0	0	1	1	1
11	1	0	0	0	1	0	0	0	0	0
12	1	1	1	0	1	0	1	1	1	1
16	1	1	1	1	0	1	1	0	1	1

Table 5-3. Representation by block of seed sources (fragments) and open-pollinated families within fragments in a common garden experiment of *Anacardium excelsum* (continued)

Fragment/ Family	Block									
Congojas										
17	1	1	1	1	1	1	1	1	1	1
20	1	1	1	0	1	1	1	0	1	0
22	1	1	1	0	1	1	1	0	1	1
23	1	1	1	0	0	1	1	1	0	1
25	1	1	1	1	1	1	1	1	1	1
R.S.R.										
2	0	0	1	0	1	0	1	0	1	0
3	1	1	1	1	1	0	1	1	0	1
4	1	1	4	0	1	1	1	1	0	0
7	1	1	7	1	1	1	1	1	1	0
13	1	1	1	0	1	1	1	1	1	0
1000	0	1	1	1	0	1	1	1	1	0
2000	0	0	1	1	1	1	1	0	0	0
Ojoche										
10	1	0	0	0	1	1	0	0	0	1
12	1	1	0	1	1	0	0	1	1	1
16	0	1	0	0	1	1	0	1	1	1
19	1	0	1	0	0	1	1	1	0	1
21	1	1	1	1	1	1	1	1	0	0
Isla										
3	1	1	0	1	1	1	1	1	1	1
4	1	1	0	1	1	1	1	1	1	1
15	1	1	0	1	1	1		1	1	1
Palmira										
1	1	1	1	1	1	1	1	1	1	1
2	1	1	1	1	1	1	1	1	1	1
3	1	1	1	1	1	1	1	1	1	1
4	1	0	1	1	1	0	1	1	1	1

Table 5-4. Estimates of allele frequencies, allelic richness, expected heterozygosity, fixation indices and G-test for Hardy-Weinberg disequilibria in populations of *Anacardium excelsum* from 10 forest fragments in northwestern Costa Rica, based on inferred maternal genotypes.

Population / locus	allele frequencies			A^1	H_e^2	F_{is}^3
	p_A	q_B	r_C			
Bosque Duquesa						
AK2	0.88	0.12	n.a.	2	0.22	-0.14
LAP	0.75	0.25	0.0	2	0.38	-0.33
PGD	0.62	0.38	n.a.	2	0.47	-0.60
PGM	0.75	0.25	n.a.	2	0.38	-0.33
UGPU	0.88	0.12	n.a.	2	0.22	-0.14
means (S.D.)				2.0(.00)	0.33(0.11)	
El Cepo						
AK2	0.76	0.24	n.a.	2	0.36	0.02
LAP	0.82	0.13	0.05	3	0.30	-0.21
PGD	0.72	0.28	n.a.	2	0.40	-0.38*
PGM	0.31	0.69	n.a.	2	0.42	-0.44*
UGPU	0.82	0.18	n.a.	2	0.29	0.13
means (S.D.)				2.2 (.45)	0.35 (0.06)	
El Ojoche						
AK2	0.92	0.08	n.a.	2	0.15	-0.09
LAP	0.50	0.25	0.25	3	0.63	-1.0***
PGD	0.83	0.17	n.a.	2	0.29	-0.20
PGM	0.58	0.42	n.a.	2	0.49	-0.03
UGPU	0.82	0.18	n.a.	2	0.28	-0.20
means (S.D.)				2.20 (.45)	0.37 (0.19)	
La Isla						
AK2	0.62	0.37	n.a.	2	0.47	0.47
LAP	0.50	0.40	0.10	3	0.58	-0.20
PGD	0.70	0.30	n.a.	2	0.42	-0.43
PGM	0.50	0.50	n.a.	2	0.50	-0.20
UGPU	1.00	0.00	n.a.	1	0.00	M
means (S.D.)				2.0 (0.71)	0.39(0.23)	
Las Congojas						
AK2	0.76	0.24	n.a.	2	0.36	-0.31
LAP	0.84	0.16	0.00	2	0.26	-0.18
PGD	0.91	0.09	n.a.	2	0.16	-0.10
PGM	0.53	0.47	n.a.	2	0.50	-0.61*
UGPU	0.88	0.12	n.a.	2	0.21	-0.13
means (S.D.)				2.0 (0.00)	0.29 (.13)	

¹Allelic richness; ²Nei's expected heterozygosity (gene diversity); ³fixation index. Asterisks indicate significance of associated G-test of genotypic frequencies (*=0.05, **=0.01, *** \leq 0.001); M = monomorphic.

Table 5-4. Estimates of allele frequencies, allelic richness, expected heterozygosity, fixation indices and G-test for Hardy-Weinberg disequilibria in populations of *Anacardium excelsum* from 10 forest fragments in northwestern Costa Rica, based on inferred maternal genotypes (continued)

population locus	allele frequencies			A^1	H_e^2	F_{is}^3
	p_A	q_B	r_C			
Marcela						
AK2	0.96	0.03	n.a.	2	0.07	-0.04
LAP	0.46	0.54	0.0	2	0.50	-0.58*
PGD	0.85	0.14	n.a.	2	0.24	0.42
PGM	0.61	0.39	n.a.	2	0.48	-0.05
UGPU	0.89	0.11	n.a.	2	0.19	-0.12
means (S.D.)				2.0(.00)	0.29(0.19)	
Palmira						
AK2	0.25	0.75	n.a.	2	0.38	-0.33
LAP	0.50	0.50	0.0	2	0.50	-1.00*
PGD	0.25	0.75	n.a.	2	0.38	-0.33
PGM	0.12	0.88	n.a.	2	0.22	-0.14
UGPU	0.62	0.38	n.a.	2	0.47	-0.60
means (S.D.)				2.0 (.00)	0.39 (0.11)	
Paso Hondo						
AK2	0.25	0.75	n.a.	2	0.38	-0.33
LAP	0.75	0.25	0.0	2	0.38	-0.33
PGD	0.75	0.25	n.a.	2	0.38	-0.33
PGM	0.75	0.25	n.a.	2	0.38	-0.33
UGPU	1.00	0.00	n.a.	1	0.00	M
means (S.D.)				1.80 (.45)	0.30 (0.17)	
River Santa Rosa						
AK2	0.80	0.20	n.a.	2	0.32	-0.25
LAP	0.78	0.16	0.06	3	0.36	-0.29
PGD	0.75	0.25	n.a.	2	0.38	-0.33
PGM	0.15	0.85	n.a.	2	0.25	-0.18
UGPU	0.70	0.30	n.a.	2	0.42	0.05
means (S.D.)				2.2 (0.45)	0.35(0.06)	
Toronja						
AK2	0.93	0.07	n.a.	2	0.13	-0.08
LAP	0.25	0.58	0.17	3	0.57	0.55
PGD	0.57	0.43	n.a.	2	0.49	-0.75
PGM	0.43	0.57	n.a.	2	0.49	0.42
UGPU	1.00	0.00	n.a.	1	0.00	M
means (S.D.)				2.0 (0.71)	0.34 (.25)	

¹Allelic richness; ²Nei's expected heterozygosity (gene diversity); ³fixation index. Asterisks indicate significance of associated G-test of genotypic frequencies (*=0.05, **=0.01, *** \leq 0.001); M = monomorphic.

Table 5-5. Homogeneity statistics for five loci of inferred maternal genotypes in populations of *Anacardium excelsum* from 10 forest fragments in northwestern Costa Rica

locus	chi-square, df, probability	G, df, probability
Overall		
AK	36.0, 9, <0.0001	33.6, 9, <0.0001
LAP	28.7, 9, <0.0001	28.8, 9, <0.0001
PGD	21.0, 9, 0.01	20.0, 9, 0.02
PGM	24.2, 9, 0.004	25.8, 9, 0.002
UGPU	13.7, 9, 0.13	16.6, 9, 0.05
Corobicí Group		
AK	1.43, 3, 0.69	1.69, 3, 0.64
LAP	6.8, 3, 0.08	5.8, 3, 0.12
PGD	4.5, 3, 0.21	4.9, 3, 0.18
PGM	10.3, 3, 0.01	10.9, 3, 0.01
UGPU	2.87, 3, 0.41	2.71, 3, 0.44
Taboga group		
AK	0.92, 2, 0.63	0.83, 2, 0.66
LAP	4.84, 2, 0.09	5.04, 2, 0.08
PGD	4.61, 2, 0.10	4.63, 2, 0.10
PGM	2.35, 2, 0.31	2.39, 2, 0.30
UGPU	1.72, 2, 0.42	2.78, 2, 0.25

Table 5-6. Estimates of Wright's statistics and mN for five loci of inferred maternal genotypes in populations of *Anacardium excelsum* from 10 forest fragments in northwestern Costa Rica

locus	sample size	F_u	F_w	F_{st}	mN
Overall					
AK	174	-0.0964	0.2400	0.3068	0.56
LAP	176	-0.3913	-0.2069	0.1325	1.63
PGD	182	-0.3692	-0.1632	0.1505	1.41
PGM	174	-0.1825	-0.0279	0.1779	1.15
UGPU	182	-0.1736	-0.0311	0.1214	1.81
Mean	178	-0.2594	-0.0359	0.1775	1.16
			Corobici group		
AK	100	-0.1658	-0.1361	0.0255	9.6
LAP	102	-0.3557	-0.2570	0.0728	3.2
PGD	106	-0.2865	-0.2422	0.0344	7.0
PGM	98	-0.3294	-0.1586	0.1285	1.7
UGPU	106	-0.0205	0.0095	0.0294	8.2
Mean	102	-0.2415	-0.1613	0.0646	3.6
			Talaboga group		
AK	50	-0.1047	-0.0839	0.0189	13.0
LAP	48	-0.2385	0.0814	0.1268	1.7
PGD	50	-0.4542	-0.3506	0.0712	3.3
PGM	50	0.0418	0.1106	0.0718	3.2
UGPU	50	-0.1322	-0.0839	0.0427	5.6
Mean	50	-0.1866	-0.0878	0.0832	2.8

Table 5-7. Pairwise estimates of Nei's unbiased genetic distance (below diagonal) and geographic distances (km, above diagonal) between ten populations of *Anacardium excelsum* located in northwestern Costa Rica, based on inferred maternal genotypes.

	Duquesa	Cepo	Ojoche	Isla	Congojas	Marcela	Paso Hondo	Palmira	R.S.R.	Toronja
Duquesa	0	17	17.25	7.9	16.75	2.75	9.6	24.75	15.25	4.5
Cepo	0.0533	0	2.92	9.25	1.85	18	11.1	8.1	3.1	19.3
Ojoche	0.0133	0.0457	0	11	2	21	13.75	6.35	3.6	21.5
Isla	0.0262	0.0367	0.0215	0	8.85	10.25	6.75	17.1	7.25	10.85
Congojas	0.0244	0.0197	0.0221	0.0298	0	19	11.85	8.25	1.7	19.4
Marcela	0.0281	0.0858	0.0063	0.0306	0.0491	0	12.5	27.3	17.25	5.8
Paso Hondo	0.3565	0.2211	0.4055	0.1934	0.3595	0.4182	0	19.3	11	9.6
Palmira	0.1	0.1439	0.1587	0.0451	0.0907	0.1813	0.259	0	10	27.8
R.S.R.	0.1111	0.0028	0.0736	0.0783	0.0534	0.115	0.2026	0.2315	0	18
Toronja	0.0739	0.1073	0.0442	0.0235	0.1272	0.0349	0.2754	0.2388	0.1293	0

Table 5-8. Estimates of allele frequencies, allelic richness, expected heterozygosity, fixation indices and G-test for Hardy-Weinberg disequilibria in populations of *Anacardium excelsum* from 12 forest fragments in northwestern Costa Rica (progeny data)

population / locus	Allele frequencies			A^1	$\hat{\lambda}_m^2$	H_e^3	F_{is}^4
	p_A	q_B	r_C				
Bosque Duquesa							
AK2	0.90	0.10	n.a.	2	1.56	0.18	0.48***
LAP	0.77	0.23	0.0	2	1.88	0.35	-0.06
PGD	0.67	0.33	n.a.	2	1.96	0.44	-0.01
PGM	0.73	0.27	n.a.	2	1.93	0.40	0.17
UGPU	0.94	0.06	n.a.	2	1.41	0.11	0.17
means (S.D.)				2.0(.00)	1.75	0.30(0.14)	0.15
El Cepo							
AK2	0.72	0.28	n.a.	2	2.00	0.40	0.46***
LAP	0.8	0.06	0.05	3	2.72	0.21	0.22***
PGD	0.80	0.20	n.a.	2	2.00	0.32	0.25***
PGM	0.30	0.70	n.a.	2	2.00	0.42	0.22***
UGPU	0.84	0.16	n.a.	2	2.00	0.28	0.28***
means (S.D.)				2.2 (.45)	2.14	0.32 (0.09)	0.29
El Ojoche							
AK2	0.99	0.01	n.a.	2	1.14	0.02	-0.01
LAP	0.47	0.35	0.18	2	2.91	0.63	0.04
PGD	0.93	0.07	n.a.	2	1.59	0.13	0.04
PGM	0.60	0.40	n.a.	2	2.00	0.48	-0.11
UGPU	0.87	0.13	n.a.	2	1.82	0.23	0.28**
means (S.D.)				2.2 (.45)	1.89	0.30 (0.25)	0.05
El Rodeo							
	p_A	q_B	r_C				
AK2	0.48	0.52	n.a.	2	n.a.	0.50	0.04
LAP	0.98	0.02	0.0	2	n.a.	0.05	-0.02
PGD	0.95	0.05	n.a.	2	n.a.	0.09	-0.05
PGM	0.21	0.79	n.a.	2	n.a.	0.34	0.57**
UGPU	1.00	0.0	n.a.	1	n.a.	0.00	M
means (S.D.)				1.80 (.45)	n.a.	0.20 (0.21)	0.14
La Isla							
AK2	0.56	0.44	n.a.	2	2.00	0.49	0.59***
LAP	0.42	0.52	0.06	3	2.48	0.55	-0.05
PGD	0.79	0.21	n.a.	2	1.91	0.33	0.18
PGM	0.43	0.57	n.a.	2	2.00	0.49	0.17
UGPU	0.99	0.01	n.a.	2	1.12	0.02	-0.01
means (S.D.)				2.2 (.45)	1.90	0.38(0.21)	0.18

¹Allelic richness; ²rarefied allelic richness standardized to the sample size used for estimation of maternal allelic richness;

³Nei's expected heterozygosity (gene diversity); ⁴fixation index; asterisks indicate significance of associated G-test of genotypic frequencies (*=0.05, **=0.01, ***≤0.001); M = monomorphic.

Table 5-8. Estimates of allele frequencies, allelic richness, expected heterozygosity, fixation indices and G-test for Hardy-Weinberg disequilibria in populations of *Anacardium excelsum* from 12 forest fragments in northwestern Costa Rica (progeny data) (continued)

population / locus	Allele frequencies			A^1	λ_{w}^2	H_e^3	F_{is}^4
Las Congojas							
AK2	0.71	0.29	n.a.	2	2.00	0.41	0.44***
LAP	0.86	0.13	0.01	3	2.17	0.24	0.24***
PGD	0.92	0.08	n.a.	2	1.96	0.15	0.15*
PGM	0.68	0.32	n.a.	2	2.00	0.43	0.03
UGPU	0.85	0.15	n.a.	2	1.99	0.25	0.09
means (S.D.)				2.2 (.45)	2.02	0.30 (.12)	0.19
EJN							
AK2	0.72	0.28	n.a.	2	n.a.	0.40	-0.08
LAP	0.09	0.91	0.0	2	n.a.	0.16	-0.63*
PGD	0.97	0.03	n.a.	2	n.a.	0.06	-0.03
PGM	0.47	0.53	n.a.	2	n.a.	0.50	-0.42
UGPU	1.00	0.0	n.a.	1	n.a.	0.00	M
means (S.D.)				1.80(.45)	n.a.	0.22(0.22)	-0.29
Marcela							
AK2	0.94	0.06	n.a.	2	1.85	0.12	0.16
LAP	0.63	0.36	0.01	3	2.25	0.47	0.23**
PGD	0.91	0.09	n.a.	2	1.94	0.17	0.37***
PGM	0.61	0.39	n.a.	2	2.00	0.48	0.08
UGPU	0.89	0.11	n.a.	2	1.96	0.19	0.17
means (S.D.)				2.2 (.45)	2.00	0.28(0.18)	0.20
Palmira							
AK2	0.23	0.77	n.a.	2	1.88	0.35	0.39***
LAP	0.47	0.52	0.01	3	2.1	0.51	0.09
PGD	0.31	0.69	n.a.	2	1.95	0.43	0.41***
PGM	0.17	0.83	n.a.	2	1.75	0.29	0.08
UGPU	0.64	0.36	n.a.	2	1.97	0.46	0.48***
means (S.D.)				2.2 (.45)	1.93	0.41 (0.09)	0.29

¹Allelic richness; ²rarefied allelic richness standardized to the sample size used for estimation of maternal allelic richness;

³Nei's expected heterozygosity (gene diversity); ⁴fixation index; asterisks indicate significance of associated G-test of genotypic frequencies (*=0.05, **=0.01, ***≤0.001); M = monomorphic.

Table 5-8. Estimates of allele frequencies, allelic richness, expected heterozygosity, fixation indices and G-test for Hardy-Weinberg disequilibria in populations of *Anacardium excelsum* from 12 forest fragments in northwestern Costa Rica (progeny data) (continued)

population / locus	Allele frequencies			A^1	λ_m^3	H_e^3	F_{is}^4
	p_A	q_B	r_C				
Paso Hondo							
AK2	0.25	0.75	n.a.	2	1.91	0.38	0.43**
LAP	0.87	0.11	0.02	2	1.72	0.23	0.34*
PGD	0.71	0.29	n.a.	2	1.94	0.41	0.83***
PGM	0.72	0.28	n.a.	2	1.94	0.40	-0.08
UGPU	0.99	0.01	n.a.	1	1.07	0.02	-0.01
means (S.D.)				2.2 (.45)	1.72	0.29 (0.17)	0.30
River Santa Rosa							
AK2	0.81	0.19	n.a.	2	1.99	0.31	0.17*
LAP	0.74	0.24	0.02	3	2.34	0.40	0.37***
PGD	0.81	0.19	n.a.	2	1.99	0.31	0.22**
PGM	0.19	0.81	n.a.	2	1.99	0.31	0.03
UGPU	0.74	0.26	n.a.	2	2.00	0.38	0.42***
means (S.D.)				2.2 (.45)	2.06	0.34(0.04)	0.24
Toronja							
AK2	0.86	0.15	n.a.	2	1.88	0.24	0.07
LAP	0.38	0.51	0.11	3	2.77	0.58	0.57***
PGD	0.65	0.35	n.a.	2	2.00	0.46	0.03
PGM	0.64	0.36	n.a.	2	2.00	0.46	0.11
UGPU	0.96	0.04	n.a.	2	1.48	0.07	-0.04
means (S.D.)				2.2 (.45)	2.03	0.36 (.20)	

¹Allelic richness; ²rarefied allelic richness standardized to the sample size used for estimation of maternal allelic richness;

³Nei's expected heterozygosity (gene diversity); ⁴fixation index; asterisks indicate significance of associated G-test of genotypic frequencies (*=0.05, **=0.01, ***≤0.001); M = monomorphic.

Table 5-9. Homogeneity statistics for five loci of progeny genotypes in populations of *Anacardium excelsum* from 12 forest fragments in northwestern Costa Rica

locus	chi-square, df, probability	G-square, df, probability
<i>Overall (excludes EJN, Rodeo)</i>		
AK2	555.3, 2, <0.00001	561.3, 2, <0.00001
LAP	459.8, 3, <0.00001	456.7, 3, <0.00001
PGD	365.8, 3, <0.00001	327.6, 3, <0.00001
PGM	416.8, 3, <0.00001	435.8, 3, <0.00001
UGPU	161.4, 3, <0.00001	175.9, 3, <0.00001
<i>Corobici group</i>		
AK2	87.0, 3, <0.00001	123.2, 3, <0.00001
LAP	204.1, 3, <0.00001	175.6, 3, <0.00001
PGD	4.5, 3, <0.00001	4.9, 3, <0.00001
PGM	10.3, 3, <0.00001	10.9, 3, <0.00001
UGPU	2.87, 3, <0.00001	2.71, 3, <0.00001
<i>Taboga group</i>		
AK2	8.1, 2, 0.02	8.1, 2, 0.02
LAP	57.2, 2, <0.00001	58.2, 2, <0.00001
PGD	61.1, 2, <0.00001	65.9, 2, <0.00001
PGM	5.7, 2, 0.06	5.8, 2, 0.05
UGPU	9.2, 2, 0.01	9.4, 2, 0.01

Table 5-10. Estimates of Wright's statistics and mN for five loci of progeny genotypes in populations of *Anacardium excelsum* from 10 forest fragments in northwestern Costa Rica.

Locus	Sample size	F_s	F_H	F_{st}	mN
AK	2792	0.3905	0.5811	0.3127	0.55
LAP	2712	0.1989	0.3104	0.1392	1.55
PGD	2926	0.2610	0.3796	0.1604	1.31
PGM	2686	0.0459	0.2059	0.1676	1.24
UGPU	2934	0.2938	0.3633	0.0983	2.29
Mean	2810	0.2148	0.3563	0.1803	1.14
Corobicí group					
AK	1728	0.3690	0.4185	0.0785	2.9
LAP	1680	0.2019	0.3169	0.1442	1.5
PGD	1822	0.1919	0.2161	0.0300	8.1
PGM	1600	0.0378	0.1967	0.1652	1.3
UGPU	1836	0.2838	0.2949	0.0156	15.8
Mean	1733	0.2148	0.2847	0.0981	2.3
Taboga group					
AK	676	0.2284	0.2360	0.0099	25.1
LAP	636	0.2775	0.3578	0.1111	2.0
PGD	680	0.0690	0.1373	0.0734	3.2
PGM	684	0.0430	0.0531	0.0105	23.6
UGPU	672	0.1294	0.1401	0.0123	20.0
Mean	670	0.1438	0.1908	0.0549	4.3

Table 5-11. Pairwise estimates of Nei's unbiased genetic distance (below diagonal) and geographic distances (km, above diagonal) between ten populations of *Anacardium excelsum* located in northwestern Costa Rica

	Duquesa	Cepo	Ojoché	Isla	Congojas	Marcela	Paso Hondo	Palmira	R.S.R.	Toronja
Duquesa	0	17	17.25	7.9	16.75	2.75	9.6	24.75	15.25	4.5
Cepo	0.0771	0	2.92	9.25	1.85	18	11.1	8.1	3.1	19.3
Ojoché	0.0481	0.0946	0	11	2	21	13.75	6.35	3.6	21.5
Isla	0.1024	0.0881	0.0831	0	8.85	10.25	6.75	17.1	7.25	10.85
Congojas	0.0317	0.0471	0.0583	0.0923	0	19	11.85	8.25	1.7	19.4
Marcela	0.0257	0.0691	0.0082	0.0705	0.0313	0	12.5	27.3	17.25	5.8
Paso Hondo	0.3893	0.2561	0.4645	0.1839	0.3888	0.4204	0	19.3	11	9.6
Palmira	0.1295	0.1304	0.2342	0.1129	0.0802	0.1794	0.2522	0	10	27.8
R.S.R.	0.1072	0.0158	0.085	0.0854	0.0864	0.0702	0.2359	0.2165	0	18
Toronja	0.0384	0.1311	0.0361	0.0524	0.0935	0.0359	0.3173	0.1896	0.1249	0

Table 5-12. Estimates of current (1999) gene flow into the Paso Hondo population based on two alleles absent from the Paso Hondo population

Allele	q_i^1	q_i^2	m_i^3
UGPU-B	0.16	0.01	0.06
LAP-C	0.07	0.02	0.28
Mean	0.17		0.16

¹ Allele frequency in source population, i , overall mean allele frequency in Paso Hondo progeny, m_i =proportion of immigrant alleles

Table 5-13. Estimates of mating system parameters of *A. excelsum* in populations located in 10 forest fragments in northwestern Costa Rica

population	\hat{F}_1	t_m (s.e.) ²	t_i (s.e.) ³	$t_m - t_i$	r_i^4	r_p^5
Bosque Daguessa	0.40	0.430* (0.199)	0.647* (0.170)	-0.217* (0.044)	0.124* (0.060)	0.418* (0.121)
El Cepo	0.28	0.566* (0.078)	0.656* (0.068)	-0.090* (0.018)	0.220* (0.088)	0.878* (0.026)
El Ojoché	0.15	0.744* (0.106)	0.709* (0.093)	0.036 (0.049)	0.108 (0.080)	0.866* (0.055)
La Isla	0.43	0.396* (0.222)	0.656* (0.188)	-0.260* (0.058)	0.161* (0.063)	0.665* (0.068)
Las Congojas	0.15	0.739* (0.094)	0.811* (0.073)	-0.073* (0.028)	0.139* (0.045)	0.766* (0.064)
Marcela	0.19	0.684* (0.092)	0.726* (0.074)	-0.042 (0.034)	0.093* (0.023)	0.469* (0.116)
Palмира	0.58	0.266* (0.088)	0.258* (0.083)	0.008 (0.015)	0.230* (0.102)	0.868* (0.040)
Paso Hondo	0.26	0.581* (0.193)	0.665* (0.186)	-0.084 (0.042)	0.154 (0.307)	0.978* (0.087)
River Santa Rosa	0.40	0.427* (0.085)	0.487* (0.064)	-0.060* (.025)	0.101 (0.052)	0.480* (0.088)
Torroxja	0.22	0.638* (0.089)	0.668* (0.069)	-0.030 (0.027)	0.092 (0.111)	0.980* (0.012)

¹expected equilibrium value of F (see text); ²estimated multilocus outcrossing rate; * indicates significant estimate, $i.e.$ $\pm 2s.e. < 1.0$; ³average of estimated single-locus outcrossing rate; ⁴estimated correlation of outcrossing rates; * indicates significant estimate, $i.e.$ $r > 2s.e.$; ⁵estimated correlation of outcrossed paternity



Table 5-14. Mean population flowering indices and associated data in 24 forest fragments of *Anacardium excelsum* in northwestern Costa Rica.

Fragment	mean flowering index	mean dbh (\bar{x}) of sampled trees	Shannon's equitability (H') for flowering
<i>Fragments of type 1 (sheltered, on watercourses)</i>			
Canateca	133.4	112.4 (15.8)	0.81
El Cepo	137.5	136.4 (26.0)	0.85
El Ojoche	184.9	81.6 (16.4)	0.74
Finca La Gotera	53.2	113.4 (31.4)	0.77
Hcda. Tenorio	67.3	91.2 (18.9)	0.83
Las Congojas	30.4	84.7 (17.1)	0.44
Q. Reventado (south)	121.9	132.7 (18.9)	0.76
Quebrada El Lajero	94.9	107.3 (21.4)	0.80
River Santa Rosa	136.0	120.2 (38.8)	0.62
<i>Other fragments (type 0)</i>			
Bosque La Duquesa	18.0	104.2 (29.3)	0.41
COSTASEM	2.14	100.9 (25.8)	0.55
EJN	10.4	91.1 (29.2)	0.52
El Rodeo	211.2	114.2 (34.6)	0.61
Hacienda Los Ríos	54.9	96.1 (43.5)	0.48
La Isla	38.1	104.9 (18.4)	0.54
Las Avispas	1.9	97.8 (37.1)	0.65
Marcela	40.6	153.4 (33.9)	0.68
Oldemar	20.3	98.1 (23.3)	0.71
Paso Hondo	170.2	151.2 (19.6)	0.66
Q. Reventado (north)	41.6	104.9 (21.6)	0.67
Quebrada La Duquesa	33.8	124.0 (32.4)	0.17
Salitral-Libertad	13.1	93.4 (34.6)	0.41
Sitio Cascante	30.8	102.3 (23.3)	0.69
Toroja	66.6	101.9 (19.1)	0.62

Table 5-15. Linear regression of fragment mean tree flowering index on fragment type and mean dbh of sampled trees in 24 fragment populations of *A. excelsum* in northwestern Costa Rica.

Source of variation	degrees of freedom	Mean square	F	significance ¹ , R ²
Regression	2	14691.0	5.24	$p = .014, R^2 = 0.33$
Residual	21	2800.8		
Total	23			
	parameter estimates	std error	t	
intercept	-78.18	64.82	-1.21	0.24
type	56.77	22.31	2.54	0.018
mean dbh	1.18	0.58	2.03	0.056

Table 5-16. Linear regressions of flowering equitability on fragment type and mean dbh and equitability on type and dbh standard deviation of sampled trees in 24 fragment populations of *A. excelsum* in northwestern Costa Rica.

Source of variation	degrees of freedom	Mean square	F	significance ¹ , R ²
<i>Equitability v. type and mean dbh</i>				
Regression	2	0.098	5.02	p= .016, R ² =0.32
Residual	21	0.020		
Total	23			
	parameter estimates	std error	t	
intercept	0.39	0.17	2.29	0.032
type	0.18	0.06	3.02	0.006
mean dbh	0.002	0.001	0.99	0.330
<i>Equitability v. type and standard deviation of dbh</i>				
Regression	2	0.103	5.40	p= .013, R ² =0.34
Residual	21	0.019		
Total	23			
	parameter estimates	std error	t	
intercept	0.69	0.11	6.03	<0.0001
type	0.15	0.06	2.43	0.024
dbh s	-0.005	0.004	-1.23	0.232

Table 5-17. Analysis of variance of effect of origin (fragment) on weight of seeds collected from seven populations of *Anacardium excelsum* located in northwestern Costa Rica.

source of variation	degrees of freedom	mean square	F	p
Fragment	6	57603.14	35.56	p<0.0001
Residual	378	1619.79		
Total	384			

Table 5-18. Means, standard errors and Bonferroni groupings of weight of seeds collected from seven populations of *Anacardium excelsum* located in northwestern Costa Rica.

Fragment	Mean weight (g), Bonferroni grouping ¹	Standard error
El Cepo	A 2.59	0.0546
Canateca	A 2.49	0.0641
Marcela	B 2.19	0.0480
Quebrada Reventado	CB 2.10	0.0549
Toronja	CD 1.92	0.0412
Salitral	CD 1.87	0.0579
Duquesa	D 1.72	0.0561

¹Same letter indicates means not significantly different, $\alpha=0.05$

Table 5-19. Analysis of variance of effect of origin (fragment) on arcsin germination

Table 5-20. Germination percentage, least square means for height at four months (cm) and number of days to germination for *Anacardium excelsum* seedlings from seven forest fragments in northwestern Costa Rica

Fragment	Germination percentage ¹	Fragment	Height, 4 months (cm) (S.E.)	Fragment	germination day (S.E.)	Fragment	Height (with covariates), 4 months (cm) S.E.
Marcela	A	Marcela	31.5 (1.05)	Reventado	A	Marcela	A
El Cepo	B	El Cepo	29.4 (1.28)	Marcela	A	El Cepo	A
Canateca	B	Canateca	25.6 (1.21)	Toronja	A	Canateca	B
Duquesa	B	Salitral	25.0 (2.39)	El Cepo	A	Duquesa	B
Toronja	B	Reventado	23.4 (1.82)	Canateca	A	Salitral	A
Reventado	B	Duquesa	23.0 (1.46)	Salitral	A	Reventado	B
Salitral	B	Toronja	22.1 (1.90)	Duquesa	A	Toronja	B

¹Data presented are overall, untransformed experimental means. Bonferroni grouping based on ANOVA of arcsin transformed data (see Table 5-19)

Table 5-21. Analysis of covariance of effect of origin (fragment), seed weight and germination day on four-month of seedlings from seven populations of *Anacardium excelsum* located in northwestern Costa Rica.

trait / source	degrees of freedom	mean squares (type 3)	F	p
Height, four months				
Block	4	57.6	1.7	0.15
Fragment	6	172.4	5.2	<0.0001
germination day	1	755.9	22.6	<0.0001
seed weight	1	183.0	5.48	0.02
error	119	33.4		

Table 5-22. Analysis of variance of a greenhouse experiment comparing growth of *Anacardium excelsum* seedlings from 9 forest fragments in northwestern Costa Rica

trait / source	degrees of freedom	expected mean squares	mean squares (type 3)	F	p
Height, 81 days					
Block	9	$\sigma^2_{\text{error}} + 43.222\sigma^2_{\text{block}}$	35.78	4.12	<0.0001
Provenance	8	$\sigma^2_{\text{error}} + 5.9104\sigma^2_{\text{fam}(p)}$	46.94	5.41	<0.0001
Family-in-provenance	58	$\sigma^2_{\text{error}} + 6.6671\sigma^2_{\text{fam}(p)}$	22.51	2.59	<0.0001
Error	380		8.68		
Height, 167 days					
Block	9	$\sigma^2_{\text{error}} + 43.333\sigma^2_{\text{block}}$	43.7	2.1	0.03
Provenance	8	$\sigma^2_{\text{error}} + 5.958\sigma^2_{\text{fam}(pr)}$	114.9	5.52	<0.0001
Family-in-provenance	58	$\sigma^2_{\text{error}} + 5.6803\sigma^2_{\text{fam}(p)}$	55.7	2.68	0.0009
Error	381		20.8		
Diameter, 167 days					
Block	9	$\sigma^2_{\text{error}} + 43.333\sigma^2_{\text{block}}$	4.7	2.7	0.005
Provenance	8	$\sigma^2_{\text{error}} + 5.959\sigma^2_{\text{fam}(pr)}$	5.8	3.34	0.001
Family-in-provenance	58	$\sigma^2_{\text{error}} + 5.6803\sigma^2_{\text{fam}(p)}$	3.90	2.26	<0.0001
Error	381		1.73		
Testa length					
Block	9	$\sigma^2_{\text{error}} + 56.222\sigma^2_{\text{block}}$	9.6	1.0	0.44
Fragment	8	$\sigma^2_{\text{error}} + 8.225\sigma^2_{\text{fam}(pr)}$	43.8	4.6	<0.0001
Family(fragment)	56	$\sigma^2_{\text{error}} + 8.7419\sigma^2_{\text{fam}(p)}$	30.2	3.2	<0.0001
Error	497		9.6		

Table 5-23. Least square means of total height at 81 and 167 days and diameter 2cm above root collar at 167 days of *Anacardium excelsum* seedlings from 9 forest fragments in northwestern Costa Rica

Fragment	Height, 81 days (cm)	Fragment	Diameter, 167 days (mm)	Fragment	Height at 167 days (cm)	Fragment	Testa length (mm) (S.E.)
Marcela	A 18.3 (0.56)	Marcela	A 7.5 (0.25)	Marcela	A 25.0 (0.87)	R.S.R.	A 32.4 (0.43)
Congojas	B A 17.6 (0.37)	Congojas	B 7.1 (0.17)	Congojas	B A 23.6 (0.58)	Congojas	A 32.3 (0.31)
La Isla	C B A 17.3 (0.60)	El Ojoché	B 7.0 (0.26)	Isla	C B A 22.4 (0.90)	Isla	B A 32.0 (0.50)
Palmita	C B A 16.6 (0.49)	Isla	B 7.0 (0.26)	Palmita	C B A 22.3 (0.76)	El Cepo	B A 31.8 (0.24)
El Cepo	C 15.8 (0.30)	Palmita	B 6.9 (0.22)	Toronja	C 20.8 (0.93)	Ojoché	B A 31.7 (0.51)
El Ojoché	C B 15.7 (0.55)	Toronja	B 6.6 (0.27)	El Cepo	C 20.8 (0.46)	Marcela	B A 31.4 (0.43)
Toronja	C B 15.7 (0.60)	El Cepo	B 6.5 (0.13)	R.S.R.	C 20.4 (0.80)	Palmita	B A 31.2 (0.51)
R.S.R.	C 15.5 (0.52)	B. Duguesa	B 6.5 (0.28)	Ojoché	C 20.2 (0.86)	Toronja	B 29.8 (0.53)
B. Duguesa	C 14.6 (0.62)	R.S.R.	B 6.2(0.23)	B. Duguesa	C 19.2 (0.97)	B. Duguesa	B 29.5 (0.55)

Table 5-24. Analysis of covariance of a greenhouse experiment comparing growth of *Anacardium excelsum* seedlings from 9 forest fragments in northwestern Costa Rica

trait / source	degrees of freedom	expected mean squares	mean squares (type 3)	F	p
Height, 81 days					
Block	9	$\sigma^2_{\text{error}} + 39.385\sigma^2_{\text{block}}$	22.2	3.51	<0.0004
Provenance	8	$\sigma^2_{\text{error}} + 5.4588\sigma^2_{\text{fam}(p)}$	36.3	5.76	<0.0001
Family-in-provenance	56	$\sigma^2_{\text{error}} + 6.133\sigma^2_{\text{fam}(p)}$	15.9	2.52	<0.0001
testa length	1	$\sigma^2_{\text{error}} + Q_{\text{length}}$	475.7	75.4	<0.0001
testa shedding date	1	$\sigma^2_{\text{error}} + Q_{\text{shed}}$	615.4	97.5	<0.0001
Error	346		6.31		
Height, 167 days					
Block	9	$\sigma^2_{\text{error}} + 39.945\sigma^2_{\text{block}}$	17.7	1.17	0.3129
Provenance	8	$\sigma^2_{\text{error}} + 5.5001\sigma^2_{\text{fam}(p)}$	84.4	5.6	<0.0001
Family-in-provenance	56	$\sigma^2_{\text{error}} + 6.3255\sigma^2_{\text{fam}(p)}$	38.2	2.53	<0.0001
testa length	1	$\sigma^2_{\text{error}} + Q_{\text{length}}$	1109.0	73.5	<0.0001
testa shedding date	1	$\sigma^2_{\text{error}} + Q_{\text{shed}}$	1562.5	103.5	<0.0001
Error	347		15.1		
Diameter, 167 days					
Block	9	$\sigma^2_{\text{error}} + 39.945\sigma^2_{\text{block}}$	2.1	1.59	0.12
Provenance	8	$\sigma^2_{\text{error}} + 5.5001\sigma^2_{\text{fam}(p)}$	4.4	3.33	0.0011
Family-in-provenance	56	$\sigma^2_{\text{error}} + 6.3255\sigma^2_{\text{fam}(p)}$	2.54	1.92	0.0002
testa length	1	$\sigma^2_{\text{error}} + Q_{\text{length}}$	34.1	25.9	<0.0001
testa shedding date	1	$\sigma^2_{\text{error}} + Q_{\text{shed}}$	168.4	127.5	<0.0001
Error	381		1.73		

Table 5-25. Least square means of total height at 81 and 167 days and diameter 2cm above root collar at 167 days of *Anacardium excelsum* seedlings from 9 forest fragments in northwestern Costa Rica (based on analysis of covariance)

Fragment	Height, 81 days (cm)	Fragment	Diameter, 167 days (mm)	Fragment	Height at 167 days (cm)
Congojas	A 17.4 (0.29)	El Ojoche	A 7.1 (0.23)	Marcela	A 23.4 (0.77)
Marcela	A 17.3 (0.50)	Congojas	A 7.0 (0.13)	Congojas	A 23.1 (0.45)
La Isla	A 16.9 (0.54)	Marcela	A 7.0 (0.23)	El Cepo	A 22.0 (0.38)
El Cepo	A 16.6 (0.25)	Isla	B A 6.9 (0.24)	Palmira	C A 22.0 (0.69)
Palmira	A 16.5 (0.45)	Palmira	B A 6.8 (0.20)	Isla	C B A 21.7 (0.82)
El Ojoche	B A 16.1 (0.50)	El Cepo	A 6.8 (0.11)	Ojoche	C B A 20.8 (0.77)
Toronja	B A 15.7 (0.60)	Toronja	B A 6.4 (0.24)	Toronja	C B 20.8 (0.81)
R.S.R.	B 14.8 (0.52)	B. Duquesa	B A 6.1 (0.26)	R.S.R.	C B 19.1 (0.80)
B.	B 14.0 (0.57)	R.S.R.	B 6.0(0.24)	B. Duquesa	B 18.2 (0.88)
Duquesa					

Table 5-26. Numbers of albino and normal seedlings in progeny of three trees of *Anacardium excelsum* from populations located in forest fragments in northwestern Costa Rica

Family	Number of seed sown	Number of albino germinants	Number of normal germinants
1505	75	0	19
1541	45	1	24
Ojoche 21	70	13	31

Table 6-1. Segregation ratios in pooled progeny arrays of five loci of *Plumeria rubra*.

Enzyme	Maternal genotype	Classes for G-test	Observed numbers per class	n_{total}	G_{adj}^1
<i>AAT1</i>	AB	AA+BB	231	441	0.99
		AB	210		
	AB	AC	9	21	0.42
		BC	12		
	AC	AA+CC	76	138	1.42
		AC	62		
	AC	AB	15	28	0.14
		BC	13		
	BC	BB+CC	3	7	0.13
		BC	4		
BC	AB	15	33	0.27	
	BC	18			
<i>ADH3</i>	AB	AA+BB	774	1546	0.003
		AB	772		
<i>PGI3</i>	AC	AA+CC	78	140	1.83
		AC	62		
<i>PGM1</i>	AB	AA+BB	43	86	0.00
		AB	43		
	AC	AA+CC	56	115	0.08
		AC	59		
	AB	AC	6	9	0.97
		BC	3		
	AC	AC	2	2	n.a.
BC		0			
<i>PGD1</i>	AB	AA+BB	834	0.76	0.76
		AB	870		

¹Critical value of G for $\beta=0.05$ is 3.841, with Williams's adjustment (Sokal and Rohlf, 1995)

Table 6-2. Segregation ratios and associated probabilities in progeny of mother trees of *Plumeria rubra* putatively heterozygous for the AAT1 locus.

Population, maternal genotypes	Family	Observed numbers per class				n_{total}	n_{exp}	p^1
AB mothers		n_{AA}	n_{BB}	$n_{AA}+n_{BB}$	n_{AB}			
Corobici	1001	14	1	15	15	30	15	1.00
Corobici	1017	15	10	25	28	53	26.5	0.78
Corobici	1018	2	8	10	10	20	10	1.00
La Pachanga	8	2	1	3	2	5	2.5	1.00
Magdalena	4003	13	6	19	22	41	20.5	0.76
Magdalena	4009	16	9	25	19	44	22	0.45
Magdalena	4010	17	4	21	19	40	20	0.87
Magdalena	4014	17	4	21	10	31	15.5	0.07
Magdalena	4016	22	6	28	30	58	29	0.90
Palmira	1-300	11	2	13	9	22	11	0.52
Palmira	1-307	5	4	9	14	23	11.5	0.40
Palmira	7	8	5	13	7	20	10	0.26
San Isidro	2021	1	8	9	7	16	8	0.80
San Isidro	2027	19	1	20	18	38	19	0.87
AC mothers		n_{AA}	n_{CC}	$n_{AA}+n_{CC}$	n_{AC}			
Palmira	1-200	16	5	21	12	33	16.5	0.16
Praderas	2048	2	21	23	22	45	22.5	1.00
Praderas	2049	3	7	10	7	17	8.5	0.63
Praderas	2040	9	2	11	10	21	10.5	1.00
Praderas	2047	1	10	11	11	22	11	1.00
AB mothers		n_{AC}	n_{BC}					
Corobici	1018	3	5			8	4	0.73
La Pachanga	8	0	1			1	0.5	1.00
Magdalena	4009	3	0			3	1.5	0.25
Magdalena	4014	2	2			4	2	1.00
San Isidro	2027	1	4			5	2.5	0.38
AC mothers		n_{AB}	n_{CB}					
Palmira	1-200	13	10			23	11.5	0.68
Praderas	2040	2	1			3	1.5	1.00
Praderas	2047	0	2			2	1.0	0.5

¹two-tailed exact binomial cumulative probability under the null hypothesis, of observed numbers of progeny in either class

Table 6-3. Segregation ratios and associated probabilities in progeny of mother trees of *Plumeria rubra* putatively heterozygous for ADH³ alleles A and B.

Population	Family	Observed numbers per class				n_{total}	n_{exp}	p^1
		n_{AA}	n_{BB}	$n_{AA}+n_{BB}$	n_{AB}			
Corobici	1000	1	17	18	21	39	19.5	0.75
Corobici	1003	15	14	29	19	48	24	0.19
Corobici	1005	13	12	25	20	45	22.5	0.55
Corobici	1006	12	17	29	41	70	35	0.19
Corobici	1007	17	10	27	13	40	20	0.04
Corobici	1009	7	11	18	25	43	21.5	0.36
Corobici	1010	7	11	18	16	34	17	0.86
Corobici	1011	9	16	25	23	48	24	0.88
Corobici	1012	5	22	27	25	52	26	0.89
Corobici	1017	17	16	33	30	63	31.5	0.80
La Pachanga	C4	6	13	19	13	32	16	0.38
La Pachanga	C7	3	8	11	10	21	10.5	1.00
La Pachanga	5	2	5	7	1	8	4	0.07
La Pachanga	6	7	9	16	12	28	14	0.57
La Pachanga	7	3	4	7	17	24	12	0.06
La Pachanga	3	1	4	5	6	11	5.5	1.00
Magdalena	4000	8	17	25	16	41	20.5	0.21
Magdalena	4005	3	6	9	9	18	9	1.00
Magdalena	4007	7	13	20	29	49	24.5	0.25
Magdalena	4008	3	20	23	22	45	22.5	1.0
Magdalena	4009	8	17	25	28	53	26.5	0.78
Magdalena	4011	2	26	28	40	68	34	0.18
Magdalena	4014	2	15	17	18	35	17.5	1.00
Magdalena	4018	2	8	10	7	17	8.5	0.63
Palmira	1-300	6	1	7	12	19	9.5	0.36
Palmira	1-306	11	1	12	12	24	12	1.00
Palmira	1-307	4	7	11	11	22	11	1.00
Praderas	2037	12	6	18	22	40	20	0.64
Praderas	2040	2	9	11	13	24	12	0.84
Praderas	2042	8	17	25	34	59	29.5	0.30
Praderas	2043	5	8	13	7	20	10	0.26
Praderas	2047	10	3	13	12	25	12.5	1.00
Praderas	2048	7	21	28	17	45	22.5	0.14
Praderas	2054	3	1	4	5	9	4.5	1.00
Praderas	2055	15	13	28	25	53	26.5	0.78

¹two-tailed exact binomial cumulative probability under the null hypothesis, of observed numbers of progeny in either class

Table 6-3. Segregation ratios and associated probabilities in progeny of mother trees of *Plumeria rubra* putatively heterozygous for ADH3 alleles A and B (continued)

Population	Family	Observed numbers per class				n_{total}	n_{exp}	p^1
		n_{AA}	n_{BB}	$n_{AA+n_{BB}}$	n_{AB}			
San Isidro	2010	6	12	18	24	42	21	0.44
San Isidro	2011	15	10	25	22	47	23.5	0.77
San Isidro	2017	4	5	9	9	18	9	1.00
San Isidro	2018	11	6	17	22	39	19.5	0.52
San Isidro	2022	9	4	13	16	29	14.5	0.71
San Isidro	2026	8	9	17	15	32	16	0.86
San Isidro	2030	7	6	13	7	20	10	0.26
San Isidro	2034	10	9	19	28	47	23.5	0.24

¹two-tailed exact binomial cumulative probability under the null hypothesis, of observed numbers of progeny in either class.

Table 6-4. Segregation ratios and associated probabilities in progeny of mother trees of *Plumeria rubra* putatively heterozygous for locus PGI3.

Population, maternal genotypes	Family	Observed numbers per class				n_{total}	n_{exp}	p^1
		n_{AA}	n_{CC}	$n_{AA+n_{CC}}$	n_{AC}			
AC mothers								
Corobici	1018	13	3	16	10	26	13	0.33
La Pachanga	8	2	1	3	1	4	2	0.63
Magdalena	4016	21	1	22	24	46	23	0.88
Magdalena	4017	15	1	16	9	25	12.5	0.23
Quebrada Palmira	1-307	1	9	10	8	18	9	0.81
Quebrada Palmira	qp12	9	2	11	10	21	10.5	1.00

¹two-tailed exact binomial cumulative probability under the null hypothesis, of observed numbers of progeny in either class

Table 6-5. Segregation ratios and associated probabilities in progeny of mother trees of *Plumeria rubra* putatively heterozygous for the PGM1 locus

Population, maternal genotypes	Family	Observed numbers per class				n_{total}	n_{exp}	p^1
		n_{AA}	n_{BB}	$n_{AA}+n_{BB}$	n_{AB}			
AB mothers								
Corobici	1004	25	1	26	28	54	27	0.66
San Isidro	2022	10	6	16	15	31	15.5	1.00
AC mothers								
		n_{AA}	n_{CC}	$n_{AA}+n_{CC}$	n_{AC}			
Corobici	1006	23	2	25	30	55	27.5	0.59
	1010	22	1	23	22	45	22.5	1.00
Q. Palmira	1	5	1	6	6	12	6	1.00
AB mothers								
		n_{AC}	n_{BC}					
1004		6	2			8	4	0.29
AC mothers								
		n_{AB}	n_{BC}					
Q. Palmira	1	2	0			2	1	0.50

¹two-tailed exact binomial cumulative probability under the null hypothesis, of observed numbers of progeny in either class

Table 6-6. Segregation ratios and associated probabilities in progeny of mother trees of *Plumeria rubra* putatively heterozygous for locus PGD1.

Population	Family	Observed numbers per class				n_{total}	n_{exp}	p^1
		n_{AA}	n_{BB}	$n_{AA}+n_{BB}$	n_{AB}			
Qbrda. Palmira	100	31	1	32	17	49	24.5	0.04
	300	2	2	4	12	16	8	0.08
	309	7	1	8	11	19	9.5	0.65
	10	5	2	7	4	11	5.5	0.55
	12	5	4	9	11	20	10	0.82
	2	6	1	7	16	23	11.5	0.09
	3	3	2	5	5	10	5	1.00
	7	2	6	8	9	17	8.5	1.00
	9	2	5	7	5	12	6	0.77
Corobicí	1000	12	5	17	16	33	16.5	1.00
	1001	14	1	15	31	46	23	0.03
	1004	24	3	27	21	48	24	0.47
	1005	11	7	18	20	38	19	0.87
	1006	13	13	26	33	59	29.5	0.43
	1008	24	5	29	23	52	26	0.49
	1010	11	7	18	20	38	19	0.87
	1011	6	19	25	15	40	20	0.15
	1012	3	11	14	21	35	17.5	0.31
	1013	8	6	14	11	25	12.5	0.69
	1017	22	6	28	31	59	29.5	0.79
	1018	6	14	20	17	37	18.5	0.74
	1021	14	6	20	28	48	24	0.31
San Isidro	2010	2	17	19	14	33	16.5	0.49
	2011	13	9	22	20	42	21	0.88
	2013	1	2	3	6	9	4.5	0.51
	2018	13	6	19	7	26	13	0.03
	2021	5	4	7	11	18	9	0.48
	2021	5	2	7	11	18	9	0.48
	2022	12	0	12	14	26	13	0.85
	2027	11	3	14	29	43	21.5	0.03
	2034	0	25	25	13	38	19	0.07
	2041	22	5	27	31	58	29	0.69
	2047	8	6	14	8	22	11	0.29
2051	11	5	16	20	36	18	0.62	

¹two-tailed exact binomial cumulative probability under the null hypothesis, of observed numbers of progeny in either class

Table 6-6. Segregation ratios and associated probabilities in progeny of mother trees of *Plumeria rubra* putatively heterozygous for locus PGD (continued)

Population	Family	Observed number per class				n_{total}	n_{exp}	p^1
		n_{AA}	n_{BB}	n_{AA+BB}	n_{AB}			
Magdalena	4000	7	2	9	10	19	9.5	1.00
	4003	11	6	17	17	34	17	1.00
	4007	15	4	19	24	43	21.5	0.54
	4008	18	2	20	25	45	22.5	0.55
	4009	12	10	22	29	51	25.5	0.40
	4010	19	2	21	15	36	18	0.41
	4013	21	6	27	19	46	23	0.30
	4016	6	8	14	16	30	15	0.86
	4019	16	4	20	24	44	22	0.65
	4020	12	4	16	20	36	18	0.62
Pachanga	CS7	5	9	14	7	21	10.5	0.19
	P1	10	0	10	10	20	10	1.00
	P2	7	7	14	9	23	11.5	0.40
	P5	3	2	5	4	9	4.5	1.00
	P6	5	2	7	21	28	14	0.01
	p7	3	10	13	7	20	10	0.26
Sandillal	r00	14	7	21	25	46	23	0.66
	r02	19	3	22	27	49	24.5	0.57

¹two-tailed exact binomial cumulative probability under the null hypothesis, of observed numbers of progeny in either class

Table 6-7. Significant estimates of Burrows's composite linkage disequilibrium coefficient, correlation coefficient and significance of associated chi-squared test in 7 populations of *Plumieria rubra* in Guanacaste province, Costa Rica.

Locus / alleles ¹	Number ² +/-	Population	Δ_i	r	P	n
AAT-A/PGM-A	5/2	Corobici	-0.0113	0.0171	.0165	720
ADH-A/PGI-A	3/4	Palмира	-0.0213	-0.1139	0.0391	328
ADH-A/PGM-A	3/4	San Isidro	0.0233	0.124	0.007	460

¹All alleles except the most frequent were pooled to one synthetic allele; i.e. number of positive and negative correlations, irrespective of significance, over all seven populations.

Table 6-8. Estimates of Ohta's multiple population linkage disequilibrium coefficients for 7 populations of *Plumeria rubra* in northwestern Costa Rica.

Loci	D_{IT}^2	D_{IS}^2	D'_{IS}^2	D_{ST}^2	D'_{ST}^2
AAT/ADH	.06148	.00036	.059083	.06132	.00166
AAT/PGI	.02186	.00011	.02183	.02205	.00002
AAT/PGM	.02695	.00065	.02693	.02478	.00001
AAT/PGD	.02869	.00050	.02855	.02644	.00014
ADH/PGI	.04148	.00025	.04145	.04304	.00003
ADH/PGM	.04945	.00052	.04943	.04559	.00002
ADH/PGD	.04005	.00067	.03953	.03879	.00052
PGI/PGM	.00802	.00001	.00800	.00813	.00003
PGI/PGD	.02059	.00009	.02042	.02039	.00017
PGM/PGD	.02215	.00026	.02202	.02091	.00013
Overall	.03207	.00034	.03180	.03114	.00027

Table 7-1. Description of populations sampled in a study of the genetic effects of forest fragmentation on *Plumeria rubra* in northwestern Costa Rica.

Population	Coordinates, population centre	Population size	isolation ¹	disturbance ²	matrix	ndi ³ (\bar{x} , CV%)	dbh ⁴ (\bar{x} , CV%)	Comments
Quebrada Palmira	85°05.7' 10°32.3'	<30	L	L-M	pastureland	0.95,127	29.6,39	Includes two, probably planted, outlying trees in Palmira cemetery and neighbouring pasture
River Corobici	85°05.9' 10°32.7'	22	L	L-M	pastureland	2.15,82	32.2,48	
San Isidro	85°08.5' 10°30.0'	35 (300+)	H	L (M-H)	pastureland, open forest	7.76,44	29.2,35	upstream of sampled area, there is open forest with individual <i>P. rubra</i> trees until a large concentration of <i>P. rubra</i> located near Panales (Figure 1)
Praderas Norte	del 85°08.6' 10°31.9'	30-50	H	L (M)	plateau pastureland, open forest	2.25	22 ⁶	
Magdalena	85°06.6' 10°28.5'	21	H	H (M)	mixed agriculture, pastureland, rural dwellings	2.31,72%	21.9,54	

¹L=low, M=moderate, H=high; ²L=low, M=moderate, H=high, based on tree cover of fragment where population is located; parentheses refer to localized atypical sectors; ³neighbourhood density index, see text; ⁴diameter at breast height; ⁵terrestrial; ⁶estimate based on mean fruit production tree⁻¹

Table 7-1. Description of populations sampled in a study of the genetic effects of forest fragmentation on *Plumeria rubra* in northwestern Costa Rica (continued)

Population	Coordinates, population centre	Population size	isolation ²	disturbance ³	matrix	ndi ⁴ (\bar{x} , CV%)	dbh ⁵ (\bar{x} , CV%)	Comments
La Pochanga	85°06.4' 10°29.3'	20-50	L-M	L-M	mixed agriculture, 0.71, 55% pastureland, secondary forest	30.9, 46		
Sandlial	85°06.1' 10°27.8'	4	H	M-H	pastureland, semi-urban, hydroelectric plant reservoir and	0.50, 41%	39.2, 20	

¹L=low, M=moderate, H=high; ²L=low, M=moderate, H=high, based on tree cover of fragment where population is located, parentheses refer to localized atypical sectors; ³neighbourhood density index, see text; ⁴diameter at breast height; ⁵stermate; ⁶stermate based on mean fruit production tree⁻¹

Table 7-2. Sampling schedule in a study of genetic effects of forest fragmentation in *Plumeria rubra* in northwestern Costa Rica (continued)

Fragment, sampling descriptors	Collection			
	1997	1998	1999	Pooled
Sandillal				
Number of families assayed	0	3	4	4
Number of progeny assayed	0	127	86	213
Mean number progeny family ⁻¹	0	42.3	21.5	53.2
Mean number of capsules family ⁻¹	0	4.7	5.2	8.8
Mean number progeny capsule ⁻¹	0	9.2	4.2	6.0
All				
Number of families assayed	50	69	37	94
Number of progeny assayed	1141	1405	726	3272
Mean number progeny family ⁻¹	22.8	20.4	19.6	34.3
Mean number of capsules family ⁻¹	3.4	3.4	4.0	5.8
Mean number progeny capsule ⁻¹	7.6	6.2	5.4	6.2

Table 7-3. Estimates of allele frequencies, number of alleles, expected heterozygosity, fixation indices and G-test for Hardy-Weinberg disequilibria in populations of *P. rubra* from 7 forest fragments in northwestern Costa Rica, based on inferred maternal genotypes.

Population / Locus	Allele frequencies			A	$H_e(N_{el})$	F_{is}^1
	p	q	r			
Quebrada Palmira						
AAT	0.69	0.23	0.07	3	0.48	-0.08
ADH	0.54	0.46	n.a.	2	0.50	-0.01
PGI	0.82	0.18	n.a.	2	0.29	-0.22
PGM	0.95	0.035	0.035	3	0.14	-0.08
PGD	0.50	0.50	n.a.	2	0.50	-0.38
means				2.40(.55)	0.38(0.16)	
River Corobicí						
AAT	0.79	0.15	0.06	3	0.34	0.10
ADH	0.35	0.65	n.a.	2	0.46	-0.29
PGI	0.97	0.03	n.a.	2	0.06	-0.03
PGM	0.88	0.03	0.08	3	0.22	-0.13
PGD	0.56	0.44	n.a.	2	0.49	-0.55*
means				2.4 (.55)	0.31 (0.18)	
San Isidro						
AAT	0.71	0.18	0.12	3	0.46	-0.13
ADH	0.50	0.50	n.a.	2	0.50	-0.11
PGI	1.00	0.0	n.a.	1	0.00	M
PGM	0.78	0.19	0.03	3	0.36	0.04
PGD	0.61	0.39	n.a.	2	0.48	-0.17
means				2.20 (.84)	0.36 (0.21)	
Praderas						
AAT	0.69	0.06	0.25	3	0.46	-0.08
ADH	0.44	0.56	n.a.	2	0.49	-0.42*
PGI	1.00	0.0	n.a.	1	0.00	M
PGM	0.88	0.12	0.0	2	0.22	-0.13
PGD	0.75	0.25	n.a.	2	0.38	0.00
means				2.0 (0.71)	0.31(0.20)	
Magdalena						
AAT	0.79	0.18	0.03	3	0.34	-0.26
ADH	0.26	0.74	n.a.	2	0.39	-0.36
PGI	0.91	0.08	n.a.	2	0.16	-0.10
PGM	0.88	0.12	0.0	2	0.21	-0.13
PGD	0.65	0.35	n.a.	2	0.46	-0.29
Mean				2.20 (0.45)	0.31(.12)	

¹Asterisks indicate significance of G-test for Hardy-Weinberg disequilibrium: ***p<.001, **<.01, *>.05; M=monomorphic

Table 7-3. Estimates of allele frequencies, number of alleles, expected heterozygosity, fixation indices and G-test for Hardy-Weinberg disequilibria in populations of *P. rubra* from 7 forest fragments in northwestern Costa Rica, based on inferred maternal genotypes (continued).

Population / Locus	Allele frequencies			A	$H_{e(Nel)}$	F_{is}
Pachanga						
AAT	0.75	0.19	0.06	3	0.40	0.33
ADH	0.50	0.50	n.a.	2	0.50	-0.50
PGI	0.94	0.06	n.a.	2	0.12	-0.07
PGM	1.0	0.0	0.0	1	0.00	M
PGD	0.50	0.50	n.a.	2	0.50	-0.50
Mean				2.0 (0.71)	0.30(0.23)	
Sandillal						
AAT	1.0	0.0	0.0	1	0.00	M
ADH	0.0	1.0	n.a.	1	0.00	M
PGI	0.88	0.12	n.a.	2	0.22	-0.14
PGM	1.0	0.0	0.0	1	0.00	M
PGD	0.75	0.25	N.A	2	0.38	-0.33
Mean				1.4 (0.55)	0.12(0.17)	

¹Asterisks indicate significance of G-test for Hardy-Weinberg disequilibrium: ***p<.001, **<.01, *>.05; M=monomorphic

Table 7-4. Homogeneity statistics for five loci of inferred maternal genotypes seven populations in a study of effects of forest fragmentation on genetics of *Plumeria rubra* in northwestern Costa Rica

Locus	chi-square, df, probability	G-square, df, probability
AAT	4.9, 6, 0.56	6.8, 6, 0.34
ADH	12.7, 6, 0.05	15.6, 6, 0.02
PGI	13.2, 6, 0.04	14.7, 6, 0.02
PGM	7.6, 6, 0.27	9.8, 6, 0.13
PGD	6.0, 6, 0.42	6.14, 6, 0.441

Table 7-5. Estimates of Wright's statistics and mN between seven populations (inferred maternal genotypes) (estimates in parentheses based on data set without Pachanga and Sandliall populations)

Locus	Sample size	F_{is}	Fit	F_{st}	mN
AAT	184 (160)	-0.0951 (-0.1248)	-0.0322 (-0.0946)	0.0575 (0.0269)	4.1 (9.0)
ADH	188 (164)	-0.2954 (-0.2516)	-0.1254 (-0.2015)	0.1312 (0.0400)	1.6 (6.0)
PGI	188 (164)	-0.1418 (-0.1586)	-0.0742 (-0.0630)	0.0592 (0.0825)	4.0 (2.8)
PGM	188 (164)	-0.1074 (-0.1074)	-0.0371 (-0.0728)	0.0635 (0.0312)	3.7 (7.8)
PGD	186 (162)	-0.3309 (-0.2938)	-0.2770 (-0.2549)	0.0404 (0.0312)	5.9 (8.1)
Mean	187 (163)	-0.2225 (-0.2066)	-0.1313 (-0.1638)	0.0747 (0.0354)	3.1 (6.8)

Table 7-6. Pairwise estimates of Nei's unbiased genetic distance (below diagonal) and geographic distances (km, above diagonal) between seven populations of *Plumeria rubra* located in Guanacaste province, northwestern Costa Rica

	Q. Palmira	R. Corobicí	San Isidro	Praderas	Magdalena	Pachanga	Sandillal
Q. Palmira		0.99	6.574	4.119	7.464	5.914	8.733
R. Corobicí	0.011		6.472	3.423	8.072	5.727	8.66
San Isidro	0.013	0.008		3.593	6.472	3.959	6.454
Praderas	0.033	0.017	0.006		8.429	5.646	8.745
Magdalena	0.026	0.001	0.017	0.018		1.85	1.304
Pachanga	-0.01	0.001	0.007	0.023	0.017		3.129
Sandillal	0.097	0.041	0.088	0.065	0.019	0.079	

Table 7-7. Estimates of allele frequencies, number of alleles, expected heterozygosity, fixation indices and G-test for Hardy-Weinberg disequilibria in 1997 progeny of four populations of *P. rubra* in northwestern Costa Rica

Population / Locus	Allele frequencies			A	H_e (Nei)	Fis ¹
	p	q	r			
Río Corobicí						
AAT	0.77	0.17	0.06	3	0.37	0.04
ADH	0.40	0.60	n.a.	2	0.48	0.05
PGI	0.96	0.04	n.a.	2	0.08	0.15*
PGM1	0.88	0.04	0.07	3	0.21	-0.04
PGD	0.61	0.39		2	0.48	-0.05
Means (s.d.)				2.4 (0.55)	0.37 (0.20)	
San Isidro						
AAT	0.76	0.18	0.06	3	0.39	-0.06
ADH	0.51	0.49	n.a.	2	0.50	-0.04
PGI	0.96	0.04	n.a.	2	0.08	-0.05
PGM1	0.79	0.17	0.04	3	0.35	-0.03
PGD	0.54	0.46	n.a.	2	0.50	0.03
Means (s.d.)				2.4 (0.55)	0.36 (0.15)	
Praderas						
AAT	0.77	0.06	0.17	3	0.38	-0.07
ADH	0.48	0.51	n.a.	2	0.50	-0.02
PGI	1.00	0.0	n.a.	1	0.00	M
PGM	0.90	0.09	0.01	3	0.18	-0.09
PGD	0.78	0.22	n.a.	2	0.34	-0.03
Means (s.d.)				2.20 (0.84)	0.31 (0.19)	
Magdalena						
AAT	0.78	0.19	0.03	3	0.36	0.10*
ADH	0.21	0.79	n.a.	2	0.34	-0.03
PGI	0.93	0.07	n.a.	2	0.12	-0.03
PGM	0.96	0.04	0.005	3	0.08	-0.04
PGD	0.69	0.31	n.a.	2	0.42	0.01
Means (s.d.)				2.4 (0.55)	0.32 (.19)	

¹Asterisks indicate significance of G-test for Hardy-Weinberg disequilibrium: ***p<.001, **<.01, *>.05; M=monomorphic.

Table 7-8. Homogeneity statistics for six loci of four *Plumeria rubra* populations in a study of effects of forest fragmentation in northwestern Costa Rica (1997 progeny)

Locus	chi-square, probability	df,	G-square, probability	df,
AAT	22.20, 3, <0.000		24.8, 3, <0.000	
ADH	128.5, 3, <0.000		132.5, 3, <0.000	
PGI	21.2, 3, <0.000		32.8, 3, <0.000	
PGM1	88.0, 3, <0.000		73.9, 3, <0.000	
PGD	41.2, 3, <0.000		42.1, 3, <0.000	

Table 7-9. Estimates of Wright's statistics and mN between 1997 progeny of four populations of *P. rubra* located in Guanacaste province, northwestern Costa Rica.

Locus	Sample size	F_{is}	F_{it}	F_{st}	mN
AAT	2098	-0.0381	-0.0234	0.0142	17.35
ADH	2132	-0.0070	0.0484	0.0550	4.29
PGI	2244	0.0158	0.0313	0.0158	15.60
PGM	2252	-0.0534	-0.0169	0.0347	6.96
PGD	1778	-0.0070	0.0303	0.0371	6.49
Mean	2101	-0.0189	0.0172	0.0355	6.70

Table 7-10. Pairwise estimates of Nei's unbiased genetic distance (below diagonal) and geographic distances (km, above diagonal) between 1997 progeny of four populations of *Plumeria rubra* located in Guanacaste province, northwestern Costa Rica

	R. Corobicí	San Isidro	Praderas	Magdalen a
R. Corobicí		6.472	3.423	8.072
San Isidro	0.0087		3.593	6.472
Praderas	0.0144	0.0232		8.429
Magdalena	0.0120	0.0375	0.0280	

Table 7-11. Estimates of allele frequencies, number of alleles, expected heterozygosity, fixation indices and G-test for Hardy-Weinberg disequilibria in 1998 progeny of six populations of *P. rubra* in northwestern Costa Rica

Population / Locus	Allele frequencies			A	H_e (Ncl)	F_{is}^1
	p	q	r			
Qbrda. Palmira						
AAT	0.65	0.29	0.06	3	0.49	-0.08
ADH	0.41	0.59	n.a.	2	0.48	0.02
PGI	0.86	0.14	n.a.	2	0.24	0.36***
PGM	0.97	0.03	0.0	2	0.06	-0.03
PGD	0.55	0.45	n.a.	2	0.50	-0.20*
means				2.2(0.45)	0.36 (0.20)	
Río Corobicí						
AAT	0.81	0.15	0.04	3	0.31	0.13*
ADH	0.38	0.62	n.a.	2	0.47	0.09
PGI	0.94	0.06	n.a.	2	0.12	-0.07
PGM	0.89	0.04	0.07	3	0.20	-0.09
PGD	0.52	0.48		2	0.50	0.01
means				2.4 (0.55)	0.32 (0.16)	
San Isidro						
AAT	0.62	0.25	0.13	3	0.53	-0.12
ADH	0.50	0.50	n.a.	2	0.50	0.05
PGI	0.93	0.07	n.a.	2	0.13	-0.07
PGM	0.84	0.13	0.02	3	0.27	-0.16**
PGD	0.53	0.47		2	0.50	-0.11
means				2.4 (0.55)	0.39 (.18)	
Praderas						
AAT	0.65	0.05	0.30	3	0.49	0.13*
ADH	0.43	0.57	n.a.	2	0.49	0.01
PGI	0.99	0.01	n.a.	2	0.02	-0.01
PGM	0.97	0.03	0.0	2	0.06	-0.03
PGD	0.74	0.26	n.a.	2	0.38	-0.08
means				2.20 (0.45)	0.29 (0.23)	
Magdalena						
AAT	0.83	0.16	0.01	3	0.29	0.21**
ADH	0.23	0.76	n.a.	2	0.36	0.04
PGI	0.94	0.06	n.a.	2	0.12	0.02
PGM	0.986	.005	0.009	3	0.03	-0.01
PGD	0.66	0.34		2	0.45	0.01
Mean				2.44 (0.55)	0.25(.17)	
Sandillal						
AAT	0.98	0.02	0.0	2	0.04	-0.02
ADH	0.03	0.97	n.a.	2	0.06	-0.03
PGI	0.85	0.15	n.a.	2	0.25	-0.17*
PGM	0.96	0.04	0.0	2	0.08	-0.04
PGD	0.64	0.36	N.A.	2	0.46	-0.16
Mean				2.0 (0.00)	0.18(.18)	

¹Asterisks indicate significance of G-test for Hardy-Weinberg disequilibrium: ***p<.001, **<.01, *>.05; M=monomorphic locus.

Table 7-12. Estimates of Wright's statistics and mN between 1998 progeny of six populations of *P. rubra* located in Guanacaste province, northwestern Costa Rica.

Locus	Sample size	F_{is}	F_{it}	F_{st}	mN
AAT	2356	0.0218	0.1118	0.0919	2.46
ADH	2902	0.0391	0.1437	0.1089	2.04
PGI	2704	0.0304	0.0601	0.0306	7.91
PGM	2732	-0.0837	-0.0405	0.0399	6.02
PGD	2346	-0.0900	-0.0595	0.0279	8.69
Means	2608	-0.0161	0.0539	0.0689	3.4
Means without Sandillal	2396	-0.0036	0.0358	0.0393	6.11

Table 7-13. Pairwise estimates of Nei's unbiased genetic distance (below diagonal) and geographic distances (km) between six populations of *P. rubra* located in Guanacaste province, Costa Rica

	Q. Palmira	R. Corobicí	San Isidro	Praderas	Magdalena	Sandilial
Q. Palmira		0.99	6.574	4.119	7.464	8.733
R. Corobicí	0.0110		6.472	3.423	8.072	8.66
San Isidro	0.0089	0.0138		3.593	6.472	6.454
Praderas	0.0325	0.0336	0.0274		8.429	8.745
Magdalena	0.0195	0.0123	0.0351	0.0307		1.304
Sandilial	0.0626	0.0415	0.0897	0.0763	0.0174	

Table 7-14. Estimates of allele frequencies, number of alleles, expected heterozygosity, fixation indices and G-test for Hardy-Weinberg disequilibria in 1999 progeny of five populations of *P. rubra* in northwestern Costa Rica

Population / Locus	Allele frequencies			A	H_e (Nei)	F_{is} ¹
	p	q	r			
Qbrda. Palmira						
AAT	0.68	0.22	0.09	3	0.47	0.10
ADH	0.52	0.48	n.a.	2	0.50	0.17*
PGI	0.90	0.10	n.a.	2	0.19	0.00
PGM	0.95	0.03	0.02	3	0.10	0.06
PGD	0.61	0.39		2	0.48	0.02
means				2.4 (0.55)	0.35 (0.19)	
Río Corobicí						
AAT	0.76	0.16	0.08	3	0.39	-0.20
ADH	0.48	0.52	n.a.	2	0.50	-0.42***
PGI	0.99	0.01	n.a.	2	0.01	0.00
PGM	0.86	0.07	0.07	3	0.24	-0.05
PGD	0.59	0.41		2	0.48	-0.09
means				2.4 (0.55)	0.32 (0.21)	
Magdalena						
AAT	0.95	0.04	0.01	3	0.09	-0.05
ADH	0.25	0.75	n.a.	2	0.37	-0.03
PGI	0.97	0.03	n.a.	2	0.06	-0.03
PGM	0.95	0.05	0.0	2	0.10	-0.05
PGD	0.84	0.16		2	0.27	-0.10
Mean				2.2 (0.45)	0.18(0.14)	
Pachanga						
AAT	0.75	0.17	0.08	3	0.40	-0.09
ADH	0.42	0.58	n.a.	2	0.49	-0.20**
PGI	0.99	0.01	n.a.	2	0.02	0.66**
PGM	0.98	0.02	0.0	2	0.04	-0.02
PGD	0.44	0.56		2	0.49	0.15
Mean				2.20 (0.45)	0.29(0.24)	
Sandillal						
AAT	0.97	0.03	0.0	2	0.06	-0.03
ADH	0.04	0.96	n.a.	2	0.08	-0.04
PGI	0.91	0.09	n.a.	2	0.16	-0.09
PGM	0.98	0.02	0.0	2	0.04	-0.02
PGD	0.75	0.25		2	0.37	-0.18
Mean				2.0 (0.0)	0.14(0.14)	

¹Asterisks indicate significance of G-test for Hardy-Weinberg disequilibrium: ***p<.001, **<.01, *>.05; M=monomorphic locus.

Table 7-15. Estimates of Wright's statistics and mN between 1999 progeny of five populations.

Locus	Sample size	F_{is}	F_{it}	F_{st}	mN
AAT	1294	-0.0417	0.0299	0.0687	3.39
ADH	1332	-0.1197	0.0347	0.1379	1.56
PGI	1402	-0.0126	0.0252	0.0373	6.45
PGM	1360	-0.0028	0.0244	0.0271	8.98
PGD	1072	-0.0283	0.0570	0.0829	2.76
Mean	1292	-0.0559	0.0393	0.0902	2.52
Mean without Sandillal, Pachanga	793	-0.0513	0.0053	0.0538	4.39

Table 7-16. Pairwise estimates of Nei's unbiased genetic distance (below diagonal) and geographic distances (km) between five populations of *P. rubra* located in Guanacaste province, Costa Rica.

	Q. Palmira	R. Corobicí	Magdalena	Pachanga	Sandillal
Q. Palmira		0.99	7.464	5.914	8.733
R. Corobicí	0.0058		8.072	5.727	8.66
Magdalena	0.0452	0.0355		1.85	1.304
Pachanga	0.0137	0.0088	0.0551		3.129
Sandillal	0.0769	0.0643	0.0122	0.0688	

Table 7-17. Estimates of current (1998, 1999) gene flow from the Magdalena to Sandillal populations based on four alleles absent from the Sandillal population

Allele	q_i^1	q_i^2 1998	m^3 (1998)	q_i 1999	m (1999)
AAT-B	0.18	0.02	0.11	0.03	0.17
AAT-C	0.03	0	0	0	0
ADH-A	0.26	0.03	0.12	0.04	0.15
PGM-B	0.12	0.04	0.33	0.02	0.17
Mean			0.14		0.12
s			0.14		0.08

¹allele frequency in source population, *i.e.* Magdalena; ²allele frequency in target (Sandillal) progeny; ³ m =proportion of immigrant alleles

Table 7-18. Estimates of mating system parameters in seven population of *Plumeria rubra* located in forest fragments in northwestern Costa Rica

Population / year	F (s.e.)	t_m (s.e.)	t_s (s.e.)	tm-ts	r_t	r_p
Quebrada Palmira						
pooled data	0.001 (0.001)	0.944 (0.076)	0.894 (0.068)	0.050 (0.025)	0.192 (0.114)	0.99 (0.038)
1998	0.002 (0.001)	0.997 (0.026)	0.945* (0.019)	0.052 (0.019)	0.105 (0.014)	0.99 (0.043)
1999	0.001 (0.102)	0.944 (0.083)	0.894 (0.065)	0.050 (0.036)	0.192 (.033)	.990 (.033)
Río Corobicí						
pooled	0.001 (.000)	0.998 (.009)	0.947* (0.010)	0.051 (0.009)	0.112 (0.000)	0.990 (0.055)
1997	0.000 (.000)	0.996 (0.015)	0.942* (0.018)	0.053 (0.013)	0.110 (0.004)	0.976 (0.053)
1998	0.000 (0.000)	0.984 (0.020)	0.924* (0.014)	0.060 (0.013)	0.109 (0.013)	0.990 (0.003)
1999	0.000 (0.000)	1.000 (0.018)	0.975* (0.010)	0.025 (0.011)	0.120 (0.012)	0.99 (0.012)
Río Tenorio (San Isidro)						
pooled	0.001 (0.000)	0.997 (0.013)	0.966* (0.009)	0.031 (0.009)	0.120 (0.001)	0.784 (0.105)
1997	0.002 (0.002)	0.950 (0.069)	0.932 (0.046)	0.018 (0.032)	0.165 (0.106)	0.911 (0.108)
1998	0.001 (0.000)	1.0 (0.000)	0.979* (0.005)	0.021 (0.005)	0.116 (0.000)	0.739 (0.073)
Río Tenorio (Praderas)						
pooled	0.000 (0.000)	0.889 (0.092)	0.908* (0.041)	-0.019 (0.056)	0.179 (0.127)	0.990 (0.018)
1997	0.007 (0.010)	0.973 (0.139)	0.950 (0.065)	0.023 (0.086)	0.125 (0.037)	0.990 (0.029)
1998	0.000 (0.000)	0.879 (0.070)	0.893* (0.031)	-0.014 (0.041)	0.210 (0.001)	0.990 (0.001)
Río Magdalena						
pooled	0.000 (0.000)	0.914 (0.054)	0.901* (0.024)	0.013 (0.031)	0.125 (0.034)	0.964 (0.071)
1997	0.000 (0.000)	0.951 (0.039)	0.910* (0.023)	0.041 (0.019)	0.131 (0.034)	0.99 (0.007)
1998	0.000 (0.000)	0.730 (0.170)	0.859* (0.063)	-0.129 (0.073)	0.252 (0.194)	0.99 (0.023)
1999	0.002 (0.001)	0.988 (0.085)	0.945 (0.034)	0.043 (0.056)	0.105 (0.008)	0.593 (0.169)
La Pachanga						
1999	0.000 (0.000)	1.0 (0.003)	0.959* (0.015)	0.041 (0.015)	0.107 (0.001)	0.764 (0.106)
Sandíbal						
pooled	0.013 (0.068)	0.999 (0.000)	0.960* (0.007)	0.040 (0.007)	0.103 (0.000)	0.82 (0.044)
1998	0.027 (0.122)	0.986* (0.006)	0.944* (0.012)	0.042 (0.007)	0.103 (0.004)	0.969 (0.019)
1999	0.013 (0.010)	1.0 (0.000)	0.974* (0.005)	0.026 (0.005)	0.103 (0.000)	0.646 (0.147)

* indicates significant tm and ts estimates (i.e. $t \pm 2s.e. < 1.0$).

Table 7-19. Linear regression of log of number of capsules tree⁻¹ on dbh and neighbourhood density index of trees of *Plumeria rubra* located in six forest fragments in northwestern Costa Rica

Source	df	Mean square	F	R ²	significance ¹
Model	2	20.46	28.83		<0.0001
Error	73	0.71			
Variable	df	parameter estimate and standard error	t		
Intercept	1	1.470±0.2631	5.59		<0.0001
dbh	1	0.056±0.0079	7.12		<0.0001
density index	1	0.056±0.0253	2.22		0.0295

¹i.e. probability of a higher value of 't' or 'F' under the null hypothesis

Table 7-20. Mean numbers of capsules tree⁻¹ in trees of *Plumeria rubra* located in seven forest fragments in northwestern Costa Rica

Fragment	sample size	mean number of capsules tree ⁻¹ (standard error)
Quebrada Palmira	10	39.8±12.19
Corobici	21	28.1±6.64
San Isidro	31	56.3±10.46
Praderas	19	24.0±4.74
Magdalena	20	23.0±5.39
Pachanga	6	43.8±12.45
Sandillal	4	98.0±25.64

Table 7-21. Analysis of covariance of effects of fragment and dbh on mean of log numbers of capsules tree⁻¹ in six forest fragments of *Plumeria rubra* located in northwestern Costa Rica

Source	df	Mean square (Type III)	F	significance ¹
Fragment	5	1.0	1.47	0.21
dbh	1	29.2	40.71	<0.0001
Error	71	0.72		

¹i.e. probability of a higher value of 'F' under the null hypothesis

Table 8-1. Summary of breeding systems and self-compatibility of 114 tree species of Hacienda La Pacífica, Cañas, Guanacaste province, Costa Rica

Sexual System	Incompatibility			Totals
	Self-incompatible	Self compatible	No data	
Hermaphroditic	26	9	44	79
Monoecious	7	0	4	11
Andromonoecious	1	1	0	2
Gynodioecious	0	1	0	1
Totals	34	11	48	93
Dioecious	21			114

Table 8-2. Summary pollen and diaspore vectors of 114 tree species of Hacienda La Pacífica, Cañas, Guanacaste province, Costa Rica

Pollen vector	Diaspore vector			Totals
	Biotic	Abiotic or no apparent dispersal mechanism	Insufficient data	
bats	4	5		9
hawkmoths	4	7		11
medium-to-large bees	11	14	1	26
wind	2	0		2
fig-wasps	4	0		4
<i>Subtotal</i>	25 (23) ²	26	1	52
hummingbirds	2	0		2
moths, small bees, diverse small insects	33	11		44
Beetles	3	0		3
<i>Subtotal</i>	36	11		47
insufficient data	3	7	5	15
Totals	66(64)	44	6	116(114) ¹

¹figures in parentheses are totals after adjusting for repetition of *Inga vera* and *Bombacopsis guineata*, both of which appear as both hawkmoth and bat pollinated

Table 8-3. Two by two classification of pollen and diaspore dispersal vectors of x forest tree species in Cañas, Guanacaste province, Costa Rica

Pollen vector	Diaspore vector		Totals
	Biotic	Abiotic	
	<i>observed (expected) numbers of species by class</i>		
Long-distance ¹	25 (31.7)	26 (19.3)	51
Short-distance	36 (29.3)	11 (17.7)	47
Totals	61	37	98

¹bats, hawkmoths, medium to large bees, fig-wasps, wind; 2moths, small bees, small diverse insects, beetles. Hummingbirds not considered long-distance pollinators because of lack of information on territoriality of the species in question (Bawa, 1990). Territoriality could lead to low expected pollen movement even for strong flyers

APPENDIX II: DETAILS OF LABORATORY PROCEDURES

Anacardium excelsum

Enzyme extraction

Samples were prepared in batches of 20. Fresh, healthy leaves were cut with scissors from seedlings in the greenhouse, placed in plastic ziplock bags and taken immediately to the laboratory. During final preparation of sample tubes in the laboratory, the leaves were held for approximately five minutes at 4°C. From each leaf, 4 sample disks were cut using the lid of an 1.5ml Eppendorf tube, for an approximate area of 2cm² (for tough leaves, an equivalent area was cut with scissors; for ease of preparation, succulent leaves were used whenever possible). The 20 samples were then placed on ice in 1.5ml Eppendorf microcentrifuge tubes. For grinding, each tube with its contained sample was dipped momentarily in liquid nitrogen, using tongs. When the nitrogen had almost evaporated, the sample was crushed and ground for around five seconds using plastic pestles. Subsequently, 120µl of Liengsiri *et al.*'s (1990) extraction buffer #9 (Tris-HCl pH 8.0 0.1M, Ascorbic acid 0.01M, Cysteine HCl 5.4mM, MgCl₂ 0.2%, CaCl₂ 0.2%, PEG 20M 1%, Sucrose 0.5M, Tween 80 1%, Tergitol 1%, β-mercaptoethanol 0.3%, pH adjusted with 1M tris) was added to the tube using a Gilson pipette. Grinding was continued until the extraction buffer melted and a thick slurry was obtained. The tube was then capped and placed back on ice. After centrifugation (10000 rpm, 10 minutes, 4°C), two filter paper (Whatman #3) wicks, each 2.5x8.0mm, were dipped in each sample extract and then placed in a different Eppendorf tube. The tubes with wicks were then packed in ice in ziplock brand plastic containers and placed at -80°C until needed.

Gel preparation

The pH5.7 histidine-citrate buffer system described by Wendel and Weeden (System 1) was used. The electrode buffer of this system has molarities 0.065 (histidine free base) and 0.019M, (citric acid, monohydrate), whilst the gel buffer is prepared as a 1:6 dilution of the electrode buffer. According to Wendel and Weeden, this results in 0.009M molarity histidine (i.e. 0.065/7) and 0.006M citric acid. However, the latter molarity does not correspond to the dilution ratio, which implies final molarity of 0.019/7 = 0.0027M. In the preparation of

the gel buffer, the dilution ratio specified by Wendel and Weeden was retained, i.e. molarities used were 0.009M histidine and 0.0027M citric acid.

For preparation of each 12.5% starch gel, 170ml of the gel buffer was placed in a 1/ Ehrlemeyer flask and heated for 2 minutes and 40 seconds in a microwave oven (Magic Chef, 850W), whilst 110ml of the buffer was mixed with 35g of hydrolized potato starch (Connaught Laboratories, Ontario) in a 1/ vacuum flask. The boiling buffer was then poured into the starch suspension, and shaken vigourously for about 4 seconds. It was then returned to the microwave for a further 2 minutes and 10 seconds (i.e. until 'glassy'), removing after 30, 60 and 90 seconds in order to swirl the mixture, for greater uniformity. After removal, the mixture was degassed using a Nalgene tap-attached aspirator pump for approximately 25 seconds, and then poured into a previously assembled gel mould with internal dimensions 8x22x1cm. After 1 hour, the gel was covered with cling film and cooled to 4°C before use (when poured in the evening for use the next morning, the mould with gel was stored in a ziplock bag in the dark at room temperature).

Gel loading and running

Gels were loaded on ice with the previously prepared wicks, and run initially for 15-20 minutes at 25mA on one of the following power units: Bio-Rad Model 250/2.5 Power Supply, E-C Apparatus Corporation VWR 105, E-C Apparatus Corportaion EC105, Gelman Sciences Inc. Deluxe Regulated Power Supply. The gels were then dewicked, and current raised to 50mA. Gels were then run for 4 hours after dewicking. As gels were run in a 25°C laboratory, ziplock bags with crushed ice were placed on each gel and changed every 15 minutes.

Staining

On termination of the run, each gel was sliced into 1mm or 2mm slices using monofilament sewing thread and 1mm thick plastic guide strips. The slices were then transferred to staining trays (Rubbermaid stackable storage trays, 22.5x7.5cm), and stained using the following buffers (quantities given are for one gel slice).

Adenylate kinase

Tris HCl 0.2M, pH8.0: 7.5ml
ADP: 80mg

Glucose: 90mg
 G6PDH: 40 units
 Hexokinase: 264 units
 1% BSA: 0.5ml
 10% MgCl₂: 0.5ml
 1% MTT: 1ml
 1% NADP: 1ml
 1% PMS: 0.25ml

After mixing, the stain buffer was combined with 12.5ml 2% agarose. The latter had been prepared previously and held in a 60°C waterbath until needed.

Source: adapted from Liengsiri *et al.* (1990) and Wendel and Weeden (1990).

Phosphogluconate dehydrogenase

Tris HCl 0.2M, pH 8.0: 5ml
 Phosphogluconic acid, barium salt: 15mg
 1% MgCl₂: 0.5ml
 1% MTT: 1ml
 1% NADP: 1ml
 1% PMS: 0.25ml

After mixing, the stain buffer was combined with 12.5ml 2% agarose. The latter has been prepared previously and held in a 60°C waterbath until needed.

Source: Liengsiri *et al.*, 1990.

UTP-glucose-1-phosphate uridylyltransferase

Tris HCl 0.2M, pH 8.0: 12.5ml
 Tetrasodium pyrophosphate: 10mg
 Phosphogluconic acid, barium salt: 15mg
 UGDP: 25mg
 1% BSA: 0.5ml
 PGM: 50 units
 G6PDH: 40 units
 1% MgCl₂: 0.5ml
 1% MTT: 0.5ml
 1% NADP: 0.5ml
 1% PMS: 0.25ml

After mixing, the stain buffer was combined with 12.5ml 2% agarose. The latter has been prepared previously and held in a 60°C waterbath until needed.

Source: Hodgkiss, 2001

Phosphoglucosmutase

Tris HCl 0.2M, pH 8.0: 25ml

Glucose-1-phosphate: 250mg

G6PDH: 25 units

1% BSA: 0.5ml

1% MgCl₂: 0.5ml

1% MTT: 1.0ml

1% NADP: 1.0ml

1% PMS: 0.25ml

Source: adapted from Hodgkiss (1991), Liengsiri *et al.*, 1990.

Malate dehydrogenase

Tris HCl 0.2M, pH 8.0: 25ml

DL-Malic acid 0.5M pH7.0: 2ml

1% MTT: 0.5ml

1% NAD: 0.5ml

1% PMS: 0.25ml

Source: adapted from Liengsiri *et al.*, 1990.

Leucine aminopeptidase

LAP stain buffer, pH6.0 (1:1 0.2M Tris HCl pH8.0, 0.2M Maleic anhydride): 25ml

4% L-leucine β-naphthyl-amide: 2ml

Fast K salt: 25mg in 2ml N,N-dimethylformamide

10% MgCl₂: 0.5ml

Source: adapted from Liengsiri *et al.*, 1990, Wendel and Weeden, 1990.

After adding the stain buffers, the gels were incubated for 1 hour (MDH) or 2 hours (others) at 37°C. After removal from the incubator, LAP gels were rinsed in 50% glycerol before photographing.

Plumeria rubra*Enzyme extraction*

Seeds of each pod-within-family were germinated on moistened chromatography paper. One to two days after germination, radicle tips of approximately 0.5cm length were cut using a scalpel (samples were prepared in batches of 20). The tips were transferred to prelabelled 1.5ml Eppendorf microcentrifuge tubes held on ice. Subsequently, 30µl of Liengsiri *et al.*'s (1990) extraction buffer #9 (Tris-HCl pH 8.0 0.1M, Ascorbic acid 0.01M, Cysteine HCl 5.4mM, MgCl₂ 0.2%, CaCl₂ 0.2%, PEG 20M 1%, Sucrose 0.5M, Tween 80 1%, Tergitol 1%, β-mercaptoethanol 0.3%, pH adjusted with 1M tris) was added to the tube using a Gilson pipette and the sample was crushed and ground briefly (10-15 seconds) using

a plastic pestle. Two filter paper (Whatman #3) wicks, each 2.5x8.0mm, were dipped in each sample extract and then placed in a different Eppendorf tube. The tubes with wicks were then packed in ice in ziplock brand plastic containers and placed at -80°C until needed.

Gel preparation

Two running buffer systems were used: Pitel and Cheliak's (1984) pH 7.0 Histidine system (electrode buffer: 0.125M Tris, pH adjusted to 7.0 with 1M citric acid; gel buffer: Histidine HCl 0.05M, EDTA 1.4mM, pH adjusted to 7.0 with 1M Tris) and Ridgeway *et al.*'s (1970) lithium borate / tris citrate system (electrode buffer: lithium hydroxide 0.06M, boric acid 0.3M; Tris 0.03M, citric acid 0.005M, 1% electrode buffer, pH adjusted to 8.5 with 1N NaOH). The 12.5% starch gels were prepared as for *A. excelsum* (see above).

Gel loading and running

Gels were loaded on ice with the previously prepared wicks, and run initially for 15-20 minutes at 70V and 100V for respectively the histidine and lithium borate systems, using one of the following power units: Bio-Rad Model 250/2.5 Power Supply, E-C Apparatus Corporation VWR 105, E-C Apparatus Corporation EC105, Gelman Sciences Inc. Deluxe Regulated Power Supply. The gels were then dewicked, and voltage doubled. Gels were then run for 4 hours after dewicking. The gels for the 1998 and 1999 collections run in a 25°C laboratory; ziplock bags with crushed ice were placed on each gel and changed every 15 minutes. Gels for the 1997 collection were run in a 4°C cold room. In this case, icepacks were changed every hour.

Staining

On termination of the run, each gel was sliced into 2mm slices using monofilament sewing thread and 1mm thick plastic guide strips. The slices were then transferred to staining trays, and stained using the following buffers (quantities given are for one gel slice):

Histidine system

Alcohol dehydrogenase

0.2M Tris-HCl, pH 8.0: 25ml
95% ethanol: 5ml
1% NAD: 0.5ml
1% NBT: 0.5ml
0.5% PMS: 0.5ml

Source: Liengsiri *et al.*, 1990

Phosphoglucomutase

Tris HCl 0.2M, pH 8.0: 25ml

Glucose-1-phosphate, disodium salt: 250mg

G6PDH: 25 units

1% MgCl₂: 0.5ml

1% MTT: 0.5ml

1% NADP: 0.5ml

0.5% PMS: 0.5ml

Source: adapted from Liengsiri *et al.*, 1990, Wendel and Weeden, 1990.

Phosphogluconate dehydrogenase

Tris HCl 0.2M, pH 8.0: 5ml

Phosphogluconic acid, barium salt: 15mg

1% MgCl₂: 1ml

1% MTT: 1ml

1% NADP: 1ml

0.5% PMS: 0.5ml

Source: Liengsiri *et al.*, 1990

Lithium borate / tris citrate system:

Aspartate aminotransferase

Pyridoxal-5-phosphate: 2mg

Fast Blue BB salt: 50mg

AAT substrate solution: 25ml

(For 500ml of AAT substrate solution: dissolve 2.65g L-aspartic acid, 0.35g L-ketoglutaric acid in about 300ml of 0.2M Tris-HCl, adjust to pH 8.0 with 1N NaOH and top up to 500ml with 0.2M Tris-HCl pH 8.0).

Source: Liengsiri *et al.*, 1990

Glucose-phosphate isomerase

Tris HCl 0.2M, pH 8.0:

Fructose-6-phosphate, disodium salt: 12.5mg

Glucose-6-phosphate dehydrogenase: 5 units

1% MgCl₂: 0.5ml

1% MTT: 0.5ml

1% NADP: 0.5ml

0.5% PMS: 0.5ml

Source: Liengsiri *et al.*, 1990

After adding the stain buffers, the gels were incubated for 1 hour at 37°C.

APPENDIX III: MODIFIED MATERNAL GENOTYPES OF *PLUMERIA RUBRA*

As explained in Chapter 7, some inferred maternal genotypes of *P. rubra* were reassigned based on capsular segregation ratios. These are detailed below (Table A2-1).

Table A3-1. Reassigned maternal inferred genotypes of *Plumeria rubra*.

Population	Family	Inferred maternal multilocus genotype	Modified maternal multilocus genotype	Justification for modification
Quebrada Palmira	Cemetery tree	BBABAAAABAB	BBABAAA Δ AB	Ratios of AA:AB progeny in pods 563 (7:2), pods 564 (6:1).
Río Corobicí	1000	Δ BABAAAAAB	$\Delta\Delta$ ABAAAAAB	Ratios of AA:AB progeny in pod 139 (6:0).
	1001	ABBBA Δ ABAB	ABBBA $\Delta\Delta$ AB	Ratios of AA:AB progeny in pods 19 (4:1), 20(7:0), 21 (8:0).
	1002	ACBBA $\Delta\Delta$ BAA	ACBBA $\Delta\Delta\Delta$ AA	Ratios of AA:AB progeny in pod 506 (6:0).
	1004	Δ CBBA Δ ABAB	$\Delta\Delta$ BBA Δ ABAB	Ratios of AA:AC progeny in pods 147 (5:0), 513 (4:0), 515 (6:0).
	1005	AAABAA Δ CAB	AAABAAA $\Delta\Delta$ AB	Ratios of AA:AC progeny in pods 518 (4:0), 520 (6:0).
	1010	AAAB Δ BACAB	AAAB Δ BACAB	Ratios of AA:AB progeny in pod 169 (19:0).
	1011	Δ BABAAAAAB	Δ BABAAAAAB	Ratios of AA:AB progeny in pods 6 (6:0), 550 (7:0).
	1021	AB Δ BAAAAAB	ABBBAAAAAB	Ratio of BB:AB progeny in pod 180 (13:0).
Magdalena	4010	ABBB Δ BAAAAB	ABBB $\Delta\Delta$ AAAAB	Ratio of AA:AB progeny in pod 73 (12:0).
La Pachanga	CS04	Δ BABAAAAAB	$\Delta\Delta$ ABAAAAAB	Ratio of AA:AB progeny in pod 476(5:0).
	CS07	AAABAA Δ BAB	AAABAAA $\Delta\Delta$ AB	Ratios of AA:AB progeny in pods 480 (5:0), 484 (4:0).

Table A3-1. Reassigned maternal inferred genotypes of *Plumeria rubra*. (continued)

Population	Family	Inferred maternal multilocus genotype	Modified maternal multilocus genotype	Justification for modification
Sandillal	00	AABB A BAAAB	AABB A A A AAAB	Ratios of AA:AB progeny in pods 305 (5:0), 229 (4:0), 300 (5:0)
	03	A BBBAAAAAA	A A BBBAAAAAA	Ratio of AA:AB progeny in pod 318 (7:0).

APPENDIX IV: CLASSIFIED SPECIES LISTS FOR HACIENDA LA PACÍFICA

Introduction

Jiménez et al. (1987) listed plant species of the *hacienda* La Pacífica. The species in question are listed below according to breeding system and self-incompatibility (Table A4-1), pollen and diaspore dispersal vectors (Table A4-2) and density (number of trees hectare⁻¹). The following exotic species and possible partial domesticates were omitted: *Mangifera indica*, *Spondias purpurea* (Anacardiaceae), *Thevetia peruviana* (Apocynaceae), *Jacaranda mimosifolia* (Bignoniaceae), *Bixa orellana* (Bixaceae), *Caesalpinia pulcherrima*, *Cassia fistula*, *Delonix regia*, (Caesalpinoidea), *Leucaena glauca*, *Tamarindus indica* (Mimosoideae), *Encalyptus* spp., *Eugenia salamensis*, *Psidium* spp. (Myrtaceae), *Glicinia sepium* (Papilionoideae), *Citrus* spp. (Rutaceae), *Gmelina arborea*, *Tectona grandis* (Verbenaceae), Palmae, identifications disputed by Janzen and Liesner (1980) (*Agonandra obtusifolia* (Opiliaceae)) and identification made to generic level only (*Maytenus* spp. (Celastraceae), *Inga* spp. (Mimosoideae), *Ficus* spp. (Moraceae) were also omitted. Sources and methodological details specific to each table are detailed below.

Species by breeding system and self-compatibility

Species are listed according to breeding system and incompatibility in Table A4-1. Family classifications follow Janzen and Liesner (1980). Identification of breeding systems was based on the following sources: Area de Conservación Guanacaste (undated); Bawa and Opler (1975), Bawa *et al.* (1985a), Burger and Huft (1995); Center for Tropical Forest Science (undated); Croat (1979), Bullock (1985), Frankie *et al.* (1983), Haber and Frankie (1989), Janzen and Liesner (1980), Little and Wadsworth (1964), Little *et al.* (1974), Witsberger *et al.* (1982). Self-compatibility classification was based on both published controlled pollination data (Bawa (1974), Bawa *et al.* (1985b), Bullock (1985)) and data from genetic markers, as referenced in Table A4.1.

Species by pollen and seed dispersal vectors

Species were classed as pollinated by bat, hawkmoth, medium to large bees, hummingbirds, small insects (e.g. small bees, moths, butterflies), beetles and wind (Table A4-2). Many of the classifications were made on the basis of detailed published studies of the zone (particularly, Heithaus *et al.* (1975), Haber and Frankie (1989) and Frankie *et al.* (1983)). Classifications unattributed to these or other specific sources (see Table A4-2) were based on floral

morphology, as noted in Table A4-2. In the great majority of cases, these were small-flowered species classed as pollinated by small insects. The following species could not be classified due to lack of information on floral morphology: *Erythroxylon havanense* (Erythroxylaceae), *Bernardia nicaraguensis*, *Capparis frondosa*, *Capparis incana* (Capparidaceae), *Croton niveus* (Euphorbiaceae), *Albizia caribaea*, *A. adinocephala* (Mimosoideae), *Willardia schiedeana* (Papilionoideae).

Diaspore dispersal mechanisms were classified as either biotic or abiotic, the latter including species with no apparent dispersal mechanisms. Classifications unattributed to specific sources were based on fruit morphology, as noted in Table A4-2.

Species density

The mean species density for riparian and upland sites, based on Glander and Nisbett's (1996) data was 2.81 ha⁻¹ and 8.93 ha⁻¹, respectively (Table A4-3). The respective medians were 0.5 ha⁻¹ and 0.2ha⁻¹. The divergence between the two measures reflects individual extreme observations (e.g., *Guaruma ulmifolia*, *Lonchocarpus minimiflorus*). As species listed by Jiménez *et al.* (1987) but not registered in Glander and Nisbett's inventories were assigned zero densities in both site types, the densities in the upland site are likely to be biased downwards by inclusion of species unlikely to occur in dry conditions, e.g. *Anacardium excelsum*, *Pithecellobium longifolium*. This does not apply so clearly in the case of the riparian sites, as these may contain dry microsites, particularly at the forest margins.

Table A4-1. Tree species of Hacienda La Pacifica, Guanacaste province, Costa Rica, classified according to breeding system and self-compatibility

Breeding System Family	Incompatibility		
	Self-incompatible	Self compatible	No data
Hemaphroditic			
Annonaceae	<i>Sapranthus balangá</i> ¹		<i>Annona purpurea</i> , <i>A. reticulata</i>
Apocynaceae	<i>Plumeria rubra</i>		<i>Stemmadenia obovata</i> , <i>Thevetia ovata</i>
Araliaceae	<i>Godmania aesculifolia</i> ¹ , <i>Tabebuia ochracea</i> ¹ , T. <i>rosea</i> ¹		<i>Siadodendron excelsum</i> <i>Crescentia alata</i> , <i>Tabebuia impetiginosa</i>
Bignoniaceae	<i>Cordia alliodora</i>		<i>Cordia bicolor</i>
Boraginaceae	<i>Bombacopsis quinata</i> , <i>Ochroma pyramidalis</i> ¹	<i>Ceiba pentandra</i> ³	<i>Ceiba aesculifolia</i> , <i>Pseudobombax septentratum</i>
Bombacaceae			
Capparidaceae			<i>Capparis frondosa</i> , <i>C. incana</i>
Celastraceae			<i>Merytenius segouianum</i>
Chrysobalanaceae			<i>Licania arborea</i>
Cochlospermaceae	<i>Cochlospermum utrifolium</i> ¹	<i>Curatella americana</i> ¹	
Dilleniaceae		<i>Muntingia</i>	
Elaeocarpaceae		<i>calabura</i> ¹ , <i>Sloanea</i> <i>terniflora</i> ¹	

¹Bawa, 1974; ²Murawski and Hamrick, cited in Nason and Hamrick, 1997; ³Heterostylous (Bullock, 1985), assumed self-incompatible; ⁴Bullock, 1985; ⁵Bawa *et al.* 1985; ⁶James *et al.* 1998; ⁷Loveless and Gullison; ⁸Nason and Hamrick, 1997; ⁹Murawski and Hamrick, 1991; ¹⁰Bawa and Opler 1975; ¹¹Wintersberger *et al.* 1982; ¹²Bawa and Webb, 1984

Table A4-1. Tree species of Hacienda La Pacifica, Guanacaste province, Costa Rica, classified according to breeding system and self-compatibility (continued)

Breeding System Family	Incompatibility		
	Self-incompatible	Self compatible	No data
Hemaphroditic Erythroxylaceae Flacourtiaceae	<i>Erythroxylon harenense</i> ²		
	<i>Casaria corymbosa</i> ⁴ , <i>C. tremula</i> ⁴	<i>Prokeia crua</i> ⁴	<i>Casaria aculeata</i>
			<i>Ocotea veraguensis</i>
Lauraceae			<i>Cassia emarginata</i> , <i>C. grandis</i> , <i>Schizolobium paralybium</i> , <i>Swartzia cubensis</i>
Leguminosae			<i>Acacia jamaicana</i> , <i>Albizia adinocephala</i> , <i>A. caribaea</i> , <i>A. guachepel</i> , <i>Inga vera</i> , <i>Lysiloma desmostachys</i> , <i>L. divaricatum</i> , <i>Pithecellobium longifolium</i> ,
Caesalpiinoideae	<i>Hymenaea courbari</i> ¹		<i>Acosmium panamense</i> , <i>Diphyssa robinoides</i> , <i>Lonchocarpus minimiflorus</i> , <i>Machaerium biowulatum</i> , <i>Platymiscium pleiostachys</i> , <i>Willardia schiedana</i>
Mimosoideae	<i>Enteolobium cyclocarpum</i> ¹ , <i>Pithecellobium saman</i> ¹		
Papilionoideae	<i>Andira inermis</i> ¹ , <i>Dalbergia retusa</i> ¹ , <i>Lonchocarpus costaricensis</i> ¹ , <i>L. eriocarinatus</i> ¹ , <i>Myrsopernum frutescens</i> ¹ , <i>Picadida carthagenensis</i> ¹ , <i>Pterocarpus rohr</i> ⁴		
Malpighiaceae		<i>Byronima crassifolia</i> ¹ , <i>Majipichia glabra</i> ¹	

¹Bawa, 1974; ²Murawski and Hamrick, cited in Nason and Hamrick, 1997; ³heterostylous (Bullock, 1985), assumed self-incompatible; ⁴Bullock, 1985; ⁵Bawa *et al.* 1985; ⁶James *et al.* 1998; ⁷Loveless and Gullison; ⁸Nason and Hamrick, 1997; ⁹Murawski and Hamrick, 1991; ¹⁰Bawa and Opler 1975; ¹¹Watsberger *et al.* 1982; ¹²Bawa and Webb, 1984

Table A4-1. Tree species of Hacienda La Pacifica, Guanacaste province, Costa Rica, classified according to breeding system and self-compatibility (continued)

Breeding System Family	Incompatibility		
	Self-incompatible	Self compatible	No data
Hemaphrodite			
Myrsinaceae		<i>Artisia revulvata</i> ¹	
Rubiaceae	<i>Hamelia patens</i> ⁵	<i>Calycophyllum candidissimum</i> ¹	<i>Exostema mexicanum</i> , <i>Coultara hexandra</i> , <i>Guelletaria macroperma</i>
Rutaceae			<i>Esenbeckia lionalis</i>
Sapotaceae			<i>Manilkara chicle</i> , <i>Mastichodendron capiri</i>
Sterculiaceae	<i>Guarzuma tomentosa</i> ¹		<i>Sterculia apetala</i>
Syracaceae			<i>Syzyx argenteus</i>
Tiliaceae	<i>Luehea candida</i> ¹ , <i>L. speciosa</i> ¹		<i>Apeiba tiborbou</i> , <i>Rabdera trinervis</i>
Verbenaceae			
Monoecious			
Anacardiaceae	<i>Spondias mombin</i> ¹		<i>Hura crepitans</i> , <i>Croton niveus</i>
Euphorbiaceae	<i>Cedrela odorata</i> ⁶ ; <i>Swietenia macrophylla</i> ⁷ ;		
Meliaceae			
Ulmaceae	<i>Ficus hondurensis</i> ⁸ , <i>F. elastica</i> ⁸ , <i>F. insipida</i> ⁸ , <i>F. ovalis</i> ⁸		<i>Trema micrantha</i>
Moraceae			
Sapindaceae			<i>Thouinidium decandrum</i>

¹Bawa, 1974; ²Murwski and Hamrick, cited in Nason and Hamrick, 1997; ³heterostylous (Bullcock, 1985), assumed self-incompatible; ⁴Bullcock, 1985; ⁵Bawa *et al.* 1985; ⁶James *et al.* 1998; ⁷Lowless and Gullison; ⁸Nason and Hamrick, 1997; ⁹Murwski and Hamrick, 1991; ¹⁰Bawa and Opler 1975; ¹¹Witsberger *et al.* 1982; ¹²Bawa and Webb, 1984

Table A4-1. Tree species of Hacienda La Pacifica, Guanacaste province, Costa Rica, classified according to breeding system and self-compatibility (continued)

Breeding System Family	Incompatibility		
	Self-incompatible	Self compatible	No data
Andromonoecious			
Anacardiaceae		<i>Anacardium excelsum</i>	
Leguminosae			
Caesalpinioideae	<i>Caesalpinia eriostachys</i> ¹		
Gynodioecious			
Moraceae		<i>Brosimum alcastrum</i> ⁹	
Diocious			
Anacardiaceae	<i>Astronium graveolens</i> ¹⁰		
Boraginaceae	<i>Cordia alliodora</i> ¹⁰		
Burseraceae	<i>Bursera simaruba</i> ¹⁰ , <i>B. tomentosa</i> ¹⁰		
Ebenaceae	<i>Dyospyros nicanrensis</i>		
Euphorbiaceae	<i>Bernardia nicaraguensis</i> ¹⁰		
Meliaceae	<i>Guarea excelso</i> ¹⁰ , <i>Trichilia americana</i> ¹ ; <i>T. birdi</i> ¹⁰ ; <i>T. martiana</i>		
Menispermaceae	<i>Hyperbaena tonduzi</i> ¹¹		
Moraceae	<i>Chlorophora tinctora</i> ¹⁰ , <i>Cecropia peltata</i> ¹⁰ , <i>Tropis racemosa</i> ⁸		
Polygonaceae	<i>Coccoloba caracasana</i> ¹⁰		
Rubiaceae	<i>Geniba americana</i> ⁵		

¹Bawa, 1974; ²Murawski and Hamrick, cited in Nason and Hamrick, 1997; ³heterostylous (Bullock, 1985), assumed self-incompatible; ⁴Bullock, 1985; ⁵Bawa *et al.* 1985; ⁶James *et al.* 1998; ⁷Loveless and Gullison; ⁸Nason and Hamrick, 1997; ⁹Murawski and Hamrick, 1991; ¹⁰Bawa and Opler 1975; ¹¹Witsberger *et al.* 1982; ¹²Bawa and Webb, 1984

Table A4-1. Tree species of Hacienda La Pacífica, Guanacaste province, Costa Rica, classified according to breeding system and self-compatibility (continued)

Breeding System Family	Incompatibility		
	Self-incompatible	Self compatible	No data
dioecious			
Sapindaceae	<i>Allophylus occidentalis</i> ³⁰		
Simaroubaceae	<i>Alvarado amorphoides</i> ¹⁰ , <i>Picramnia latifolia</i> ¹⁰ , <i>Quassia amara</i> , <i>Simarouba glauca</i> ¹⁰		

¹Bawa, 1974; ²Murawski and Hamrick, cited in Nason and Hamrick, 1997; ³heterostylous (Bullock, 1985), assumed self-incompatible; ⁴Bullock, 1985; ⁵Bawa *et al.* 1985; ⁶James *et al.* 1998; ⁷Loveless and Gullison; ⁸Nason and Hamrick, 1997; ⁹Murawski and Hamrick, 1991; ¹⁰Bawa and Opler 1975; ¹¹Watsberger *et al.* 1982; ¹²Bawa and Webb, 1984

Table A4-2. Tree species of Hacienda La Pacifica, Cañas, Guanacaste, Costa Rica, classified by pollinator type and diaspore dispersal vector

Pollen vector ¹	Diaspore dispersal vector ²		
	Biotic	Abiotic or no apparent dispersal mechanism	No data
Bats			
Bombacaceae		<i>Bombacopsis quinata</i> ^{3,22} <i>Caiba aesculifolia</i> ^{3,22} <i>C. pentandra</i> ^{3,22} , <i>Ochroma lagopus</i> ^{3,22} , <i>Pseudobombax septenarium</i> ^{3,22}	
Bignoniaceae	<i>Crescentia alata</i> ^{3,23}		
Leguminosae			
Caesalpinioidea	<i>Hymenocourbari</i> ^{3,23}		
Mimosoideae	<i>Inga vera</i> ^{3,24}		
Sapotaceae	<i>Manilkara chicle</i> ^{3,25}		
Hummingbirds			
Rubiaceae	<i>Hamelia patens</i> ^{4,58}		
Simaroubaceae	<i>Quassia amara</i> ^{3,27}		
Hawkmoths			
Apocynaceae		<i>Plumeria rubra</i> ^{5,57}	
Bombacaceae		<i>B. quinata</i> ^{5,22}	
Leguminosae			
Mimosoideae			
Rubiaceae	<i>Enterolobium cyclocarpum</i> ^{7,28} , <i>Inga vera</i> ^{3,24} , <i>Pithecolobium saman</i> ^{5,5}	<i>Inga A. guachepel</i> ⁵ , <i>Pithecolobium longifolium</i> ^{5,57} (water?)	
Tiliaceae	<i>Guetaria macroperma</i> ^{5,54}	<i>Conlana hexandra</i> ^{5,59} <i>Luhea candida</i> ⁵ , <i>L. speciosa</i> ⁵	

Table A4-2. Tree species of Hacienda La Pacifica, Cañas, Guanacaste, Costa Rica, classified by pollinator type and diaspore dispersal vector (continued)

	Biotic	Diaspore dispersal vector? Abiotic or no apparent dispersal mechanism	No data
Pollen vector¹ Medium-to-large bees			
Apocynaceae	<i>Stemmadenia obovata</i> ³⁰ , <i>Thevetia ovala</i> ³⁵		
Bignoniaceae		<i>Godmania aciculifolia</i> ³¹⁸ , <i>Tabebuia impetiginosa</i> ³²² , <i>T. ochracea</i> ³²² , <i>T. rosea</i> ³²²	
Cochlospermaceae	<i>Curatella americana</i> ³²⁴	<i>Cochlospermum vitifolium</i> ³³³	
Dilleniaceae			
Leguminosae			
Caesalpinoidea	<i>Cassia grandis</i> ³³⁵ , <i>Swerthia cubensis</i> ⁴⁵⁹	<i>Caesalpinia eriostachys</i> ³⁶⁵ , <i>Cassia emarginata</i> ³ <i>Schizolobium parathybium</i> ³⁵⁸	
Papilionoideae	<i>Anidra imermis</i> ³³	<i>Dalbergia retusa</i> ³⁶⁶ , <i>Lonchocarpus costaricensis</i> ³⁵⁷ , <i>L. eriocarinata</i> ³⁵⁵ , <i>Platymiscium pleiostachyum</i> ³⁷¹ , <i>Pterocarpus rubri</i> ³⁵ , <i>Myrsopernum frutescens</i> ³⁵ , <i>Piscidia carthagenensis</i> ³⁵	
Malpighiaceae	<i>Byrsonima crassifolia</i> ³³⁸ , <i>Malpighia glabra</i> ^{30,39}		

Table A4-2. Tree species of Hacienda La Pacífica, Cañas, Guanacaste, Costa Rica, classified by pollinator type and diaspore dispersal vector (continued)

Pollen vector ¹ Medium-large bees	Biotic	Diaspore dispersal vector ² Abiotic or no apparent dispersal mechanism	No data
Rubiaceae Sytracaceae Tiliaceae Small moths, diverse insects Anacardiaceae	<i>Gemipa americana</i> ¹⁵ <i>Sytrax argenteus</i> ^{1,40} <i>Ardisia thiborhou</i> ^{1,21} <i>Anacardium excelsum</i> ^{1,2,3} , <i>Spondias</i> <i>mombin</i> ^{1,4,3} <i>Astronium graveolens</i> ^{1,3,70} <i>Saidodendron excelsum</i> ^{1,3,42} <i>Cordia bicolor</i> ^{1,3,73} , <i>C. colococa</i> ^{1,3,74} <i>B. simarubd</i> ^{1,5,43} , <i>B. tomentosa</i> ^{1,6,26} <i>Meylenus segoianum</i> ^{4,4,44} <i>Licania arborea</i> ^{1,5,5} <i>Dyosyros nicaraguensis</i> ^{1,3,28} <i>Muntingia calabura</i> ^{1,3,5} , <i>Sloanea</i> <i>teniflora</i> ^{1,3,45} <i>Casearia aculeata</i> ^{1,5} , <i>C. corymbosa</i> ^{1,3,46} , <i>C. tremula</i> ^{1,3,5} , <i>P. crucei</i> ^{1,3,47} <i>Ocotea veraguensis</i> ^{1,3,48}	<i>Cordia alliodora</i> ^{1,15}	
Flacourtiaceae Lauraceae			

Table A4-2. Tree species of Hacienda La Pacifica, Cañas, Guanacaste, Costa Rica, classified by pollinator type and diaspore dispersal vector (continued)

Pollen vector ¹	Diaspore dispersal vector ²		
	Biotic	Abiotic or no apparent dispersal mechanism	No data
Small bees, moths, small diverse insects			
Sapindaceae	<i>Allophylus occidentalis</i> ¹³²⁶	<i>Thouinidium decandrum</i> ¹³⁵	
Sapotaceae	<i>Mastichodendron capiri</i> ¹³²⁶		
Simaroubaceae	<i>Picramnia glauca</i> ⁴²⁶	<i>Simarouba alvaradae amorphoides</i> ¹³⁵	
	<i>Gueyuma ulmifolia</i> ¹³⁵⁴ , <i>Trema micrantha</i> ¹³⁵⁵		
Sterculiaceae			
Ulmaceae			
Verbenaceae			
Beetles			
Annonaceae	<i>Sapranthus palanga</i> ²⁰²⁰ , <i>purpurea</i> ²¹²⁸ , <i>A. reticulata</i> ²¹²⁸	<i>Annona</i>	

Table A4-2. Tree species of Hacienda La Pacifica, Cañas, Guanacaste, Costa Rica, classified by pollinator type and diaspore dispersal vector (continued)

	Biotic	Diaspore dispersal vector ² Abiotic or no apparent dispersal mechanism	No data
Pollen vector¹			
Wind			
Moraceae	<i>Cecropia peltata</i> ^{3,23} , <i>Trophis racemosa</i> ^{2,26}		
Fig wasps	<i>Ficus elastica</i> , <i>F. hondurensis</i> , <i>F. insipida</i> , <i>F. ovalis</i>		
Moraceae			
Not classified	<i>Capparis frondosa</i> ²⁵ , <i>C. incana</i> ²⁵ , <i>Sterculia apetala</i> ²⁸	<i>Albizia caribaea</i> , <i>A. adinocephala</i> ; <i>Exostemma mexicanum</i> , <i>Hura bavanense</i> , <i>orepitan</i> , <i>Lysiloma divaricatum</i> ²² ; <i>L. desmostiachyis</i> ⁵ <i>Rebdera trinervis</i> ²⁶	<i>Diphyssa robinoides</i> , <i>Egythroxylon</i> <i>nicaraguensis</i> , <i>Croton niveus</i> , <i>Willardia schiedana</i>

¹Source: see first footnote for each entry; ²source: see second footnote for each entry; ³Heinhaus *et al.*, 1975; ⁴Bawa *et al.*, 1985b; ⁵Janzen and Liesner, 1980; ⁶Häber and Frankie, 1989; ⁷Apóstol *et al.*, 1997; ⁸small, diverse insects (Bawa *et al.*, 1985b); ⁹large, yellow flowers; ¹⁰moths, based on congeners listed by Bawa *et al.*, 1985b; ¹¹moths (Bawa *et al.*, 1985b); ¹²Ghazoul and McLetchy, 2001; ¹³small flowers; ¹⁴Nelson and Hamrick 21beetle pollinated genus; see congeners in Bawa *et al.*, 1985b; ¹⁵wind (kapok floaters); ¹⁶Janzen 1983b; ¹⁷Kopur, 1983; ¹⁸Janzen, 1983c; ¹⁹fleshy fruit (Janzen and Liesner, 1980); ²⁰fleshy drupe (Standley and Steyermark, 1946a); ²¹Janzen and Martin, 1982; ²²winged seed (Witsberger *et al.*, 1982; ²³Fisher and McDermid, 1983; ²⁴Gentry, 1983; ²⁵Bawa and Frankie, 1983; ²⁶birds (Janzen, 1967; ²⁷Salas Estrada, 1993); ²⁸molluscs² in pod; ²⁹dry, deliquescent pod (Janzen and Liesner, 1980); ³⁰wind-dispersal appears to be generalized in the genus, e.g. congeners listed in Augsberger (1986), Ibarra-Muniqués (2001). See also Molina, 1996; ³¹Anderson, 1983; ³²birds (Anderson, 1983); ³³Carnacho and Orozco, 1998; ³⁴Janzen 1983d; ³⁵fleshy fruit (Engquist and Sullivan, 2001); ³⁶Elizondo, 1999; ³⁷Janzen 1983d; ³⁸fleshy fruit (Engquist and Sullivan, 2001); ³⁹Stevens, 1983; ⁴⁰Engquist and Sullivan, 2001; ⁴¹Nelson and Hamrick (Engquist and Sullivan, 2001); ⁴²Stevens, 1983; ⁴³Engquist and Sullivan, 2001; ⁴⁴Elizondo, 1999; ⁴⁵Janzen 1983d; ⁴⁶Janzen, 1967; ⁴⁷Stevens, 1983; ⁴⁸Engquist and Sullivan, 2001); ⁴⁹Janzen, 2001); ⁵⁰Janzen and Steyermark, 1946a; ⁵¹Engquist and Sullivan, 2001; ⁵²Foster, 1983; ⁵³Janzen, 1983c; ⁵⁴Janzen, 1967; ⁵⁵Janzen, 1983c; ⁵⁶Hartshorn, 1983; ⁵⁷wind (Molina, 1996); ⁵⁸Janzen, 1967; ⁵⁹Janzen, 1983c; ⁶⁰oil flower² (Gottberger, 1986); ⁶¹beccate fruit (Standley and Steyermark, 1946a); ⁶²wind (Janzen, 1967); ⁶³Janzen, 1967; ⁶⁴fleshy fruit (Engquist and Sullivan, 2001); ⁶⁵explostively dehiscient pod (Bawa and Webb, 1984); ⁶⁶wind (Bawa and Webb, 1984); ⁶⁷Frankie *et al.*, 1974; ⁶⁸juicy berry (Salas Estrada, 1993); ⁶⁹Janzen, 1967; ⁷⁰Janzen and Liesner, 1980); ⁷¹wind (Augsberger, 1986); ⁷²based on similarity to wind dispersed congener *P. demorphantrum* (Witsberger *et al.*, 1982); ⁷³based on similarity with *L. desmostiachyis*; ⁷⁴fleshy seed (Janzen and Liesner, 1980); ⁷⁵wind (Lurie *et al.*, 1974); ⁷⁶pulpy fruit (Standley and Steyermark, 1946b)

Table A4-3. Population densities of tree species of Hacienda La Pacífica, Cañas, Guanacaste, Costa Rica (data based on Glander and Nisbett, 1996; Jiménez et al., 1987)

Species	Density (stems \geq 1cm dbh ha ⁻¹)	
	Riparian ¹	Upland
Anacardiaceae		
<i>Anacardium excelsum</i>	29	0
<i>Astronium graveolens</i>	6.75	8.3
<i>Spondias mombin</i>	4.15	0.2
Annonaceae		
<i>Annona purpurea</i>	1.4	0.6
<i>Annona reticulata</i>	2.35	2.1
<i>Sapranthus palanga</i>	0.2	3.7
Apocynaceae		
<i>Plumeria rubra</i>	0.1	0
<i>Stemmadenia obovata</i>	2.7	6
<i>Tibetia ovata</i>	1.3	0.2
Araliaceae		
<i>Sciadodendron excelsum</i>	0.1	0.2
Bignoniaceae		
<i>Crescentia alata</i>	0	0
<i>Godmania aesculifolia</i>	0	1.4
<i>Tabebuia impetiginosa</i>	0.1	1.9
<i>Tabebuia ochraceae</i>	6.95	52.4
<i>Tabebuia rosea</i>	4.35	0.4
Bombacaceae		
<i>Bombacopsis quinata</i>	1	1.7
<i>Ceiba aesculifolia</i>	0	0
<i>Ceiba pentandra</i>	0.8	0
<i>Ochroma lagopus</i>	0.1	0
<i>Pseudobombax septenatum</i>	0.1	0
Boraginaceae		
<i>Cordia alliodora</i>	9.25	68.4
<i>Cordia bicolor</i>	1.6	9.1
<i>Cordia colococca</i>	7.7	28.9
Burseraceae		
<i>Bursera simaruba</i>	2.5	1.2
<i>Bursera tomentosa</i>	0	0
Caesalpinaceae		
<i>Caesalpinia eriostachys</i>	0	0
<i>Cassia emarginata</i>	0.4	5
<i>Cassia grandis</i>	0.9	0
<i>Hymenaea courbaril</i>	7.4	1.9
<i>Schizolobium parahybum</i>	1.1	0
<i>Swartzia cubensis</i>	0	0

Table A4-3. Population densities of tree species of Hacienda La Pacífica, Cañas, Guanacaste, Costa Rica (data based on Glander and Nisbett, 1996; Jiménez et al., 1987) (continued)

Species	Density (stems \geq 1cm dbh ha ⁻¹)	
	Riparian ¹	Upland
Capparidaceae		
<i>Capparis frondosa</i>	0	0
<i>Capparis incana</i>	0.1	0
Celastraceae		
<i>Maytenus segoiicarum</i>	0	0
Chrysobalanaceae		
<i>Licania arborea</i>	5	0.4
Cochlospermaceae		
<i>Cochlospermum vitifolium</i>	1	1
Dilleniaceae		
<i>Coccoloba americana</i>	0	0
Ebenaceae		
<i>Dyospiros nicaraguensis</i>	7.75	8.9
Eleocarpaceae		
<i>Muntingia calabura</i>	3.6	0.8
<i>Sloanea terniflora</i>	2.6	0.2
Erythroxylaceae		
<i>Erythroxylon bavanense</i>	0	0
Euphorbiaceae		
<i>Bernardia nicaraguensis</i>	0	0
<i>Croton niveus</i>	0	0
<i>Hura crepitans</i>	0.2	0
Flacourtiaceae		
<i>Casearia aculeata</i>	2.3	8.5
<i>Casearia corymbosa</i>	2.5	11.6
<i>Casearia tremula</i>	0.1	0
<i>Prockia crucis</i>	0	0
Lauraceae		
<i>Ocotea veraguensis</i>	0	0
Malpighiaceae		
<i>Byrsonima crassifolia</i>	0.5	0.2
<i>Malpighia glabra</i>	0.2	0
Meliaceae		
<i>Cedrela odorata</i>	1	0
<i>Guarea excelsa</i>	0	0
<i>Swietenia macrophylla</i>	12.05	5.6
<i>Trichilia birta</i>	0	0
<i>Trichilia americana</i>	3.9	12.3
<i>Trichilia martiana</i>	1.95	0.2

Table A4-3. Population densities of tree species of Hacienda La Pacífica, Cañas, Guanacaste, Costa Rica (data based on Glander and Nisbett, 1996; Jiménez *et al.*, 1987) (continued)

Species	Density (stems \geq 1cm dbh ha ⁻¹)	
	Riparian ¹	Upland
Menispermaceae		
<i>Hyberbaena tonduzii</i>	0	0
Mimisoideae		
<i>Acacia farnesiana</i>	0	0
<i>Albizia adinocephala</i>	0.3	3.7
<i>Albizia caribaea</i>	1.7	25.8
<i>Albizia guachepete</i>	4.7	1.9
<i>Enterolobium cyclocarpum</i>	4.6	3.1
<i>Inga vera</i>	1.4	0
<i>Lysiloma desmostachys</i>	0	0
<i>Lysiloma divaricatum</i>	3.45	29.1
<i>Pithecellobium longifolium</i>	4.65	0
<i>Pithecellobium saman</i>	2.5	0.6
Moraceae		
<i>Brosimum alicastrum</i>	0	0
<i>Cecropia peltata</i>	0.2	0
<i>Chlorophora tinctoria</i>	1.5	3.3
<i>Ficus hondurensis</i>	0	0
<i>Ficus elastica</i>	0	0
<i>Ficus insipida</i>	0	0
<i>Ficus ovalis</i>	0.5	0.42
<i>Trophis racemosa</i>	0	0
Myrsinaceae		
<i>Ardisia revoluta</i>	4.8	0
Papilionoideae		
<i>Acosmium panamense</i>	0.4	0.6
<i>Andira inermis</i>	2.65	0
<i>Dalbergia retusa</i>	4.3	4
<i>Diphysa robinoides</i>	0	0
<i>Lonchocarpus costaricensis</i>	0	0.4
<i>Lonchocarpus eriocarinalis</i>	0	0
<i>Lonchocarpus minimiflorus</i>	31.6	546
<i>Machaerium biovulatum</i>	2.3	11
<i>Myrospermum frutescens</i>	11.75	19.3
<i>Piscidia carthagenesis</i>	4.2	6
<i>Platimiscium pleostachyum</i>	0.1	0.8
<i>Pterocarpus rohrii</i>	0	6.2
<i>Willardia schiedeana</i>	0	0

Table A4-3. Population densities of tree species of Hacienda La Pacífica, Cañas, Guanacaste, Costa Rica (data based on Glander and Nisbett, 1996; Jiménez *et al.*, 1987) (continued)

Species	Density (stems \geq 1cm dbh ha ⁻¹)	
	Riparian ¹	Upland
Polygonaceae		
<i>Coccoloba caracasana</i>	2.3	0
Rubiaceae		
<i>Calycophyllum candidissimum</i>	4.35	3.7
<i>Conzattia hexandra</i>	0	0
<i>Excostema mexicanum</i>	0	0
<i>Genipa americana</i>	3.45	2.7
<i>Guettarda macrosperma</i>	0	0
<i>Hamelia patens</i>	0	0
Rutaceae		
<i>Esenbeckia littoralis</i>	0	0
Sapindaceae		
<i>Allophylus occidentalis</i>	1.3	4.1
<i>Thouinidium decandrum</i>	4.15	7.3
Sapotaceae		
<i>Manilkara chicle</i>	5.15	0
<i>Mastichodendron capiri</i>	0.6	1.2
Simaroubaceae		
<i>Alvaradoa amorphoides</i>	0	0
<i>Picramnia latifolia</i>	0	0
<i>Quassia amara</i>	0.2	0
<i>Simarouba glauca</i>	2.75	0.8
Sterculiaceae		
<i>Guazuma tomentosa</i>	60.35	68
<i>Stercula apetala</i>	0.7	0.2
Styracaceae		
<i>Styrax argenteus</i>	0	0
Tiliaceae		
<i>Apeiba tiborbou</i>	0.4	0
<i>Luehea candida</i>	8.8	23.7
<i>Luehea speciosa</i>	0.1	0.4
Ulmaceae		
<i>Trema micrantha</i>	0.6	0.6
Verbenaceae		
<i>Rehdera trinervis</i>	0.305	0
Mean	2.81	
Median	0.50	

¹Mean of Glander and Nisbett's two riparian areas

APPENDIX V: FINANCIAL REPORTS

FINANCIAL REPORT

FOR THE PERIOD ENDED DECEMBER 31, 2001

SOURCE : CENTER FOR INTERNATIONAL FORESTRY RESEARCH (CIFOR) CG (SWGRP)

DONOR : CIFOR

**AGREEMENT : THE IMPACT OF HABITAT FRAGMENTATION ON GENETIC
DIVERSITY, MATING SYSTEMS, AND EFFECTIVE POPULATION
SIZE OF TROPICAL FOREST FRAGMENTS IN COSTA RICA**

**TROPICAL AGRICULTURE RESEARCH AND HIGHER EDUCATION CENTER
CATIE**

TROPICAL AGRICULTURE RESEARCH AND HIGHER EDUCATION CENTER**CATIE****CENTER FOR INTERNATIONAL FORESTRY RESEARCH****CIFOR****FINANCIAL REPORT****PROJECT****THE IMPACT OF HABITAT FRAGMENTATION ON GENETIC DIVERSITY
MATING SYSTEMS, AND EFFECTIVE POPULATION SIZE OF TROPICAL
FOREST FRAGMENTS IN COSTA RICA****Amount US\$**

Timing	Expenditures 01/04/96 to 08/31/1997	Expenditures 09/01/97 to 09/30/1998	Expenditures 10/01/98 to 12/31/1999	Expenditures 01/01/00 to 12/31/2001	Total Expenditures
Emoluments					
Salary, Project coordinator	56.287,73	50.596,46	11.848,06	-	118.732,25
Tree Climber		2.308,79	1.114,58	190,99	3.614,36
Equipment etc.					-
Vehicle	-	7.071,84		-	7.071,84
Lab. Supplies	-	355,00	19,68	-	374,68
Communications	-	83,54	807,88	387,60	1.279,02
Fuel	-	318,89	18,45	3,92	341,26
Vehicle maintenance	-	584,30	11,21	-	595,51
					-
Travel expenses					-
Local travel	7.812,73	4.389,67	77,33	280,40	12.560,13
International travel	2.300,62	-	1.173,78	110,00	3.584,40
				-	-
Training	3.881,44	3.745,50	2.436,21	5.926,19	15.989,34
					-
Miscellaneous					-
Vehicle insurance	1.132,74	2.180,48	2.121,79	416,01	5.851,02
Seed Collection			198,90	-	198,90
Overhead	-	5.959,20		1.345,41	7.304,61
TOTAL	71.415,26	77.593,67	19.827,87	8.660,52	177.497,32

**TROPICAL AGRICULTURE RESEARCH AND HIGHER EDUCATION CENTER
CATIE**

REMITTANCES RECEIVED

<u>DATE</u>	<u>US Dollars</u>
May 28, 1996	40.000,00
May 28, 1997	30.000,00
June 19, 1997	30.000,00
February 02, 1998	45.000,00
May 06, 1998	30.000,00
February 10, 1999	5.000,00
Total	<u><u>180.000,00</u></u>



Note:


Funds may be deposited in a/c No. 20.8619.1616
CATIE-REGULAR, BANK OF AMERICA
730 15th Street, N.W., 7th Floor
Washington, D.C. 20005-1012 USA.
A.B.A 054001204

**TROPICAL AGRICULTURE RESEARCH AND HIGHER EDUCATION CENTER
CATIE**

FINANCIAL POSITION

	<u>US Dollars</u>
Payment Received	<u>180.000,00</u>
Less: Recorded Expenses	<u>(177.497,32)</u>
Balance	<u><u>2.502,68</u></u>

I hereby certify that the information contained herein is true and correct to be the best of my knowledge


Eduardo Piccolo
Director Administration and Finances


Jonathan Cornelius
Project Coordinator



FINANCIAL REPORT

FOR THE PERIOD ENDED DICEMBER 31, 1999

SOURCE : CENTRE FOR INTERNATIONAL FORESTRY RESEARCH

DONOR : CIFOR

**AGREEMENT : THE ROLE OF REMOTE SENSING IN MONITORING TROPICAL FOREST
FRAGMENTS IN COSTA RICA (AERIAL PHOTOGRAPHY)**

**TROPICAL AGRICULTURE RESEARCH AND HIGHER EDUCATION CENTRE
CATIE**

**TROPICAL AGRICULTURE RESEARCH AND HIGHER EDUCATION CENTER
CATIE**

**PROJECT
CENTRE FOR INTERNATIONAL FORESTRY RESEARCH
CIFOR**

FINANCIAL REPORT

**AERIAL PHOTOGRAPHY
Amount US \$**

ACTIVITY	TOTAL EXPENDITURES
National Travels	198,94
Fuel	25,22
Equipment	3.874,44
Aereal photography	10.881,40
Total	14.980,00



Account Code

4470

REMITTANCES RECEIVED

DATE	US\$
December 31, 1997	12.000,00
September 27, 1999	<u>2.980,00</u>
Total	<u><u>14.980,00</u></u>



Account Code

4470

FINACIAL POSITION

	US\$
Payment Received	14,980.00
Recorded expenses	<u>14,980.00</u>
Balance	<u><u>0.00</u></u>

I hereby certify that the information contained herein is true and correct
to be the best of my knowledge


Viviana Sanchez Chaves
Director Administration and Finances


Jonathan Cornelius
Project Coordinator



APPENDIX VI: PROJECT PERSONNEL

Researcher: Jonathan Cornelius

Research assistant: Oldemar Baeza

Field worker: Manuel Sojo

Volunteer: Sofia Ryder