

LEGUMINOUS TREES FOR SHADE

✓
**G. Budowski
D.C.L. Kass
R.O. Russo**

**Paper presented at Symposium on Nitrogen Fixing trees for
the tropics National Research Program on Soil Biology
(PNPBS) EMBRAPA - UFRRJ, Km 47, Rio de Janeiro, Brasil
19 - 24 September, 1983.**

**The publication and distribution of this work was sponsored by the
Swiss Program of Cooperation for Development, DDA, through INFORAT:
Información y Documentación Forestal para América Tropical.**

**CENTRO AGRONOMOICO TROPICAL DE INVESTIGACION Y ENSEÑANZA, CATIE
Department of Natural Renewable Resources
Turrialba, Costa Rica, 1983**

LEGUMINOUS TREES FOR SHADE

G. Budowski^{1/}
D.C.L. Kass^{1/}
R.O. Russo^{1/}

Centro Interamericano
Documentación e Información
Agrícola

16 OCT 1974

C I D I A
Turrialba, Costa Rica

ABSTRACT

Use of leguminous trees to provide shade, especially in perennial crops, would appear to be an ancient agricultural practice, probably coinciding with the domestication of perennials such as tea, coffee, and cocoa. Leguminous trees can also be found in association with annual crops, in pastures, and in situations such as living fenceposts, where shade is not clearly being provided to an associated species. The factors which might influence the choice of leguminous as opposed to non-leguminous trees to provide shade in such situations are considered. Various such associations are described in some detail, in an attempt to elucidate some of the ecological interrelationships that are probably reflected in traditional agroforestry practices. It is concluded that not only the possibility of improving nitrogen nutrition but also other characteristics such as type of shade, coppicing ability, and ease of husbandry might have favored the selection of leguminous trees for shade. Many of these areas would appear to merit considerably more research with the objective of obtaining more quantitative data than is presently available.

^{1/}Head of Department of Renewable Natural Resources, Soil Management Specialist and Forester respectively, Centro Agronómico Tropical de Investigación y Enseñanza, CATIE, Turrialba, Costa Rica.

LEGUMINOUS TREES FOR SHADE^{1/}

Gerardo Budowski^{2/}
Donald C.L. Kass^{2/}
Ricardo O. Russo^{2/}

I. INTRODUCTION

A shade tree could be defined as one used to reduce radiation (light and heat) to some part of the environment. Due to the fact that a tree rather than some inanimate object is used to produce shade and that the environment usually includes growing plants and animals, a complex interaction is set up between the environment, the shade trees, and the shaded species which extends far beyond the mere reduction of heat and light (Willey, 1975).

We shall first briefly describe the effects of shade per se and then proceed to a discussion of how the situation is altered when a tree is used to produce the shade. Finally, we shall consider the special case which is the object of the present discussion-- the effect of using a leguminous tree for shade.

II. THE EFFECTS OF SHADE

Considering first the environment when no living species is associated with the shade, the primary result of the shade will be to reduce soil and ambient temperatures. Reduced temperature will reduce evaporation from the

^{1/} Contribution from the Departments of Renewable Natural Resources and Crop Production, CATIE (Tropical Agricultural Research and Training Center), Turrialba, Costa Rica.

^{2/} Head of Department of Renewable Natural Resources; Soil Management Specialist, Crop Production Department; and Forest Management Specialist, Department of Renewable Natural Resources, respectively. CATIE, Turrialba, Costa Rica.

surface and increase the relative humidity of the environment. Reduced temperatures will reduce the reaction rates of physical and biological processes at or below the soils surface (Willey, 1975).

When plants or animals are placed in a shaded environment, the effects of reduced temperature become more complex. Reduced temperatures will reduce stress for animals in hot climates, especially those unadapted to such climates. They will also probably induce a modification in activities. However, the increased relative humidity brought on by reduced temperatures may have adverse effects.

Shading will obviously reduce the light available to plants. The effect of shade on plant growth has been the object of numerous experiments, reviewing which Grime (1979) reached the following conclusions:

1. In response to shade, the majority of plants produce less dry matter, retain photosynthate in the shoot at the expense of root growth, develop longer internodes and petioles, and produce larger thinner leaves.
2. Species differ considerably both with respect to the magnitude and rate of these responses. However, the capacity to maximize dry matter production in shade through modification of the phenotype is most apparent in species characteristic of unshaded or lightly shaded environments while plants normally found in deep shade tend to grow slowly and to show much less pronounced morphogenic responses to shade treatment (Grime, 1979).

Economic plants most commonly grown under shade are coffee (Coffea arbica L.), tea (Camellia sinensis L.), cocoa (Theobroma cacao L.), black pepper (Piper nigrum L.), vanilla (Vanilla planifolia L.), the species of the Zingiberaceae (ginger, cardamom, and turmeric) as well as tobacco (Nicotiana tabacum L.) for cigar wrappers. As expected, in the wild state, all of these species with the exception of tobacco, occur as understory plants in a tropical forest environment (Purseglove, 1968). In tobacco, the production are large, thin leaves as a consequence of shade is a desirable characteristic for cigar wrappers (Willey, 1975). Willey (1975) reviewed several investigations in which coffee and cocoa leaves were to some degree able to compensate for the reduced light produced by shading by reflecting less light, increasing the chlorophyll content, reorienting the chloroplasts to give less transmission and better utilization of incident light, and increasing the number of stomata per unit of leaf surface. Reduced light has been shown to reduce the intensity, if not the duration, of "flushing" in cocoa. Changes in the chemical composition of leaves due to shading is of special importance in tea, where the leaf is the harvested product. Recent work has shown that shade might adversely affect tea quality (Hilton, 1974).

While most researchers agree that reduction of light intensity by shade reduces photosynthesis and probably yields, reduction in leaf temperature and especially reduction in the diurnal variation of leaf temperature caused by shading is generally accepted as beneficial in cocoa, coffee and tea (Willey, 1975). The benefits of reduced soil temperature may be of additional importance at the seeding or establishment stage but no experimental

evidence was found by Willey (1975). Increased relative humidity occasioned by reduced temperatures appears to improve water balances (Fordham, 1971) by reducing transpiration.

Before considering the difference in the situation where the shade is produced by trees, two consequences of reduced light intensity should be mentioned. Firstly plants that grow less due to reduced photosynthesis will yield less but will also take up smaller amounts of essential nutrients. At low nutrient levels, shaded cocoa (Murray and Nichols, 1966) and tea (Wight, 1958) may even outyield nonshaded plants. Other workers have found higher yields for non-shaded cocoa even without fertilizer (Willey, 1975) but this will depend on the fertility level of the soil involved. In any case, shading can keep the yield level down to a level that the fertility status of the soil can maintain (Hardy, 1962). In coffee, shading also equalizes yields over years of overbearing and subsequent dieback (Huxley, 1970). Thus, for farmers without access to high levels of mineral fertilizers, shade can be of considerable value (Beer, 1982).

Secondly, shade will alter the ecological balance between the crop and its associated pests, weeds, and diseases (Willey, 1975). In some cases, this alteration has been shown to be of benefit to the crop as in the case of the cocoa capsid the coffee leaf miner, and various weeds in pastures. In other cases, shading may favor certain pests and diseases.

III. TREES FOR SHADE

When shade is provided by trees, the selection becomes complex, mostly because one is dealing with a plant that is photosynthesizing and transpiring rather than merely providing shade. Since they are photosynthesizing, shade trees will affect the quality as well as the quantity of transmitted light. In transmitted light, i.e. that which actually passes through leaves of the shade tree, most of the wavelengths used in photosynthesis are filtered out by the chloroplasts of the leaves of the shade species (Willey, 1975). Thus, the understory crop will depend to a great degree on the light which passes through the shade tree leaf canopy without being used for photosynthesis. The distinction often made in the literature between light and heavy shading thus takes on considerable significance although it is rarely specified as whether "light shading" means a fairly uniform sparse leaf canopy or a small number of widely spaced but very large shade trees (Willey, 1975). The leaf architecture (size and dispersion) of the shade tree should have a great effect on how much radiation is transmitted with loss of quality and how much is sunflecks, i.e. light which comes through the shade with little or no reduction in quality. Other complicating factors will be wind speed, cumulus clouds, and the effect on photoperiod of the light of altered quality (Allen, Sinclair, and Lemon, 1976). It would seem that a so-called "dappled" shade, with abundant sunflecks, which change with movement of the shade tree leaves would be of greatest benefit to the shaded crop since the other benefits of shading would be maintained (Willey, 1975).

Whether transpiration by the shade tree increases water stress to the associated species is one of the most controversial aspects of the use of shade trees (Willey, 1975). Again, the effect cannot readily be separated from the reduced evaporation and transpiration in the associated crop induced by the shade. In very humid environments, such as those frequently used for the production of cocoa and tea, the transpiration of the shade tree is probably of little effect and the transpiration of the shade tree may even remove excess moisture in low-lying, high water table areas (Cadima and Alvim, 1967). When moisture is more limiting, there are frequent reports of shade trees increasing moisture stress (Franco and Inforzato, 1951). Willey (1975) puts considerable emphasis on the mass and distribution of roots in the shade and shaded species. Deciduous shade species or pruning shade trees in the dry season would reduce moisture loss in the dry season, but the other benefits of shading would also be lost in this period.

Evaluation of the effect of shade trees of nutrient availability is complicated by the fact, pointed out earlier, that shade reduces growth and nutrient requirements of the associated species. The benefits of the abundant (up to $5000 \text{ kg ha}^{-1}/\text{yr}^{-1}$) of leaf litter generally produced by most shade species is generally recognized although Willey (1975) makes some qualifications to these advantages:

1. The benefits to soil temperature and erosion control of the litter are less critical in an environment which is already shaded.

2. Except for the case where the shade trees bring up nutrients from lower levels or fix nitrogen, the leaf litter merely recycles nutrients which theoretically should have been available to the associated species, if over a longer time period.
3. Most of these benefits could also be obtained by mulching.

It would appear to these authors that these qualifications have more theoretical than practical significance. The nutrients in the litter are much more readily available to the associated crop than dispersed in the soil where they can be readily lost by leaching. Mulching requires the added expense of transporting the mulch material and in any case, the effects of shading are lost.

Some other effects of shade trees which were enumerated by a series of papers by Budowski (1959, 1981a, 1981b, 1981c, 1982) and Beer (1982) include:

1. Damage to the understory crop by falling or pruned branches of the tree shade (Purseglove, 1968).
2. Effect of shade trees on wind velocities, reducing evaporation from the leaf surfaces of the shaded species, but also perhaps promoting condensation which may be of value in dry periods.
3. The shade trees may produce an economic product, which might result in spreading labor use over a longer period.
4. Reduction of damage due to hail and heavy rain.

5. Concentration of raindrops, increasing erosion hazard, which should be offset to some degree by the effect of trees on reducing raindrop impact.
6. Difficulties in harvest, especially mechanized harvest, occasioned by the presence of a shade tree in the associated crop.
7. Use of shade trees results in a more complex environment, more difficult to study and manage efficiently (Budowski, 1981b).

IV. LEGUMINOUS TREES FOR SHADE

In considering the special case, in which shade is provided by a member of the Leguminosae, four different uses of shade will be discussed separately, depending upon the type of fauna and flora associated. Some of the considerations discussed under one use will, in many cases, also be of relevance for another use.

The categories we have chosen are as follows:

- a. Shade for perennial crops
- b. Shade for animals and/or pasture species
- c. Association with annual crops, including alley cropping
- d. Situations in which there is no obvious associated species.

These would include the provision of shade as an amenity, and the use of leguminous trees for windbreaks, firebreaks and living fences.

a. Shade for perennial crops

Most of the tree species used for shade in perennial crops are legumes. All but one of the species listed by Purseglove as shade for tea are legumes. Of the six genera listed for cacao shade, four are legumes. Although only four of the nine genera listed as permanent coffee shade are legumes, most of these are used in East Africa, while in the Americas, leguminous shade species predominate. All of the five genera, Purseglove lists as temporary shades are legumes. In the case of black pepper, it is most common to plant the vines in existing plantations of other economic species; where shade trees are planted especially for the pepper, leguminous trees are generally used. Where pepper production has been introduced by oriental immigrants as in Malaysia and Brazil, more intensive, unshaded production has predominated resulting in higher yields, but also rapid soil degradation in Malaysia (Purseglove, 1968).

When confronted with such evidence, the average agronomist would conclude that leguminous trees have been chosen for their nitrogen fixing ability. It would probably be wrong however to conclude that this was the only, if indeed the major, consideration used in choosing these trees by ancient farmers in widely separated parts of the globe.

Certainly, these farmers could not have been aware of the nitrogen-fixing ability of these trees since even now, evidence for such fixation is poorly documented and mostly based on circumstantial evidence (Nair, 1982; Orchard and Darb, 1956; Salinas and Sanchez, 1981; Pak et al., 1977); Felker, 1978; Radwanski, and Wickens, 1969; Enriquez, 1983), the exception being

actual measurement of acetylene reduction by Inga jinicuil in association with coffee (Roskoski, 1982). Leguminous trees were probably an important constituent of the forests in which cocoa, coffee, cinnamon, and tea occur naturally. In selective cutting of these forests, statistical considerations might account for a large proportion of leguminous trees being left for shade even if no selection were made. Another argument against nitrogen-fixation as a criterion for selection of leguminous trees for shade is the low nitrogen requirement of crops such as coffee, cocoa, tea (Sanchez, 1976) and cinnamon, especially when grown under shade (Willey, 1975). Furthermore coffee and tea were domesticated and associated with leguminous trees at high elevations in the tropics where soils of volcanic origin often have high organic matter contents and nitrogen is unlikely to be a limiting element. Even at lower elevations in the tropics, soils of volcanic origins show considerable N-supplying power (Kass, Quinlan, and Russo, 1983).

Evidence for benefits of leguminous as opposed to non-leguminous trees for shade of cocoa was obtained by Enriquez (1983) in Costa Rica and presented in Table 1. It is impossible to tell, of course, whether the increase is due to nitrogen supplied by the leguminous trees or due to one of the other factors enumerated above.

Roskoski (1981, 1982) has shown nitrogen fixation rates of 35-40 kg⁻¹ ha⁻¹ yr⁻¹ by Inga jinicuil Schlechter as measured by acetylene reduction when used as a coffee shade in Mexico. No evidence of fixation was found for Inga vera H.B. and K. while coffee yields associated with the Inga jinicuil were 37% higher than those associated with I. vera. This study also pointed

out that not all leguminous trees bear nodules and fix nitrogen. It would thus be difficult to explain the widespread use of Inga vera as a coffee shade on the basis of its supposed N-fixing ability.

Table 1. Production and pod index of Cocoa in four years with shade of Cordia alliodora and Erythrina poeppigiana (means of 3 varieties of cocoa)

System	Year	Cocoa Yield (kg/ha ⁻¹)	Pod index (pods per kg of dry cocoa)
Cocoa + <u>Cordia</u>	1979	83.25	18.2
	1980	468.73	23.4
	1981	371.50	24.5
	1982	708.36	25.0
	Mean	407.96	22.8
Cocoa + <u>Erythrina</u>	1979	162.75	20.5
	1980	631.70	20.9
	1981	979.11	18.1
	1982	924.43	32.1
	Mean	674.50	20.7

From Enriquez, 1983

There are six reasons, other than nitrogen fixation that legumes might be favored for use as shade trees in perennial crops. These includes

1. Production of valuable products
2. Abundant litter production (which might be related to N fixation)
3. Ease of husbandry. Most leguminous trees reproduce readily from stakes or large cuttings.
4. Good coppicing ability while this might be related to nitrogen fixation, it is belied by the fact that several non-legumes also coppice well.
5. A canopy structure that permits a desirable type of shade.
6. Drought resistance; non-competitive root structure.

The leguminous genera commonly used for shade do not produce any particularly valuable products. Neither Albizia, Erythrina, Gliricidia, or Inga are prized for their timber; and one participant in a CATIE sponsored agroforestry seminar recommended replacing Inga with cedar and citrus in Mexican coffee plantations (Fuentes, 1979). All of these species, with the exception of Erythrina, can be used for fuelwood, however, (National Academy of Sciences, 1980) while Erythrina and Inga produce edible products. Inga pod pulp, Erythrina flowers, and wood from Gliricidia and Albizia might have been more important to the people who first used these trees for shade than they appear today to an economist.

While litter production under leguminous trees used for shade is considerable, not too much data is available for comparison of leguminous and non-leguminous trees for shade. Russo (1983) obtained a figure of 4200 kg/ha/yr of dry matter for leaf fall from *Erythrina poeppigiana* in a coffee plantation. Hadfield (1963) quoted by Willey (1975) gives a figure of 5000 kg/ha/yr for tea shades, which are presumably legumes; for, with the exception of *Grevillea robusta* A. Cunn., use of non-legumes as tea shades is rare (Purseglove, 1968). Nye (1961), and Greenland and Kowal (1960) give somewhat higher figures (10,500 to 12,500 kg ha⁻¹ yr⁻¹) for tropical rainforest but they were probably dealing with higher density of larger trees than would be found in a coffee or tea plantation. A comparison of litter production by *Cordia* and *Erythrina* was made in the experiment of Enriquez (1983) but the data are presently being analyzed as part of a doctoral thesis. As we shall soon see that coppicing ability is an almost universal attribute of leguminous trees, figures for litter production may be more than quadrupled if the biomass of material pruned annually is included (Russo, 1983).

The four genera we have mentioned as the most common coffee, cocoa and tea shades are all fast growing and easy to reproduce (NAS, 1979; NAS, 1980). This attribute might have favored their use as shade trees especially as we have seen that Roskoski (1982) was unable to find nodules or evidence of N fixation in *Inga vera*, a widely used leguminous coffee shade in Mexico and Central America, which is noted for its rapid growth (NAS, 1980).

A fairly good argument can probably be made for coppicing ability as a desirable characteristic that leguminous trees commonly used as shade trees share-- in fact, there would appear to be very few legumes that do not coppice well (NAS, 1980). There are several non-legumes such as Eucalyptus, which do coppice well, but good coppicing ability would seem to be much more common among the legumes. Interestingly, some non-legumes which do not coppice well such as Grevillea robusta, have recently been favored for shade. Perhaps increased labor cost has made pollarding a less common practice and provision of a light shade that can be readily maintained might be more conducive to higher yields at moderate fertilizer levels (Blom, 1980).

This brings us to another attribute of the leguminous trees which might not be related to nitrogen fixation: the provision of a desirable type of shade. The light shade provided by Gleditsia (NAS, 1979), Giricidia (NAS, 1980) and the non-leguminous Grevillea (NAS, 1980) are often mentioned, but the authors were unable to find any data with the exception an M.S. thesis by Daccarett (1967) who measured light interception by three leguminous species and Cordia alliodora. Unfortunately, populations of the trees in the study varied so that the data are only comparable for Erythrina poeppigiana and Gliricidia sepium, two legumes, which do however have differing types of crowns. No evaluation of light quality was made. The data, however, are given in Table 2 as similar data are not readily available.

Table 2. Light interception by varying populations of different species in four different directions from trees (10 observations per direction) (Daccaret, 1967)

Species	Population trees ha ⁻¹	Light interception, 2.5m from tree %				
		north	east	south	west	mean
<u>Erythrina poeppigiana</u>	24	53.83	49.82	56.21	62.57	55.60
<u>Pithecolobium saman</u>	20	9.02	10.37	27.36	28.47	18.80
<u>Gliricidia sepium</u>	24	24.53	25.73	32.56	54.90	34.44
<u>Cordia alliodora</u>	8	0.95	1.14	2.86	19.36	6.08

It would seem to the authors that pinnate leaves of the type so common in the legumes might produce a more favorable type of shade. There would appear to be very little information available on this subject although Norman, Miller and Tanner (1971) compared gap-size and light intensity distributions below canopies of sunflower (Helianthus annuus L.) which has simple leaves and sumac (Rhus typhina L.) which has compound leaves. A type of canopy permitting good light penetration has been cited as a positive attribute of species such as Leucaena (Brewbaker and Hutton, 1978) and Gliricidia sepium (National Academy of Sciences, 1980) but no quantitative data is presented. It would seem to be an area meriting further research.

Rooting habit of leguminous trees is another area of much speculation and little factual data. Again, Leucaena is supposed to have a deep

taproot and few lateral roots, thus favoring its use in alley cropping schemes (Dijkman, 1950; Blom, 1980). Kang, Wilson, and Sipkens (1981) indeed showed that *Leucaena* root mass drops off considerably between 20 and 75 cm from a hedge row but no comparative data for other species was presented. Daccaret (1967) found more roots of *Erythrina poeppigiana* and *Gliricidia sepium* than of the leguminous *Pithecolobium saman* and of the non-leguminous *Cordia alliodora* at a distance of 1 m from the respective trees in a tree-pasture association. For some reason the pasture species had a larger proportion of their roots in the upper 20 cm under *Erythrina* than under any of the other species or in the control. It would appear that there might be intra - as well as inter-family differences with regard to rooting habit among shade species.

In conclusion, it would appear that there is some evidence that nitrogen-fixing ability might not be the only attribute of leguminous trees which has favored their use for shade. There is, however, hardly enough data to draw many conclusions about these attributes because even nitrogen fixation has not been demonstrated in many leguminous species used for shade. Some of the characteristics considered here, such as litter production, pruning management and coppicing ability, amount and type of shade, rooting habit, as well as nitrogen fixation would appear to merit a considerable research effort to see if differences do exist, not only for legumes as a group but also among different leguminous species.

b. Shade for animals and/or pasture species

The practice of letting seedlings from natural forest regeneration thrive in pastures is well known in Costa Rica and other regions (Budowski, 1981a; Lagemann and Heuveldop, 1983). The benefits of shade trees in pastures are considerable; and according to Budowski (1982), it is necessary to consider both biological and sociological aspects to evaluate advantages and disadvantages of silvopastoral associations. The main biological aspects are:

- A better utilization of the vertical space is achieved and to a certain extent, natural ecological models are simulated in regard to form and structure of vegetation.
- There is greater resistance against adverse rainfall conditions (both water excess and deficiency).
- Temperature extremes are mitigated at ground level.
- A large amount of biomass returns to the soil as organic matter through fallen leaves, fruits, flowers and branches.
- Tree roots can reduce compaction and improve infiltration.

The above advantages would apply to both leguminous and non-leguminous species. With leguminous trees, there should be additional advantages such as higher nitrogen content of the litter and foliage which can be used as feed or browse (Blom, 1980; Felker and Bandurski, 1979). Table 3 shows the nutritive value of the foliage of four leguminous tree species

as compared with other forage legumes and grasses commonly used in the humid tropics.

As shade species, leguminous trees have sometimes been reported to have a favorable effect on the grass growth beneath them. According to Daccarett and Blydenstein (1968), the protein content of grasses growing beneath Erythrina poeppigiana was higher than that of grasses growing beneath non-leguminous trees. Currently, at the Centro Agronómico Tropical de Investigación y Enseñanza (CATIE) in Turrialba, Costa Rica, evaluation of production Pennisetum purpureum associated with Erythrina poeppigiana is being carried out (Benavides, 1983a). Another study concerns the grass Cynodon plectostachyus associated with Erythrina poeppigiana, Cordia alliodora, or no tree species (Bronstein, 1983). Some preliminary results are shown in Table 4. Finally, Table 5 shows several leguminous trees commonly found in pastures in Costa Rica and some of their uses.

Mention should also be made of Alnus acuminata, a nitrogen-fixing non-legume commonly found in pastures at the higher elevations in Costa Rica (Beer, 1980). Effects of the trees on pasture production have not been measured; but the frequency with which the association is encountered should imply some benefit.

Table 3. Nutritive values of 4 legume trees, compared with other forages, legumes and grasses.

	C.P. (%)	D.M. %	IVDDM (%)	ME (Mcal/kg DM)
<u>Erythrina poeppigiana</u>	23.2 a 34.4	23.4 a 24.1	56.6	2.00
<u>E. berteriana</u>	24.3	27.8		1.97
<u>Gliricidia sepium</u>	24.8 a 27.6	23.1 a 35.9		
<u>Leucaena leucocephala</u>	25.2 a 35.6	25.9		
<u>Cajanus cajan</u>	24.2	25.0		
<u>Manihot esculenta</u> - leaves	15.8 a 32.1	27.9		1.43
<u>Ipomea batatas</u> - leaves	15.3	16.4		
<u>Dolichos lablab</u>	20.2	20.4		2.58
<u>Musa</u> spp. var. pelipita-leaves	13.5	22.2		1.58
<u>Panicum maximum</u>	10.7	19.5	54.1	1.95
<u>Pennisetum purpureum</u>	10.9	16.7		1.98

C.P. = Crude protein (%); D.M. = Dry Matter %; IVDDM = In vitro digestibility of dry matter; M.E. = Metabolisable energy (Mcal/kg DM)

Sources: Benavides, 1983b; Franco, 1983; Moreno 1982; Roldan, 1981; Russo, 1983.

Table 4. Grass biomass production (kg/ha) with and without shade of *Erythrina poeppigiana* trees*

	With trees	Without trees
<i>Cynodon plectostachyus</i> <u>1/</u>	5,786 kg/ha/6 months***	2,126 kg/ha/6 months
<i>Pennisetum purpureum</i> <u>2/</u>	6,390 kg/ha/4 months	5,590 kg/ha/4 months

1/ Unpublished data, Bronstein, 1983. Personal communication

2/ Unpublished data, Benavides, 1983a. Personal communication

* Both without fertilizer

** Trees planted 6m x 6m

*** One-year old trees planted 3m x 2m and 3m x 1m

Table 5. Legume trees in pastures in Costa Rica

	Shade	Human food	Living fence	Fire Wood	Wood	Forage
<u>Erythrina poeppigiana</u>						
<u>Schizolobium parahybum</u>						
<u>Inga spp.</u>						
<u>Pithecolobium saman</u>						
<u>Euterolobium cyclocarpum</u>						
<u>Gliricidia sepium</u>						
<u>Diphysa robinoides</u>						
<u>Hymenaea courbaril</u>						
<u>Cassia grandis</u>						
<u>Pithecolobium dulce</u>						
<u>Lonchocarpus spp.</u>						
<u>Albizia adinocephala</u>						
<u>Andira inermis</u>						
<u>Caesalpinia eriotachys</u>						
<u>Platymiscium pinnatum</u>						
<u>Pterocarpus hayesii</u>						

c. Associations with annual crops

Associations of woody legumes with annual crops do not appear to be very common in traditional agricultural systems, the exceptions being leguminous tress in association with various annual crops in the Sahelian zone of Africa (Nair, 1982) and the use of living stakes of Gliricidia sepium as a support for Dioscorea yams in the forest zone of West Africa. Felker (1978) reported increased millet and peanut yeilds near Acacia albida Del., trees in Senegal while increased wheat yields near the same species were reported from the Sudan (Radwanski and Wickens, 1981). Charreau and Vidal (1965) reported higher levels of soil nitrogen, organic matter, Ca, Mg, K, Na, and available P_2O_5 close to Acacia albida trees in Senegal than in soil outside the tree canopy.

While most of the associations of leguminous trees and annual crops occur in the semiarid tropics, with the exception of the Gliricidia-yam association, most efforts at improving the system have been centered in the humid tropics (Steiner, 1982; Kang et al., 1981). The bulk of the research effort with alley cropping has been at the International Institute for Tropical Agriculture, in the forest zone of Nigeria (IITA, 1982, 1981), utilizing Leucaena leucocephala Lam., Tephrosia candida (Roxb) DC, Cajanus Cajan (L) Millsp., and Gliricidia sepium (Jacq) Steud. Recently a program has begun at CATIE in Turrialba, Costa Rica, using Gliricidia sepium and Erythrina poeppigiana (Walpers) O.F. Cook as alley crops (Kass, Russo, and Quinlan, 1983). Although Steiner (1982) saw little prospect for alley cropping in the humid

tropics due to the higher cloud cover, exacerbating effects of shading, efforts in the drier areas have often come up against problems of competition for water and slow growth of the leguminous species under these conditions (IITA, 1981).

We assume that the speakers from IITA will cover their research activities. At CATIE, we have found that in the first year, i.e. of alley crop establishment, total dry matter production of a maize-cassava-bean cropping system was increased by the introduction of both Erythrina and Gliricidia alley crops. Nitrogen recovery of the entire system was increased by 73 and 34 kg ha⁻¹ yr⁻¹ by the introduction of Erythrina and Gliricidia respectively as alley crops.

Another situation where legumes may be used for shade for both annuals and perennials is the home garden. Species found in such gardens in Costa Rica, Haiti, the Philippines and Singapore have been enumerated by Price (1982). While there is no quantitative data on the relative importance of the different species, very few of the trees reported by Price were legumes. Of 38 species of trees found in Costa Rican home gardens, only 3 were leguminous. Of 17 tree species in a Haitian garden, five were legumes. Of 52 tree species in the Singapore gardens, only eight were legumes. It may be in such situations where many fruit trees are grown, and there is considerable input of organic matter from household trash, leguminous trees may not offer too many benefits. It should be remembered, however, that numbers of plants of each species were not reported.

d. Living fenceposts and other non-associative uses

Use of living trees and hedges as fences is a common practice in the tropics (Bond, 1944; Sauer, 1979; Baggio, 1982). Plants are most frequently established from cuttings and can subsequently be pruned back to produce more cuttings, fuelwood, green manure, or fodder for animals in the case of nontoxic species. The comparative advantages and disadvantages of this practice have been summarized by Budowski (1981a). Surveys of the practice in Costa Rica (Sauer, 1979; Baggio, 1982; Lagemann and Heuvelop, 1983) and Nigeria (Bond, 1944) have shown that considerable management skill exists among local farmers which contributes considerably to their success with the practice. Researchers attempting to emulate local practices have often had less success in establishment of such living fences than traditional farmers.

The existing data cannot give a clear indication of whether leguminous trees are favored for living fence posts; but coppicing ability, rapid growth, and ease of husbandry discussed earlier would certainly be of importance in determining the suitability of a particular species for living fences. In the National Academy of Science survey of firewood species (National Academy of Sciences, 1980), where rapid production of biomass suitable for fuel is the major criterion for recommendation, 32 of the 62 recommended species are legumes while another five are of the genera Alnus, Casuarina, and Trema, for which nitrogen fixation has been demonstrated. In Costa Rica, where only eight of the 55 species mentioned by Sauer as fencepost species are legumes, of the three most commonly used genera (Erythrina, Gliricidia, and Bursera), the first two are legumes (Sauer, 1979).

An attribute of many legumes which was of more value before the introduction of barbed wire which favored their use as living fences was thorniness (Sauer, 1979): Acacia albida (Bond, 1944) and Pithecollobium dulce (National Academy of Sciences, 1980), have been cited as valuable for keeping cattle out of undersirable places.

In the provision of shade as an amenity, leguminous trees are often favored in the tropics. In addition to type of shade and ease of management mentioned earlier, the showy flowers and value as a bee forage for honey production are prized in genera such as Calliandra, Erythrina, Gliricidia, and Inga (National Academy of Sciences, 1980).

V. CONCLUSIONS

In the present paper, we have reviewed the use of leguminous trees for shade in four different contexts: with perennial crops, in pastures, with annual crops, and in living fence posts and other non-associative uses. We have noted a tendency for leguminous trees to be favored over non-leguminous trees in all these contexts although the use of nonlegumes for these purposes is often quite widespread. Emphasis was placed on whether legumes offered any particular benefits over nonlegumes as shade trees, which would justify their widespread use and a research effort to increase their use as shade trees. In general, it was found that while leguminous trees have many attributes which favor their use as shade trees, many of these attributes are not exclusive to the legumes. There are nonleguminous trees, which are easy to establish, coppice readily, produce a desirable type of shade, have



showy flowers, and produce abundant litter. Even nitrogen fixation, the factor often cited as a reason for favoring legumes, can be found in non-leguminous genera such as Alnus, Casuarina, and Trema, which are prized as shade trees (National Academy of Sciences, 1980).

The literature was searched with special reference to situations where leguminous and nonleguminous shade species were compared as to their effects on the shaded species and/or the environment. Very few such comparisons have been documented although there are certainly more such comparative studies which have either not been published or published where the authors were unable to consult them. Results of one study with perennial crops (Enriquez, 1983) and two studies with pasture species (Daccarett, 1967; Benavides, 1983a) are given. All of these studies show benefits from using leguminous as opposed to nonleguminous shade species. In the case of a study with annual crops, the shade species were cut back previous to planting maize and cowpeas so the effect of shade would be minimal. The benefits to the crop were greater with a legume (Gliricidia sepium) than a nonlegume (Acioa barteri) but the differences were not statistically significant (IITA, 1982). With living fenceposts, no comparison of leguminous versus nonleguminous species appears to have been documented.

There is obviously a great scope for a larger use of leguminous trees for shade. Foremost among the reasons is the need for reforestation with trees that restore soil fertility while binding the soil with their roots and allowing better water infiltration (Blom, 1980). The greatest need is for reforestation

of marginal lands that have been degraded and compacted and are presently being eroded (National Academy of Sciences, 1980). The use of large stakes or cuttings offers a means of producing a crown in relatively little time; and it has been suggested that trees like Gilricidia could readily be used to eradicate persistent grasses like Imperata by shading them out (Franco, 1983).

THE FUTURE OF LEGUMINOUS SHADE TREES

There is obviously a great scope for a larger use of legumes. But perhaps the greatest scope of legume trees resides in the use in agroforestry systems involving associations with grasses for pasture or animal or perennial crops that benefit from such association by taking advantage of the litter, the better microenvironmental and soil improvement. The fact that many uses can be derived from the large number of leguminous trees to benefit human needs directly or indirectly provides unequalled opportunities for research.

Perhaps the greatest need at this time is information and transfer of knowledge.

The following example of a little known legume tree Calliandra calothyrsus, taken from a forth-coming book of the National Academy of Sciences (3) illustrates the case:

"In 1936, foresters transported seed in this small Central American tree from Guatemala to Indonesia. They were interested in Calliandra and other legumes as possible green manures of shade trees in coffee plantations. In particular, they wanted an alternative to Leucaena, notably for use at high altitudes where Leucaena did not perform well. The foresters planted test plots of Calliandra in a few places in East

(3) National Research Council. Innovations in Tropical Reforestation. Calliandra; a versatile small tree for the humid tropics. Washington, D.C., National Academy Press. 1983. 44p.

Java, but World War II and the subsequent fighting in Indonesia interrupted the investigations, and for 20 years the plant remained largely forgotten by sciences.

The, in the 1960s, administrators of Perum Perhutani, the government forest corporation of Java, noted that villagers in East Java had spontaneously adopted Calliandra and were cultivating it for their firewood needs. The villagers were so successful that in 1974 Perum Perhutani began encouraging the widespread testing and planting of Calliandra. By 1981 the steadily expanding plantations, many planted by villagers themselves, covered almost 2,000 km² on Java. Today Javanese cultivate Calliandra widely, often intercropping it with fruit trees and vegetables. The tree has become so popular in rural areas that "Kaliandra" is now a widely used name for children.

However, Calliandra remains essentially unknown elsewhere, and the purpose of this report is to recount Java's experience in the hope that other countries will be encouraged to investigate Calliandra's promise for themselves".

The report shows that Calliandra fixes nitrogen, is an excellent fuelwood, improves the soil, provides good shade, covering the soil to prevent runoff and erosion. It also produces a forage with 22% protein that can be dried, pelleted, and exported. The flowers produce high quality protein. There are probably dozens of leguminous trees with a future (and probably several dozens with a past) similar to Calliandra.

In Costa Rica, we tested Calliandra for coffee shade, together with Acacia angustissima and the Brazilian "braacatinga" (Mimosa scarella). All these species were readily accepted by the coffee farmers in the San Ramón area. They asked for more seedlings; and our small nursery was not large enough to supply the demand. The irony is that Calliandra had to travel to Indonesia to get recognition in its native Costa Rica.

LITERATURE CITED

- ALLEN, L.H., T.R. SINCLAIR, and E.R. LEMON. 1976. Radiation and microclimate relations in multiple cropping systems. p. 171-200. In Papendick, Sanchez, and Trenbath eds. Multiple Cropping. ASA Spec. Publ. No. 27.
- BAGGIO, A.J. 1982. Establecimiento, manejo, y utilización del sistema agroforestal cercos vivos de Gliricidia sepium (Jacq.) Steud., en Costa Rica. M.S. Thesis. CATIE, Turrialba, Costa Rica. 91p.
- BEER, J. 1980. Alnus acuminata con pasto. CATIE, Turrialba, Costa Rica. 6p. (mimeo).
- BEER, J. 1982. Advantages, disadvantages and desirable characteristics of shade trees for coffee, cacao, and tea. S2-5 In Price, N. ed. Short course: Agroforestry for the humid tropics. Turrialba, Costa Rica, March 16-26. 1982. CATIE mimeo.
- BENAVIDES, J.E. 1983a. Utilización de forrajes de origen arbóreo en la alimentación de ruminantes menores. Turrialba, Costa Rica. CATIE mimeo. 11p.
- BENAVIDES, J.E. 1983b. Alternativas para la producción caprina y bovina en Costa Rica. CATIE mimeo 31p.
- BLOM, P.S. 1980. Leucaena, a promising versatile leguminous tree for the tropics. Abstracts on Tropical Agriculture. 6(3): 9-17.
- BOND, W.E.T. 1944. Hedge plants in Northern Nigeria. Tropical Agriculture (Trinidad). 21(12):228-230.
- BREWBAKER, J.L. and E.M. HUTTON. 1979. Leucaena versatile tropical tree legume. p. 207-259. In G.A. Ritchie ed. New Agricultural Crops. AAAS Selected symposium. Westview Press. Boulder, Colorado.
- BRONSTEIN, G.E. 1982. Los árboles en la producción de pastos. Revisión bibliográfica. CATIE mimeo.
- BRONSTEIN, G.E. 1983. Producción de pasto estrella africana (Cynodon plectostachyus) asociado con árboles de poró (Erythrina poeppigiana), con árboles de laurel (Cordia alliodora) y sin árboles. Proyecto de tesis. CATIE, Turrialba.
- BUDOWSKI, G. 1959. Forestry practices of interest to coffee growers. Coffee (Turrialba, Costa Rica) 1(3):49-52.

- BUDOWSKI, G. 1981a. Agroforestry in Central America. p. 13-21 In Heuvelodp, J. and J. Lagemann eds. Agroforestry, proceedings of a seminar held in CATIE, Turrialba, Costa Rica. 28 Feb. - 3 March, 1981.
- BUDOWSKI, G. 1981b. Applicability of agroforestry systems. p. 13-16 In L.H. MacDonald ed. Agroforestry in the African humid tropics. United Nations University. Tokyo, Japan.
- BUDOWSKI, G. 1981c. Quantification of current agroforestry practices and controlled research plots in Costa Rica. A paper submitted to the consultative meeting on plant research and agroforestry. ICRAF. Nairobi, Kenya. CATIE mimeo 24p.
- BUDOWSKI, G. 1982. The socio-economic effects of forest management on the lives of people living in the area: p. 87-102. The case of Central America and some Caribbean countries. In E.G. Hallsworth ed. Socio-economic effects and constraints in tropical forest management. London John Wiley and Sons Ltd. (Paper presented at workshop held in Dehra Dun, India, December, 1980).
- CADIMA ZEVALLOS, A. and P. de T. ALVIM. 1967. Influencia del árbol de sombra Erythrina glauca sobre algunos factores edafológicos relacionados con la producción del cacaotero. Turrialba, 17(3): 330-336.
- CHARREAU, C. and P. VIDAL. 1965. Influence de l'Acacia albida Del. sur le soil nutrition minerale et rendements des mils Pennisetum au Senegal. Agron. Trop. 20:600-626.
- DACCARET D., M. 1967. La influencia de árboles leguminosos y no leguminosos sobre el forraje que crece bajo ellos. M.S. Thesis. IICA, Turrialba, Costa Rica. 33p.
- DACCARETT D., M. and J. BLYDENSTEIN. 1968. La influencia de árboles leguminosos y no leguminosos sobre el forraje que crece bajo ellos. Turrialba, 18(4): 405-408.
- COOK, O.F. 1901. Shade in coffee culture. U.S. Dept. of Agriculture. Division of Botany. Washington, D.C. 79p.
- DIJKMAN, M.M. 1950. Leucaena- a promising soil-erosion-control plant. Economic Botany 4:337-349.
- ENRIQUEZ, G. 1983. Breve resumen de los resultados del experimento central de plantas perennes de la Montaña. CATIE, Costa Rica. Agroforestry Short Course, 1983. CATIE mimeo. 13p.
- FELKER, P. 1978. State of the art, Acacia albida as a complementary permanent intercrop with annual crops. AID. Univ. of California Riverside, mimeo. 133p.

- FELKER, P. and R.S. BANDURSKI. 1979. Uses and potential uses of leguminous trees for minimal energy input agriculture. *Econ. Bot.* 33: 172-184.
- FONSECA, M.T. 1968. El poró. *Revista de agricultura (Costa Rica)* 40(6-7): 102-112.
- FORDHAM, R. 1971. Irrigation and shade experiment, Las Hermanas A.R. *Cacao res. Univ. W. Indies*, 1971: 38-46.
- FRANCO, F. 1983. Combate de malezas, altura de planta e intensidad del primer corte sobre la productividad y rebrote de Leucaena leucocephala (Lam). DeWit en el período seco. M.S. Thesis. CATIE-UCR. 50p.
- FRANCO, C.M. and R. INFORZATO. 1951. Quantidade de agua transpirada pelo cafeeiro sobreado pelo Ingazeiro. *Bragantia* 11:12-15
- FUENTES, R. 1979. Coffee production farming systems in Mexico. p. 60-72 In G. De las Salas, ed. Workshop, Agro-forestry systems in Latin America. Turrialba, Costa Rica, March 26-30, 1979.
- GREENLAND, D.J. and J.M.L. KOWAL. 1960. Nutrient content of moist tropical forest of Ghana. *Plant and Soil* 12:154-174.
- GRIME, J.P. 1979. Plant strategies and vegetation processes. John Wiley and Sons, London. 222p.
- HARDY, F. 1962. Cacao soils III, the problem of shade for cacao. *Gordian* 62: 685-690.
- HILTON, P.J. 1974. The effect of shade upon the chemical composition of the flush of tea (Camellia sinensis L.) *Trop. Sci.* 16:15-22.
- HUXLEY, P.A. 1970. Some aspects of the physiology of arabica coffee: the central problems and the need for a synthesis. p. 255-267 In Luckwill, L.C. and C.V. Cutting eds., *Physiology of tree crops*. Academic Press.
- INTERNATIONAL INSTITUTE FOR TROPICAL AGRICULTURE (IITA) 1981. Farming systems program. p. 1-49. In Annual Report for 1980.
- _____ 1982. Farming systems program. p. 1-47. In Annual Report for 1981.
- KANG, B.T., G.F. WILSON, and L. SIPKENS. 1981. Alley cropping maize (Zea mays L.L.) and Leucaena leucocephala (Lam) in southern Nigeria. *Plant and Soil* 63: 165-179.
- KASS, D.C.L., R. RUSSO, and M.M. QUINALN. 1983. Leguminous Trees as a nitrogen source for annual crops. *Agronomy Abstracts* p. 45.

- LAGEMANN, J. and J. HEUVELDOP. 1983. Characterization and evaluation of agroforestry systems: the case for Acosta-Puriscal, Costa Rica. *Agroforestry Systems* 1: 101-115.
- LAUDELOUT, H. 1954. Investigation on the mineral element supply of forest-fallow burning. *Proc. 2nd Inter-Afr. Soils Conf.* 1: 383-388.
- MORENO, A.H. 1982. Sistemas agroforestales en la producción pecuaria, ejemplo del piñon cubano (Gliricidia sepium (Jacq) Steud.) Santo Domingo, República Dominicana, S.E.A., Proyecto de Conservación de Forrajes.
- MURRAY, D.B. and R. NICHOLS. 1966. Light, shade, and growth in some tropical plants. *In* Light as an ecological factor (Bainbridge et al, ed.) Blackwell, Oxford p. 249-263.
- NAIR, P.K.R. 1982. Soil productivity aspects of agroforestry. ICRAF. Nairobi, Kenya Mimeo, 83p.
- NATIONAL ACADEMY OF SCIENCES. 1979. Tropical legumes: resources for the future. Washington, D.C. 321p.
- NATIONAL ACADEMY OF SCIENCES. 1980 Firewood crops. Washington, D.C. 237p.
- NORMAN, J.M., E.E. MILLER, and C.B. TANNER. 1971. Light intensity and sunfleck-size distribution in plant canopies. *Agronomy J.* 63:743-748.
- NYE, P.H. 1961. Organic and nutrient cycles under a moist tropical forest. *Plant and Soil.* 13: 333-346.
- ORCHARD, E.R. and G.D. DARB. 1956. Fertility changes under continued wattle culture with special reference to nitrogen fixation and base status of the soil. *IV Int. Congr. Soil Sci.*, Paris IV 45:305-310.
- PAK, N., N. ARYA, R. VILLALON, and M.A. TAGLE.. 1977. Analytical study of tamarugo (Prosopis tamarugo), an autochthonous Chilean feed. *J. Sci. Fd. Agric.* 28:59-62.
- PRICE, N. 1982. The tropical mixed garden, an agroforestry component of the small farm. S2-8 *In* Price, N. ed. Short course: Agroforestry for the humid tropics Turrialba, Costa Rica. March 16-25, 1982. CATIE mimeo.
- PURSEGLOVE, J.W. 1968. Tropical Crops, dicotyledons. Longmans London. 712p.
- RADWANSKI, S.A. and J.E. WICKENS. 1969. The ecology of Acacia albida on mantle soils in Zalingei, Jebel Marra, Sudan. *J. appl. Ecology* 4:569-579.
- ROLDAN PÉREZ, G. 1981. Degradación ruminal de algunos forrajes protéicos en función del consumo de banano verde suplementario. M.S. Thesis. CATIE/UCR. 71p.

- ROSKOSKI, J.P. 1981. Nodulation and N₂ - fixation by Inga jinicuil, a woody legume in coffee plantations. Measurements of nodule biomass and field C₂H₂ reduction rates. Plant and Soil 59:201-206.
- ROSKOSKI, J.P. 1982. Nitrogen fixation in a Mexican coffee plantation. Plant and Soil 67:283-291.
- RUSSO, R.O. 1981. Arboles con pasto, justificación y descripción de un caso en Costa Rica. Turrialba, Costa Rica. CATIE mimeo 12p.
- RUSSO, R.O. 1983. Efecto de la poda de Erythrina poeppigiana (Walpers) O.F. Cook (poró) sobre la nodulación, producción de biomasa y contenido de nitrógeno en el suelo en un sistema agroforestal "Café-poró". M.S. Thesis. CATIE/UCR. 108p.
- SALINAS, H.E. and S.C. SANCHEZ. 1971. Estudio del tamarugo como productor de alimento del ganado lanar en la pampa del tamarugal. Inst. Forestal. Departamento Forestal Sección Silvicultura. Informe Técnico No. 38.
- SANCHEZ, P.A. 1976. Properties and management of soils in the tropics. John Wiley, NY. 618p.
- SAUER, J.D. 1979. Living fences in Costa Rican agriculture. Turrialba 29(4): 255-261.
- STEINER, K.G. 1982. Intercropping in tropical smallholder agriculture with special reference to West Africa. Deutsche Gesellschaft fur Technische Zusammenarbeit (GTZ). Eschborn, 303p.
- WIGHT, W. 1958. The shade tree tradition in the tea gardens of northern India. A.R. Tockai exp. Sta. Indian Tea Assoc p. 75-100.
- WILLEY, R.W. 1975. The use of shade in coffee, cocoa and tea. Horticultural abstracts 45(12):791-798.